

# Effect of Beetroot Juice (Beta Vulgaris L) During Training on Malondialdehyde Level in Athletes

*by* Rosmaini Hsb

---

**Submission date:** 23-Apr-2020 11:30PM (UTC-0400)

**Submission ID:** 1306191811

**File name:** ART20203027.pdf (178.4K)

**Word count:** 3400

**Character count:** 18720

# Effect of Beetroot Juice (Beta Vulgaris L) During Training on Malondialdehyde Level in Athletes

Rosmaini Hasibuan<sup>1</sup>, Fajar Apollo Sinaga<sup>2</sup>, Rika Nailuvar Sinaga<sup>3</sup>

<sup>1</sup>Universitas Negeri Medan, Faculty of Sports Science, Jl. Willem Iskandar, Medan, Indonesia

**Abstract:** Severe physical activity can cause lipid peroxidation marked by increased levels of malondialdehyde (MDA) which can reduce athlete's performance and health. Lipid peroxidation can be prevented or reduced by administering antioxidants. Beets (Beta Vulgaris L) contain various types of natural antioxidants but have not been investigated for their efficacy to reduce lipid peroxidation that is triggered by physical activity. The purpose of this study was to determine the effect of beetroot juice during exercise on lipid peroxidation during maximum physical activity. This type of research is an experimental study with a randomized control group pretest-posttest design research design. The study was conducted at the Unimed Stadium and the USU Faculty of Medicine Integrated Laboratory. The sample was 30 students of Sports Science who met the criteria. Pretest is done by checking MDA levels. Furthermore the sample was divided into 2 groups (P1 = 15, P2 = 15). During the training program group P1 was given 300 ml beet juice while P2 was given as control. Then all samples performed maximum physical activity using a beep test, and MDA levels were re-examined. Then all samples performed maximum physical activity using a beep test, and MDA levels were re-examined. The results showed a decrease in MDA levels in the training group that was given 300ml beet juice compared to the control group ( $p < 0.05$ ). The conclusion of the study was that the administration of beet juice during training could reduce MDA levels when athletes performed maximum physical activity.

**Keywords:** beet juice, malondialdehyde, antioxidant, maximum physical activity

## 1. Introduction

Heavy physical activity can increase oxygen consumption 100-200 times due to an increase in metabolism in the body [1]. Increased use of oxygen mainly by contracting muscles causes an increase in electron leakage from the mitochondria which will become reactive oxygen compounds [2]. Generally 2-5% of oxygen used in metabolic processes in the body will become superoxid ions so that during heavy physical activity an increase in free radical production occurs [3]. When free radical production exceeds cellular defense antioxidants, oxidative stress can occur, where one of the contributing factors is physical activity [4,5]. Under conditions of oxidative stress, free radicals will oxidize lipids and damage the organization of cell membranes [6,7]. Malondialdehyde (MDA) is one of the results of lipid peroxidation induced by free radicals during maximal physical exercise or endurance training with high intensity [8,9,10], so that MDA is a general indicator used to determine the amount of free radicals and indirectly assess the body's oxidant capacity [8]. Bulduk et al reported that volleyball athletes who run a 20 meter shuttle run can cause lipid peroxidation which is marked by an increase in MDA levels and a decrease in CAT and GPx antioxidant levels [11]. The results of research conducted by Eroglu on judo athletes who perform sub-maximal physical activity also concluded that an increase in MDA and a decrease in the concentration of Cu and Zn ions which is an indicator of oxidative stress and lipid peroxidation [12]. The results of a study conducted by Moflehi et al reported that increasing the intensity of exercise can cause increased levels of MDA and creatine kinase levels [13]. It is known, CK is an indicator of damage to muscle cells [14,15]. The decrease in antioxidant levels and the increase in creatine kinase and lipid peroxidation due to maximum and sub-maximal physical activity can be prevented by optimizing nutrition,

especially by increasing antioxidant content [3,16]. Accordingly, Gomez said oxidative damage due to physical activity might be prevented by optimizing nutrition, especially by increasing the antioxidant content of food [17]. According to Silalahi, antioxidant properties will be more effective when consuming antioxidant-rich vegetables or fruits of various types rather than using a single antioxidant such as vitamin E. This may be due to the presence of other components and their interactions in vegetables and fruits that play a positive role [18]. Some natural plants have been tested for antioxidant activity to prevent lipid peroxidation due to physical activity including ginseng extract [19], grape seed extract [20], green tea leaf extract [21], plant extracts of ixora parviflora [22], lemon verbena [23], licopene from tomatoes [24], rhodiola rosea [25], apple juice [26] and red fruit oil [27,28].

One alternative natural ingredient that has antioxidant content is red beet (Beta vulgaris L). Red beet is known to have betalain content which is a compound that has a very high antioxidant that is able to neutralize free radicals [29]. Other compounds in red beets (Beta vulgaris L) that function as antioxidants are betain [30], vitamin C, carotenoids, phenolic acids such as and flavonoids [31,32,33]. Phenolic acid compounds in red beets include ferulic acid, caffeic acid, p-coumaric acid, syringic acid, and vanillic acid [34]. The specialty of red beet (Beta vulgaris L) which has various types of antioxidants makes researchers feel interested and need to examine the antioxidant effect of red beet juice on MDA levels in maximum physical activity.

## 2. Methods

### 2.1 Material

Volume 8 Issue 12, December 2019

[www.ijsr.net](http://www.ijsr.net)

Licensed Under Creative Commons Attribution CC BY

Research materials include Red Beet (*Beta vulgaris* L) obtained from MMTC Medan Market, 1% EDTA solution, Aquadest, MDA Kit purchased from Shanghai Korain Biotech Co., Ltd (Shanghai, China)

## 2.2 Tool

Research tools include: Spectrophotometer (Shimadzu), Glassware, Spuit, Juicer.

## 2.3 Types of Research

This type of research is experimental research with pre-test post-test design, control group design. The location of the study was conducted at the Unimed Stadium, the UNIMED FIK Physiology Laboratory and the Integrated Laboratory of the Faculty of Medicine, University of North Sumatra, Medan.

## 2.4 Research Subject

The study used 30 sports science students with criteria of having a good VO<sub>2</sub>max level, male sex, age 20-22 years, having a good BMI, not smoking, not taking supplements and antioxidants 2 weeks before and during the study, willing to be the subject research and has received ethical clearance from the ethics committee of the Faculty of Medicine, University of North Sumatra.

## 2.5 Research Implementation

All subjects underwent hematological examination to measure MDA levels after the athlete had maximal physical activity by performing a bleptest (pretest). Furthermore, athletes are divided into 2 groups (P1 = 15, P2 = 15). During the training program with moderate intensity for one month, the treatment group (P1) was given 300 ml of beetroot juice (*Beta vulgaris* L) while P2 as a control. Then after one month, all athletes perform maximum physical activity by doing a bleep test. Furthermore, a hematologic examination was performed again to measure MDA levels (posttest).

## 2.6 Malondialdehyde Examination

MDA examination is carried out using a spectrophotometer using the enzyme-linked immunosorbent assay (ELISA) method. Inspection procedures by following the procedures set out in each kit


## 2.7 Data analysis

Data analysis with normality test, homogeneity and t test.

## 3. Result and Discussion

Based on the results of the research, the pretest malondialdehyde (MDA) level in the treatment group was  $1.98 \pm 0.37$  nmol / ml, while the control group was  $2.11 \pm 0.24$  nmol / ml. The normality and homogeneity test shows that the data is normally distributed and homogeneous ( $p > 0.05$ ). The results of the analysis using the independent t test

showed that there were no differences in pretest MDA levels between the treatment group and the control group. The results of measurement of MDA levels posttest treatment group was  $5.75 \pm 0.51$  nmol/ml, while the control group obtained MDA levels  $10.69 \pm 0.97$  nmol/ml. Normality and homogeneity test showed that MDA data posttest treatment group and control group were normally distributed and homogeneous ( $p > 0.05$ ). The results of the analysis using the t dependent test obtained differences in the levels of pretest-posttest MDA treatment groups and control groups ( $p = 0,000$ ). Statistical test results with the independent t test obtained a significant difference in posttest MDA levels between the treatment and control groups ( $p = 0,000$ ).



**Figure 1:** Effect of administration of Beet Juice (*Beta Vulgaris* L) during exercise on malondialdehyde levels on maximal physical activity. Data are mean  $\pm$  SD,  $n = 15$ ,  $p < 0.05$ .

Malondialdehyde is a dialdehyde compound with the molecular formula  $C_3H_4O_2$ , which can be produced from oxidation of unsaturated fatty acids by free radicals. Therefore, high MDA concentrations indicate an oxidation process in the cell membrane [35]. In the study obtained levels of MDA in the group that was not given beet juice was greater than the group given beet juice. High levels of MDA in the group that were not given Beet Juice showed that during maximum physical activity would produce free radicals which oxidize cell membranes (Figure 1). The results of this study are supported by several study results, that acute aerobic physical activity contributes to oxidative stress especially when exercising with high intensity. Two mechanisms that cause oxidative stress in high-intensity aerobic exercise are increased pro-oxidants through the effect of increasing oxygen consumption 10 to 15 times compared to rest and relatively insufficient antioxidants compared to pro-oxidants [36]. Meanwhile according to Ji (1999), during maximum physical activity oxygen consumption throughout the body increases to 20 times, while oxygen consumption in muscle fibers is estimated to increase to 100 times [37]. Under conditions of oxidative stress, free radicals will cause lipid peroxidation of cell membranes and damage the organization of cell membranes [7]. Increased levels of MDA triggered by physical activity have been reported by many researchers including Moflehi, who examined the effects of aerobic exercise with different intensities (low intensity; moderate and high) on increasing levels of MDA and CK in people who are not athletes [13]. The results of this study indicate that the higher the intensity of exercise the greater the MDA level. In this study, aerobic exercise for 20 minutes with an intensity of 80% can increase MDA levels compared to controls ( $9.09 \pm 2.08$  Vs

2.69±1.32μmol/L). Meanwhile, a study conducted by Bulduk, et al., (2011) reported that volleyball athletes who performed a 20 meter shuttle run test with a VO<sub>2</sub>max level of 41.78±4.91ml/kg/min turned out to increase MDA levels from 1.41±0.30 nmol/ml to 2.06±0.08 nmol/ml, while non-athletes with VO<sub>2</sub>max levels of 26.91±3.67 ml/kg/min, MDA levels increased from 0.94±0.24 ml/kg/min to 1.10±0.21 ml/kg/min.

Giving Beet Juice during exercise in this study can reduce MDA levels when athletes perform maximum physical activity when compared with MDA levels in the control group. The decrease in MDA levels is due to the antioxidant content found in beets (*Beta Vulgaris* L). It is known, beets contain antioxidants including phenolic compounds, flavonoids, vitamin C, carotenoids and betalains. Betalains contained in beets are betacyanin and betaxanthine [31,32,38]. Some in vitro research results show betalain pigments can protect cellular components from oxidative injury [39,40]. For example, in the study by Kanner et al two betalain metabolites (betanin and betanidin) were shown to reduce linoleic damage caused by cytochrome C oxidase and lipid membrane oxidation induced by methyoglobin-H<sub>2</sub>O<sub>2</sub>-activated and free iron (AA-Fe). The authors also report that betanin, the most abundant betalain found in beets (300-600 mg / kg), is the most effective inhibitor of lipid peroxidation. High antioxidant activity of Betanin is thought to originate from its extraordinary electron donation capacity and ability to diffuse radicals that very reactive that targets cell membranes [39]. In addition, beets also contain NO<sub>3</sub> compounds (nitrates) which have been shown to suppress the formation of free radicals such as superoxide and hydrogen peroxide [41,42].

#### 4. Conclusion

The conclusion of the study is that the administration of beet juice during training can reduce MDA levels when athletes carry out maximum physical activity.

#### References

- Joyner,M.J dan Casey,D.P. (2015). "Regulation of Increased Blood Flow ( Hyperemia ) to Muscles During Exercise: A Hierarchy of Competing Physiological Needs." 95(2):549-601.
- Nemes,R., Koltai,E., Taylor,A.W., Suzuki,K., Gyori,F dan Radak,Z. (2018). Reactive Oxygen and Nitrogen Species Regulate Key Metabolic, Anabolic, and Catabolic Pathways in Skeletal Muscle. *Antioxidants*. 7, 85; 1-13. doi:10.3390/antiox7070085
- Simioni, Carolina, Giorgio Zauli, Alberto M. Martelli, Marco Vitale, Gianni Sacchetti, Arianna Gonelli, dan Luca M. Neri. (2018). "Oxidative stress: role of physical exercise and antioxidant nutraceuticals in adulthood and aging." *Oncotarget* 9(24):17181-98.
- Daniel, R.M., Stelian, S., Dragomir, C. (2010). The effect of acute physical exercise on the antioxidant status of the skeletal and cardiac muscle in the Wistar rat. *Romanian Biotechnological Letters*. Vol. 15. No. 3, Supplement, p 56-61.
- Kawamura, T dan Muraoka, I. (2018). Exercise-Induced Oxidative Stress and the Effects of Antioxidant Intake from a Physiological Viewpoint. *Antioxidants* Vol: 7;119; doi:10.3390/antiox7090119
- Ayala, A., Munoz,M.F., dan Arguelles, S. (2014). Lipid Peroxidation: Production, Metabolism, and Signaling Mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxidative Medicine and Cellular Longevity*. Article ID 360438;1-31
- Evans, W. J. (2000). Vitamin E, vitamin C, and exercise. *Am J Clin Nutr*, 72, 647S-52S.
- Liu,R., Wu,L., Du,Q., Ren,J.W., Chen,Q.H., Li,D., Mao,R.X., Liu,X.R. Li,Y. (2019). Small Molecule Oligopeptides Isolated from Walnut (*Juglans regia* L.) and Their Anti-Fatigue Effects in Mice. *Molecules*. Vol. 24(45):2-15
- Bao L., Cai X, Wang J, Zhang Y, Sun B, Li Y. (2016). Anti-Fatigue Effects of Small Molecule Oligopeptides Isolated from *Panax ginseng* C. A. Meyer in Mice. *Nutrients*. 13;8;2-11
- Xu, Meihong, Rui Liang, Yong Li, dan Junbo Wang. (2017). "Anti-fatigue effects of dietary nucleotides in mice." *Food & Nutrition Research* 61(1):1334485
- Bulduk, E.O., Ergene, N., Baltaci, A.K., Gumuş, H. (2011). *Plasma antioxidant responses and oxidative stress following a 20 meter shuttle run test in female volleyball players. International Journal of Human Sciences*. Vol.8(2) 510-526
- Eroglu, Y., Daglioglu, O. (2013). The effect of submaximal exercise on oxidant and antioxidant mechanisms in judokas and sedentary. *International Journal of Sport Studies*. Vol., 3 (5), 480- 486
- Moflehi, D., Kok, L.Y., Fadilah, T., Amri, S. (2013). Effect of Single-Session Aerobic Exercise with Varying Intensities on Lipid Peroxidation and Muscle-Damage Markers in Sedentary Males. *Global Journal of Health Science; Vol. 4, No. 4. 48-54*
- Koch, A.J., Pereira,R., Machado,M. (2014). The creatine kinase response to resistance exercise. *J Musculoskeletal Neuronal Interact*. 14(1):68-77
- Figueira,T.G., Magosso,R.F., Neto,J.C., Carli,J.P.C., Pires,C.M.R. (2017). Creatine Kinase Response of Physically Active Young Men to One- and Two-Legged Cycling. *Journal of Exercise Physiologyonline*. Vol20 (3);168-176.
- Pingitore,A., Lima,G.P.P., Mastorci,F., Quinones, A., Iervasi, G. Vassalle,C. (2015). Exercise and oxidative stress: Potential effects of antioxidant dietary strategies in sports." *Nutrition* 31(7-8):916-22
- Gomez-Cabrera MC, Domenech E & Vina J (2008) Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radic Biol Med*. 44, 126-131
- Silalahi, J. (2006). Makanan Fungsional. Penerbit Kanisius Yogyakarta. Halaman 38-56
- Kim, S. H. et al. (2005) 'Effects of Panax ginseng extract on exercise-induced oxidative stress.', *The Journal of sports medicine and physical fitness*, 45(2), pp. 178-82.
- Belviranlı, M., Gokbel, H., Okudan, N. (2012). Basarali,K. Effects of grape seed extract

- supplementation on exercise-induced oxidative stress in rats. *British Journal of Nutrition*. 108, 249–256
- [21] Jowko, E. *et al.* (2015). The effect of green tea extract supplementation on exercise-induced oxidative stress parameters in male sprinters. *European Journal of Nutrition*, 54(5), pp. 783–791.
- [22] Kan, N. W. *et al.* (2013). Hepatoprotective effects of *Ixora parviflora* extract against exhaustive exercise-induced oxidative stress in mice. *Molecules*. 18(9), pp. 10721–10732.
- [23] Funes, L. *et al.* (2011). Effect of lemon verbena supplementation on muscular damage markers, proinflammatory cytokines release and neutrophils' oxidative stress in chronic exercise. *European Journal of Applied Physiology*, 111(4), pp. 695–705.
- [24] Liu, C.C., Huang, C.C., Lin, W.T., Hsieh, C.C., Lin, S.J., Yang, S.C. (2005). Lycopene supplementation attenuated xanthine oxidase and myeloperoxidase activities in skeletal muscle tissues of rats after exhaustive exercise. *British Journal of Nutrition*. (94), pp 595-601
- [25] Parisi, A. *et al.* (2010). Effects of chronic rhodiola rosea supplementation on sport performance and antioxidant capacity in trained male. *Journal of Sports Medicine and Physical Fitness*, 50(1), pp. 57–63.
- [26] Kujawska, M. *et al.* (2011). Cloudy apple juice protects against chemical-induced oxidative stress in rat. *European Journal of Nutrition*, 50(1), pp. 53–60. doi: 10.1007/s00394-010-0114-y.
- [27] Sinaga, F. A., Risfandi, M., Mesanan., Jumadin, J.P. (2017) 'The Effect of Red Fruit Oil on Hematological Parameters and Endurance Performance at the Maximal Physical Activity. *International Journal of Science and Research (IJSR)*. Vol. 6 (5), pp 155-158
- [28] Sinaga, F.A and Susanti, N. (2018). The influence of red fruit (*pandanus conoideus* lam.) Oil on glutathione peroxidase level at maximum physical activity. *Asian J Pharm Clin Res*, Vol 11, Special Issue 1; 103-106
- [29] Clifford, T., Glyn, H., Daniel J., dan Emma J. S. (2015). The Potential Benefits of Red Beetroot Supplementation in Health and Disease. *Nutrients*. 7; 2801–22.
- [30] Zhao, G., He, F., Wu, C., Li, P., Li, N., Deng, J., Zhu, G., Ren, W., dan Peng, Y. (2018). Betaine in inflammation: Mechanistic Aspects and Applications. *Frontiers in Immunology*. 9; 1070; 1-13
- [31] Georgiev, V.G.; Weber, J.; Kneschke, E.M.; Denev, P.N.; Bley, T.; Pavlov, A.I. (2010). Antioxidant activity and phenolic content of betalain extracts from intact plants and hairy root cultures of the red beetroot *Beta vulgaris* cv. Detroit dark red. *Plant Foods Hum. Nutr.* 65, 105–111.
- [32] Kujala, T.S.; Vienola, M.S.; Klika, K.D.; Loponen, J.M.; Pihlaja, K. Betalain and phenolic compositions of four beetroot (*Beta vulgaris*) cultivars. *Eur. Food. Res. Technol.* 2002, 214, 505–510.
- [33] Wootton-Beard, P.C.; Ryan, L (2011). A beetroot juice shot is a significant and convenient source of bioaccessible antioxidants. *J. Funct. Foods* 3, 329–334.
- [34] Chhikara, N.; Kushwaha, K.; Sharma, P.; Gat, Y.; Panghal, A. Bioactive Compounds of Beetroot and Utilization in Food Processing Industry: A Critical Review. *Food Chem.* 2019, 272, 192–200.
- [35] Winarsi, H. (2007). Natural Antioxidants and Free Radicals. Canisius Publisher. Yogyakarta
- [36] Alessio, H.M., Hagerman, A.E., Fulkerson, B.K., Ambrose, J., Rice, R.E., Wiley, R.L. (2000), Generation of reactive oxygen species after exhaustive aerobic and isometric exercise. *Med Sci Sports Exerc.* 32(9):1576-81
- [37] Ji, L.L. (1999). Antioxidants and Oxidative stress in exercise. *Society for Experimental Biology and Medicine*, 283: 292.
- [38] Ninfali, P.; Angelino, D. (2013). Nutritional and functional potential of *Beta vulgaris* *cicla* and *rubra*. *Fitoterapia*. 89: 188-199
- [39] Kanner, J.; Harel, S.; Granit, R. (2001). Betalains A New Class of Dietary Cationized Antioxidants. *J. Agric. Food. Chem.* 49, 5178–5185
- [40] Tesoriere, L.; Fazzari, M.; Angileri, F.; Gentile, C.; Livrea, M.A. In vitro digestion of betalainic foods. (2008). Stability and bioaccessibility of betaxanthins and betacyanins and antioxidative potential of food digesta. *J. Agric. Food. Chem.* 56: 10487-10492
- [41] Lundberg, J.O.; Carlström, M.; Larsen, F.J.; Weitzberg, E. (2011). Roles of dietary inorganic nitrate in cardiovascular health and disease. *Cardiovas Res.* 89: 525–532.
- [42] Wink, D.A.; Miranda, K.M.; Espey, M.G.; Pluta, R.M.; Hewett, S.J.; Colton, C.; Vitek, M.; Feelisch, M.; Grisham, M.B. (2001). Mechanisms of the antioxidant effects of nitric oxide. *Antioxid. Redox Signal.* 3: 203–213

# Effect of Beetroot Juice (Beta Vulgaris L) During Training on Malondialdehyde Level in Athletes

---

## ORIGINALITY REPORT

---

**24%**

SIMILARITY INDEX

**15%**

INTERNET SOURCES

**15%**

PUBLICATIONS

**16%**

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

1%

★ **herbamed.com**

Internet Source

---

Exclude quotes Off

Exclude bibliography Off

Exclude matches < 1%

# Effect of Beetroot Juice (Beta Vulgaris L) During Training on Malondialdehyde Level in Athletes

---

## GRADEMARK REPORT

---

FINAL GRADE

**/75**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---