



Oct 09, 2021
Dr. Diky Setya Diningrat,
Jalan KH Agus Salim No.4,
Kelurahan Madras Hulu
Kecamatan Medan Polonia,
Medan20151,
North Sumatra
Indonesia

Subject: Acceptance Letter for Article No. 89770-JMS-ANSI

It's a great pleasure for us to inform you that below mentioned manuscript has been accepted for publication in Journal of Medical Sciences as Research Article on the recommendation of the reviewers.

Title : Influence of Combination of Weight Training with Sport Massage on 200 Meter Runners
Author's Name : Novita Sari Harahap, Aznan Lelo, Ambrosius Purba and Diky Setya Diningrat
Receiving Date : February 24, 2018
Published on : June 04, 2018

Regards

A blue handwritten signature that appears to read 'Imran Pasha'.

M, Imran Pasha
Publication Manager

UNIVERSITAS NEGERI MEDAN
UNIMED
THE Character Building UNIVERSITY

Final Decision: Accepted After Minor Revision

Reference your article entitled “The Influence of Combination of Weight Training with Sport Massage on Glutathione Peroxides, and Lactic Acid” submitted for publication to Journal of Medical Sciences.

The current article is considered for further evaluation processing after addressing the following queries. My decision is based on the following reason(s):

- Comment 1: Remove the reference from the objective of study within the objective of study.
- Comment 2: Check the running title “Weight training and sports massage collective effect on GP and LA” and do the modifications if required.
- Comment 3: When the table 1 and fig 1 and 2 are representing the similar data then remove one of them as per journal format.
- Comment 4: Provide the reference no 28 after 27 in text and, also mention the reference no 26 citation after 25 references within the text.
- Comment 5: Mention in significance statement that how the current study is beneficial to researchers in future.



Influence of Combination of Weight Training with Sport Massage on Glutathione Peroxides, and Lactic Acid

¹Novita Sari Harahap, ²Aznan Lelo, ³Ambrosius Purba, ⁴Diky Setya Diningrat

¹The Faculty of Sports Science, Medan State University (Universitas Negeri Medan), Medan 20221, North Sumatra, Indonesia,

²Department of Pharmacology, the Faculty of Medicine, University of Sumatera Utara, Medan, North Sumatra, Indonesia

³Department of Physiology, the Faculty of Medicine, Padjajaran University, Bandung, West Java, Indonesia

⁴Department of Biology, the Faculty of Mathematic and Natural Sciences, Medan State University (Universitas Negeri Medan), Medan 20221, North Sumatra, Indonesia,

Background and Objective: Explosive power movement in a 200 meter race will potentially yield free radicals. The energy source of a 200 meter runner, based on his movement, comes from anaerobic predominant metabolism will also potentially yield high lactic acid. The increase in lactic acid causes the decrease in pH which tends to suffer from acidosis so that the establishment of ATP is hampered and cause pain in muscles. This condition can influence the performance of the 200 meter runners. The objective of the research was to find out the influence of combination of weight training with sport massage, on the change in glutathione peroxides and lactic acid elimination in the 200 meter runners. **Materials and Methods:** The research used experimental method with pre and post test design. The research subjects were 20 students of Faculty of Sports Science Medan State University who had not been trained as 200 meter runners and had fulfilled the inclusive and exclusive criteria, taken by using random sampling technique. They were divided into 2 (two) groups: WT+SM (Combination of weight training with sport massage) group and WT-SM (weight training without sport massage). The treatment was performed in 8 weeks with the frequency of exercise 3 times a week. **Results:** The research showed that there was significant influence of the treatment ($p \leq 0.05$) in the pre and post treatment in the WT+SM group and in the WT-SM group on the mean value of increase in glutathione peroxides content and the mean value of decrease in lactic acid content in 200 meter runners. There was no significant disparity ($p \geq 0.05$) of the increase in glutathione peroxides in 200 meter runners in the post treatment done by WT+SM group and by WT-SM group. There was significant disparity ($p \leq 0.5$) in the mean value of decrease in lactic acid content in the 200 meter runners in the post treatment done by WT+SM group and by WT-SM group. **Conclusion:** Combination weight training with sport massage was better in increasing glutathione peroxides content and decreasing lactic acid content in the 200 meter runners than that of only weight training.

Keywords: Weight Training, Sport Massage, Glutathione Peroxides, Lactic Acid, 200 meter Runner

Corresponding Author: Diky Setya Diningrat, Department of Biology, Faculty of Mathematics and Natural Sciences, Medan State University (Universitas Negeri Medan), 20221 Medan, North Sumatra, Indonesia Tel: +6181361362400 LiveDNA: 62.20093

Competing Interest: The authors have declared that no competing interest exists.

Name of the author and e-mail ID	Types of contribution
Dr. Novita Sari Harahap	Dr. Novita Sari Harahap researched on measuring antioxidant glutathione peroxides

	content and Accutrend Plus to measure lactic acid content and also measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure.
Dr. Aznan Lelo	Dr. Ambrosius Purba researched on data analysis on measuring antioxidant glutathione peroxides content and Accutrend Plus to measure lactic acid content
Dr. Ambrosius Purba	Dr. Ambrosius Purba researched on data analysis on measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure.
Dr. Diky Setya Diningrat	Dr. Diky Setya Diningrat was advisor for statistical data analysis and ELISA.

INTRODUCTION

Weight training is a kind of exercise which is known as resistance exercise. The positive effects of this kind of exercise include improvement in physical condition, fitness, and performance; therefore, it is usually recommended because it can decrease the risk for injury, and it functions as prophylaxis, compared with the other exercises^{1,2}. Every physical exercise like weight training has the potency for the incidence of free radicals which can bring about the incidence of oxidative stress. Our body has a number of limited antioxidants, while the activity of free radicals can reach 20% during the training. Therefore, the body has to have its own defense to protect from the attack of free radicals by performing regular exercises. The activity of antioxidant depends upon the intensity and the duration of exercises^{3,4}. Even though exercise can increase antioxidant defense system, a long period and heavy weight training can cause imbalance between free radical production and antioxidants which is called oxidative stress⁵.

The mechanism of the establishment of free radicals in 200 meter runners begins from muscle contraction during exclusive power movement. In the incidence of explosive power, ischemia will occur in muscles and be followed by muscle relaxation alternately. This muscle movement will cause blood flow to be hampered because of muscle contraction, and when the muscles are relaxed blood will flow rapidly. Ischemic reperfusion can cause the release of one electron of respiration chains so that free radicals are established^{6,7,8}. The establishment of free radicals in 200 meter runners can disturb mitochondria and they can even damage mitochondria. The disturbance of mitochondrion function which will cause the establishment of ATP, needed by 200 meter runners, is not fulfilled so that it causes fatigue easily. Therefore, the body has to have defense system; namely, antioxidants in order to protect from the attack of free radicals⁵.

Regular and rated standardized exercises like weight training should be done in order to decrease the negative impact of oxidative stress because of the increase in free radical production. Weight training is intended to increase physical condition of the 200 meter runners. The measurement of antioxidant enzyme content like glutathione peroxides (GPx) should also be done in order to find out whether there is the increase in oxidative stress or not. In the use of energy in the 200 meter runners during the explosive power movement, most of the predominant source of energy is anaerobic laticed which potentially yields high lactic acid⁹. Physiologically, the increase in lactic acid content will decrease pH in body fluid. The decrease

in pH can cause disturbance in the activity of oxidative enzyme in muscle cells so that the capacity of metabolism to produce ATP which is needed during muscle contraction is disturbed^{9,10}.

Recovery should be done immediately after doing the exercises in order to eliminate lactic acid in muscles by conducting sport massage. Immediate recovery after doing exercise is the most crucial factor in increasing an athlete's performance^{11,12}. Researches on the effect of sport massage therapy are relatively limited, and even some of them have different results about the influence of sport massage. For example, in the research conducted by Hemming et al (2000) on 8 amateur boxers; the treatment group was provided by sport massage within 20 minutes by using effleurage and petrissage techniques on the whole body, while the other group was provided with passive-relaxing treatment. The result showed that there was no significant disparity in lactic acid content and the performance of the boxers who were given sport massage treatment from those who were given passive-relaxing treatment¹³.

Another controversial research result was conducted by Wiltshire et al (2005) on untrained individuals; it revealed that sport massage performed during the recovery in decreasing lactic acid content in blood was better than that with passive-relaxing treatment after 200 meter running activity. The two researches above indicate two different results about the influence of Sport Massage on athletes' physical condition. Therefore, this research was conducted because up to the present there has been no research which is concerned with the influence of Sport Massage on the performance of 200 meter runners that is measured with the duration of 200 meter race^{11,14}.

MATERIALS AND METHODS

Subjects: This study was not commercial research but part of doctoral dissertation. This research project was conducted from July 2016 to November 2016 in Pharmacology laboratory Faculty of Medicine North Sumatera University and Sport Science Laboratory of Medan State University. The research subjects were 20 students of Sports Science Study Program, UNIMED, with the criteria of males, 18 to 19 years old, untrained as 200 meter runners, physically healthy, willing to be the research subjects, and active in participating in the whole series of research procedure. They were divided randomly into two groups with 10 participants in the combination of weight training with sport massage (WT+SM) group, and 10 participants in weight training without sport massage (WT-SM) group.

Study Design: The research used experimental study method with Experimental Randomized Pre and Post test group design.

Materials and Device: Materials used in this research were vena blood to analyze GPx content and lactic acid content. The device used in the research was GPx kit to measure antioxidant glutathione peroxides content and Accutrend Plus to measure lactic acid content.

Research Preparation: This research had been approved by the Research Ethics Committee of the Faculty of Medicine, USU, in its Letter No. 270/KOMET/FKUSU/2016 after the research subjects had understood the objective of the research and its benefit. They were then willing to fill out their personal data and informed consent forms and to sign them. The measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure. Schedule for exercises was determined to be in 8 weeks with the frequency of exercises 3 times a week. Exercise was performed in the afternoon, starting from 4:00 PM until 5:00 PM. The research procedure was done by professionals according to their fields. The subjects' blood and lactic acid content were taken by 3 medical analysts. Weight training program was trained by certified trainers who were competent as athletic trainers, while sport

massage was done by 12 certified masseurs. During the process of the research, the subjects were not allowed to consume antioxidant substances like vitamin C and vitamin E or other supplements. They were also disallowed to do other physical activities except training program in the scope of the research. There would be no physical exercise for them, and they had to take enough rest.

Weight Training Procedure: The schedule for training was held in 8 weeks with frequency of exercise 3 times a week in the afternoon. Physical exercise was in the form of weight training plus technical training and sprint running tactics. The weight training was as follows: leg press: 8-12 RM, 3 sets; *squat jump*: 12-15 RM, 3 sets; *leg curl*: 15-25 RM, 3 sets; *sit ups*: 15-25 RM, sets; and Proprioceptive Neuromuscular Facilitation(PNF)exercise was done 3 times a week.

Sport Massage Treatment Procedure: Each time the exercise was done, WT+SM group was given specific sport massage to the 200 meter runners; it was partial massage in the lower legs by using effleurage technique. The method was by doing rubbing movement in the body, using the whole palms and fingers stuck on the lower leg areas; the movement was from the lower parts toward the upper ones. The duration of sport massage was 12 minutes (6 minutes from the right lower part of the legs, and the other 6 minutes from the left lower part of the legs)

Measuring Glutathione Peroxides and Lactic Acid: Pre-Exercise Program. Two days before measuring the Glutathione Peroxides and Lactic Acid, the research subjects were suggested that they not consume any drugs or other supplements; they were not allowed to do any physical exercise, and they had to take enough rest. They began to do warming-up in 15 minutes. After that, they did 200 meter race by recording their travel time. After they got to the finish line, ten minutes later, their blood was taken from vena mediana cubiti for assaying their Glutathione Peroxides and Lactic Acid content. The assaying of Glutathione Peroxides content was based on ELISA method with GPx Qayee-Bio Kit, stated in ng/ml unit, and spectrophotometer UV gauge with wavelength of 450 nm. The assaying of lactic acid content was done by using Accutrend Plus device with milimoles per liter (mmol/L) unit. By the end of the training program, the measurement of Glutathione Peroxides and Lactic Acid was the same as what had been done in sport massage (WT-SM).

Statistical analysis: All data were statistically analyzed with SPSS software (version 16). One-way analysis of variance (ANOVA) was used to study significant difference between means and significance level at $p = 0.05$.

RESULTS

The research subjects in the two groups, based on age were 18.65 years old, based on body weight were 61.65, based on height were 167.8, based on IMT were 22 based on systole blood pressure were 119.5, based on diastole blood pressure were 78., and based on pulse while in rest were 64.8. All in all, they showed no significant disparity ($p \geq 0.05$); therefore, all of them were homogeneous; they had the same characteristics, either physically or physiologically.

Table 1. Mean value of glutathione peroxides and lactic acid content in the 200 m runners

Variables		WT+SM	WT-SM	p
		Mean±SD	Mean±SD	
Glutathione Peroxides	Pre-exercise	218.35± 10.31	220.17 ±12.66	0.155
	Post-exercise	248.57 ± 16.59	232.82± 23.38	
	t-dependent	0.000*	0.01*	
Lactic Acid	Pre-exercise	8.23 ±0.27	8.23 ± 0.46	0.000*

Post-exercise	6.34 ± 0.31	7.82 ± 0.21
t-dependent	0.000*	0.02*

Explanation: * = significance ($p \leq 0,05$), SD : standard deviation.

Glutathione peroxides of the 200 meter runners increased significantly ($p \leq 0.05$) in the two groups in the pre-exercise, compared with that in the post-exercise; in the WT±SM group and in the WT-SM group (Table 1). The increase in glutathione peroxides in the 200 meter runners in the post-exercise in WT±SM group was higher than that WT-SM group. The percentage of the increase in glutathione peroxides content of the 200 meter runners. However, there was no significant disparity ($p \geq 0.05$) in the increase in glutathione peroxides content in the 200 meter runners in the post-exercise between WT+SM group and WT-SM group (Fig. 1).




Fig. 1: Mean value of glutathione peroxides content in pre and post exercise in the 200 m runners.

Lactic acid content in the 200 meter runners decreased significantly ($p \leq 0.05$) in the two groups in pre-exercise, compared with that in the post-exercise: in the WT+SM group and the WT-SM (Table 1). The decrease in lactic acid content in the 200 meter runners in the post-exercise in the WT+SM group was higher than that in the WT-SM group at the percentage of the decrease in the lactic acid content in the 200 meter runners. There was significant disparity ($p \leq 0.05$) in the decrease of lactic acid content in the 200 meter runners between the WT+SM group and the WT-SM group (Fig. 2).




Fig.2: Mean value of lactic acid content in the pre and post-exercise in the 200 m runners. *= significance ($p < 0,05$).

DISCUSSION

Disparity in glutathione peroxides content in the 200 meter runners: Weight training is a type of sport which is intended to develop strength which uses gravitation against the gravity which is yielded by muscles through concentric or eccentric contraction. In this exercise, body muscles undergo contraction by using body weight or other devices to stimulate muscle growth/work strength, and resistance and by targeting certain muscle groups and types of movement. The exercise program can give benefit as maximal as possible and the risk as minimal as possible if it fulfills the training formulation which is called FITT (Frequency of Training, Intensity of Training, Time Spent, Type of Exercise)¹⁵.

Standardized exercise which is measured, guided, and programmed well can increase the physical condition of the 200 meter runners so that free radicals which exist as the result of the high intensity of physical exercise can be eliminated by the runners who have good physical condition. If they have good physical condition, their capability of producing endogenous antioxidants like Glutathione Peroxides will be better than the 200 meter runners who do not have good physical condition⁴. Weight training program is more focused on anaerobic exercise so that it will potentially yield relatively more lactic acid and can cause purine catabolism to become xanthenes and veins. This indicates the transition and acute vein deoxygenating which has the same problem with ischemic reperfusion. Purine catabolism and ischemic reperfusion will activate xanthenes oxides system which, in turn, establishes free radicals (O_2)^{4,16,17,18}. The research which was done by Best, et. al. (2002) on 100 meter swimmers revealed that there was the increase in oxidative stress biomarker like TBARS (4.1 ± 0.7 versus 4.9 ± 1.1), *Creatinine Kinase* (206 ± 170.7 versus 244 ± 176.9) and glutathione peroxides (0.52 ± 0.06 versus 0.62 ± 0.05). This research concluded that anaerobic exercise could cause oxidative stress; this increase in oxidative stress could hamper the establishment of Glutathione Peroxides and increased the use of endogenous antioxidants like Glutathione peroxides¹⁹.

The result of this research was in accordance with the research done by Groussard, et. al. (2003); it was aimed to determine the effect of implementing the combination of exercise with massage regularly on oxidant and antioxidant content. In their research, the research subjects were 25 people who did not do their sport regularly, and they did not receive any sport massage. They were randomly divided into three groups: control group (CG, $n=9$), exercise group (EG, $n=8$), and massage and exercise group (MEG, $n=8$). The result of the research showed that there was significant decrease ($p < 0.05$) in MDA content in the EG and MEG groups. It was also found that there was significant increase ($p < 0.05$) in glutathione peroxides in the EG and MEG groups, and there was significant increase ($p < 0.05$) in SOD content in the EG and MEG groups, compared with that in CG group. This indicated that physical activity which is done regularly and combined with massage can cause positive effect on the increase in antioxidants and endogen – GPx and SOD²⁰.

The result of this research was also in accordance with another research done by Azizbeigi et al. (2013); it was aimed to find out the influence of progressive resistance exercise on oxidative weight and the activity of enzymatic antioxidant in erythrocyte¹⁴. The research subjects were 20 men who were divided into two groups: progressive resistance exercise group with moderate intensity (50%) within 8 weeks and control group. The result of the research showed that progressive resistance exercise with moderate intensity significantly increased the activity of SOD erythrocyte ($p=0.014$), decreased MDA concentration ($p=0.030$), and GPx content tended to increase. The role of sport massage in the WT+SM group could give positive effect on the increase in GPx content even though statistically there was no significant disparity from WT-SM group, but there was the disparity in the increase in WT+SM group.

Disparity in lactic acid content in the 200 m runners: Exercise in WT+SM group was more significant in decreasing lactic acid in the 200 meter runners than that in the WT-SM group. This indicated that LPK and sport massage had significant influence on the decrease in lactic acid content in the 200 meter runners. The increase in lactic acid content is more significant in the WT+SM because weight training with standardized exercise which is measured, regular, and programmed well can increase tolerance against lactic acid and increase the capacity of bicarbonate buffer and phosphate in muscles. Besides that, sport massage can be efficient to eliminate lactic acid in the 200 meter runners²¹.

200 meter race athletics is the category of sprint and most of its predominant energy source is lactacyd anaerobic lactacyd which will yield relatively high lactic acid. The result of measuring lactic acid in the pre-exercise program in all treatment groups indicated the average increase of 8.23 mmol/L measured 10 minutes in the post 200 meter race. The increase in lactic acid was caused by the incidence of hypoxia in muscle tissues so that anaerobic metabolism which yields lactic acid occurs. The increase in lactic acid is in accordance with the result of the research done by Kawczyński et al., (2014) which found that there was the increase in lactic acid content in the 100 meter runners at the mean value of 9.51 mmol/L which was measured in the 10th minute in the post 100 meter race²². The higher the physical activity is, the more increasing the need for oxygen. The need for oxygen can be increased by increasing the function of heart and lungs. When physical activity is high while oxygen intake from heart and lungs is not optimal, anaerobic metabolism exist to fulfill the need for energy. This condition increases lactic acid content in blood and muscles. The increase in lactic acid can change reactive oxygen compound to be more reactive so that the physical condition of the 200 meter runners will not maximal¹⁹.

Physiologically, the increase in lactic acid content will decrease pH in the body liquid. The decrease in pH will affect the establishment of ATP which is needed during muscle contraction. Muscle contraction occurs because nervous impulse will reach axon terminal, followed by the release of acetylcholine (Ach) which immediately potentially stimulates muscle action in the membrane of muscles sarcolemma. Potential action in muscles spreads through tubules T and causes reticulum sarcoplasm to release calcium ion to sarcoplasm. When there was the increase in lactic acid, the releasing process of acetylcholine and calcium ion from reticulum sarcoplasm will be disturbed. In consequence, there is no calcium bond with troponin. This bond will open tropomyosin which covers cross-bridge receptor. The opening of cross-bridge receptor from tropomyosin bond causes the actin to shift and the caput myosin to be stuck; in this process, ATP is needed²³. The increase in lactic acid will disturb the ATP splitting metabolism to be the energy.

Calcium ion low content in cytosol will decrease the muscle contraction power (leg muscle strength, abdominal and leg muscle resistance, leg muscle explosive power, running speed, and reaction speed). During the resting period, calcium ion enters again into reticulum sarcoplasm, and tropomyosin will close cross-bridge receptor again so the bond of actins and myosin is released and contraction will not occur. Physical activity will cause the bond of actins and myosin to become smooth so that muscle contraction highly depends on the higher or lower content of lactic acid in the 3200 meter runners²³.

Sport massage given in the pre-exercise to the WT+SM group in the 200 meter runners was an active recovery. It can cause muscles to be relaxed and regain the condition of blood vessel which used to be pinched by muscle contraction. Optimal relaxation can increase the backflow of vena blood in the lower muscle part of legs in bringing optimally lactic acid to the hearts of the 200 meter runners; therefore, it could increase the oxygen of muscle tissues and

expedite the elimination of lactic acid²⁴. The increase in tissue oxygenation will increase the conversion of lactic acid in heart through the Cori cycle to pyruvate acid which, in turn, will be changed to glucose so that it will expedite the supply for energy and expedite recovery⁹.

Physiologically, massage is proved to be able to decrease heartbeat, to increase blood pressure, to increase blood circulation and lymph, to decrease muscle tension, to increase the range of joint movement, and to decrease pain. Physiology is widely used by athletes, either for their physical performance or for prevention, therapy, and rehabilitation for injuries and the side effect of sport^{19,25}. Sport massage on the body will give physiological effect such as increasing blood flow and lymphatic flow, stimulating nervous system, and increasing vena blood backflow. The increase in this vena blood backflow will efficiently return blood to the heart and drain off lactic acid which is deposited in muscles, and expedite the elimination of lactic acid in blood and muscles. It has been proved that effleurage technique is more effective in decreasing lactic acid deposit in muscle cells than that with moderate intensity sport^{24,26}.

This glucose can be reused as metabolism in active muscles or kept in muscles through the process of gluconeogenesis to become glycogen and can be used as an energy source. Pyruvate acid will enter mitochondria when the exercise is done below the lactate threshold (LT) and the establishment of energy is processed anaerobically, while in the exercise which is done over the lactate threshold (LT) like conventional exercise (heavy intensity exercise), pyruvate acid will be changed to lactic acid because the supply of oxygen is not enough to enter mitochondria²⁷. In a research done by Groussard, et. al., (2003) by using Wingate test, the research subjects pedaled bicycles within 30 seconds. Wingate test would stimulate anaerobic glycolysis metabolism by activating purine catabolism and lactic acid production. The data showed that anaerobic exercise by using Wingate test is the potential factor which causes oxidative stress. It is relatively the same as its movement pattern to increase anaerobic glycolysis in the 200 meter runners²⁰.

According to Best et al., (2008), an athlete can carry on his exercise in relatively high intensity in the condition of lactic acid of 6-7 mmol/L¹⁹. Therefore, the research of this research with the mean value of lactic acid content in the 200 meter runners in the LPK+SM of 6.34 mmol/L showed that the level of fatigue was still low so the performance in the WT+SM group was better than that in the WT-SM group. The research done by Weerapong et al (2005) on individuals who were not trained found that sport massage during the recovery period could decrease lactic acid content in blood more than passive rest after the activity in 200 meter race²⁵. The research done by Monedero and Donne (2000) also found that sport massage could increase the elimination of lactic acid during the recovery period after doing high intensity exercise²⁸.

Pyne, et al., (2005) pointed out that sport massage done within 20 minutes could cause the decrease in lactic acid content in blood of 36.21%, and sport massage done within 45 minutes could decrease lactic acid in blood of 72.4%²⁷. The research done by Hanon, et al., (2010) was aimed to evaluate the influence of three methods for recovery – massage, active and passive recovery on the change in lactic acid concentration in blood, and the time of performance in swimming athletes²⁹. The research subjects were 17 swimming athletes. The result of the research showed that there was no disparity between active recovery and massage, and massage was more effective than passive recovery to eliminate lactic acid in blood. The research result was also in accordance with the research done by Devlin, et al., (2014)³⁰. It stated that active recovery after heavy exercise would eliminate more quickly the accumulation of lactic acid in blood than passive recovery.

The explanation above indicates that in regular physical exercise and measured standardized exercise the body will be adjusted so that the capacity to work in higher intensity

will be able to use lactic acid during the exercise^{31,32}. This will cause more lactic acid to enter mitochondria and to be used for aerobic metabolism so that it can decrease the amount of lactic acid released to the blood flow. The combination with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage.

CONCLUSION

Combination of weight training with sport massage was better in increasing glutathione peroxides and decreasing lactic acid content in the 200 meter runners than weight training only.

SIGNIFICANCE STATEMENTS

This research showed that there is significant impact of combination of weight training with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage. Further studies are needed to investigate the correlation between weight training with sport massage and weight training without sport massage.

ACKNOWLEDGMENT

The authors would like to thank Ministry of research and higher education Republic Indonesia for providing Research Grants (Doctoral Research Grant, grant no. 038A/UN33.8/LL/2017.). The authors would also like to thank Medan State University (Universitas Negeri Medan) and Faculty of Medicine, North Sumatra University for providing the necessary facilities.

REFERENCES

- [1] Westcott, W., 2009. ACSM strength training guidelines: Role in body composition and health enhancement. *ACSM's Health & Fitness Journal*, 13(4), pp.14-22.
- [2] Novatchkov, H. and Baca, A., 2013. Artificial intelligence in sports on the example of weight training. *Journal of sports science & medicine*, 12(1), p.27.
- [3] Gomez-Cabrera, M.C., Domenech, E. and Viña, J., 2008. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radical Biology and Medicine*, 44(2), pp.126-131.
- [4] El Abed, K., Rebai, H., Bloomer, R.J., Trabelsi, K., Masmoudi, L., Zbidi, A., Sahnoun, Z., Hakim, A. and Tabka, Z., 2011. Antioxidant status and oxidative stress at rest and in response to acute exercise in judokas and sedentary men. *The Journal of Strength & Conditioning Research*, 25(9), pp.2400-2409.
- [5] Escribano, B.M., Tunez, I., Requena, F., Rubio, M.D., De Miguel, R., Montilla, P., Tovar, P. and Aguera, E.I., 2010. Effects of an aerobic training program on oxidative stress biomarkers in bulls. *Veterinari Medicina*, 55(9), pp.422-8.
- [6] Kostaropoulos, I.A., Nikolaidis, M.G., Jamurtas, A.Z. and Ikonomidou, G.V., 2006. Comparison of the blood redox status between long-distance and short-distance runners. *Physiological research*, 55(6), p.611.
- [7] Güzel, N.A., Hazar, S. and Erbas, D., 2007. Effects of different resistance exercise protocols on nitric oxide, lipid peroxidation and creatine kinase activity in sedentary males. *Journal of sports science & medicine*, 6(4), p.417.
- [8] Sahlin, K., Shabalina, I.G., Mattsson, C.M., Bakkman, L., Fernström, M., Rozhdestvenskaya, Z., Enqvist, J.K., Nedergaard, J., Ekblom, B. and Tonkonogi, M., 2010. Ultraendurance exercise increases the production of reactive oxygen species in

- isolated mitochondria from human skeletal muscle. *Journal of applied physiology*, 108(4), pp.780-787.
- [9] Astrand, P.O., Rodahl, K. (2003) *Textbook of Work Physiology, Physiological Base of Exercise*. New York: McGraw-Hill.
- [10] Hall, J.E., 2015. *Guyton and Hall textbook of medical physiology e-Book*. Elsevier Health Sciences.
- [11] Wiltshire, E.V., Poitras, V., Pak, M., Hong, T., Rayner, J. and Tschakovsky, M.E., 2010. Massage impairs postexercise muscle blood flow and "lactic acid" removal. *Medicine & Science in Sports & Exercise*, 42(6), pp.1062-1071.
- [12] Pinar, S., Kaya, F., Bicer, B., Erzeybek, M.S. and Cotuk, H.B., 2012. Different recovery methods and muscle performance after exhausting exercise: comparison of the effects of electrical muscle stimulation and massage. *Biology of sport*, 29(4), p.269.
- [13] Hemmings, B., Smith, M., Graydon, J. and Dyson, R., 2000. Effects of massage on physiological restoration, perceived recovery, and repeated sports performance. *British journal of sports medicine*, 34(2), pp.109-114.
- [14] Azizbeigi, K., Azarbayjani, M.A., Peeri, M., Agha-Alinejad, H. and Stannard, S., 2013. The effect of progressive resistance training on oxidative stress and antioxidant enzyme activity in erythrocytes in untrained men. *International journal of sport nutrition and exercise metabolism*, 23(3), pp.230-238.
- [15] Loturco, I., Ugrinowitsch, C., Roschel, H., Tricoli, V. and González-Badillo, J.J., 2013. Training at the optimum power zone produces similar performance improvements to traditional strength training. *Journal of sports science & medicine*, 12(1), p.109.
- [16] Valado, A., Pereira, L. and Paula, C., 2007. Effect of the intense anaerobic exercise on nitric oxide and malondialdehyde in studies of oxidative stress. *International Journal of Biology and Biomedical Engineering*, 1(1), pp.32-26.
- [17] Fisher-Wellman, K. and Bloomer, R.J., 2009. Acute exercise and oxidative stress: a 30 year history. *Dynamic medicine*, 8(1), p.1.
- [18] George, B.O. and Osharechiren, O.I., 2009. Oxidative stress and antioxidant status in sportsmen two hours after strenuous exercise and in sedentary control subjects. *African Journal of Biotechnology*, 8(3).
- [19] Best, T.M., Hunter, R., Wilcox, A. and Haq, F., 2008. Effectiveness of sports massage for recovery of skeletal muscle from strenuous exercise. *Clinical Journal of Sport Medicine*, 18(5), pp.446-460.
- [20] Groussard, C., Rannou-Bekono, F., Machefer, G., Chevanne, M., Vincent, S., Sergent, O., Cillard, J. and Gratas-Delamarche, A., 2003. Changes in blood lipid peroxidation markers and antioxidants after a single sprint anaerobic exercise. *European journal of applied physiology*, 89(1), pp.14-20.
- [21] Mika, A., Mika, P., Fernhall, B. and Unnithan, V.B., 2007. Comparison of recovery strategies on muscle performance after fatiguing exercise. *American journal of physical medicine & rehabilitation*, 86(6), pp.474-481.
- [22] Kawczyński, A., Kobiałka, K., Dariusz, M., Paweł, C., Adam, M., Adam, Z. and Jan, C., 2015. Blood lactate concentrations in elite Polish 100 m sprinters. *International Journal of Performance Analysis in Sport*, 15(1), pp.391-396.
- [23] McArdle, W.D., Katch, F.I. and Katch, V.L., 2010. *Exercise physiology: nutrition, energy, and human performance*. Lippincott Williams & Wilkins.
- [24] Cafarelli, E. and Flint, F., 1992. The role of massage in preparation for and recovery from exercise. *Sports medicine*, 14(1), pp.1-9.

- [25] Weerapong, P., Hume, P.A. and Kolt, G.S., 2005. The mechanisms of massage and effects on performance, muscle recovery and injury prevention. *Sports medicine*, 35(3), pp.235-256.
- [26] Goats, G.C., 1994. Massage--the scientific basis of an ancient art: Part 2. Physiological and therapeutic effects. *British journal of sports medicine*, 28(3), pp.153-156.
- [27] Pyne, D.B., Lee, H.A.M.I.L.T.O.N. and Swanwick, K.M., 2001. Monitoring the lactate threshold in world-ranked swimmers. *Medicine and Science in Sports and Exercise*, 33(2), pp.291-297.
- [28] Monedero, J. and Donne, B., 2000. Effect of recovery interventions on lactate removal and subsequent performance. *International journal of sports medicine*, 21(08), pp.593-597.
- [29] Hanon, C., Lepretre, P.M., Bishop, D. and Thomas, C., 2010. Oxygen uptake and blood metabolic responses to a 400-m run. *European journal of applied physiology*, 109(2), pp.233-240.
- [30] Devlin, J., Paton, B., Poole, L., Sun, W., Ferguson, C., Wilson, J. and Kemi, O.J., 2014. Blood lactate clearance after maximal exercise depends on active recovery intensity. *Journal of Sports Medicine and Physical Fitness*, 54(3), pp.271-278.
- [31] Leprêtre, P.M., Lopes, P., Thomas, C. and Hanon, C., 2012. Changes in cardiac tone regulation with fatigue after supra-maximal running exercise. *The Scientific World Journal*, 2012.
- [32] Ali, S.R., Koushkie, M.J., Asadmanesh, A. and Salesi, M., 2012. Influence of massage, active and passive recovery on swimming performance and blood lactate. *The Journal of sports medicine and physical fitness*, 52(2), pp.122-127.



Final Decision: Accepted After Minor Revision

Reference your article entitled “The Influence of Combination of Weight Training with Sport Massage on Glutathione Peroxides, and Lactic Acid” submitted for publication to Journal of Medical Sciences.

The current article is considered for further evaluation processing after addressing the following queries. My decision is based on the following reason(s):

Comment 1: Remove the reference from the objective of study within the objective of study.

The two researches above indicate two different results about the influence of Sport Massage on athletes’ physical condition. Therefore, this research was conducted because up to the present there has been no research which is concerned with the influence of Sport Massage on the performance of 200 meter runners that is measured with the duration of 200 meter race.

- I have fixed it and attached in the article

Comment 2: Check the running title “Weight training and sports massage collective effect on GP and LA” and do the modifications if required.

Influence of Combination of Weight Training with Sport Massage on 200 meter Runners

- I have fixed it and attached in the article

Comment 3: When the table 1 and fig 1and 2 are representing the similar data then remove one of them as per journal format.

The research subjects in the two groups, based on age were 18.65 years old, based on body weight were 61.65, based on height were 167.8, based on IMT were 22 based on systole blood pressure were 119.5, based on diastole blood pressure were 78., and based on pulse while in rest were 64.8. All in all, they showed no significant disparity ($p \geq 0.05$); therefore, all of them were homogeneous; they had the same characteristics, either physically or physiologically.

Table 1. Mean value of glutathione peroxides and lactic acid content in the 200 m runners

Variables		WT+SM	WT-SM	p
		Mean±SD	Mean±SD	
Glutathione Peroxides	Pre-exercise	218.35± 10.31	220.17 ±12.66	0.155
	Post-exercise	248.57 ± 16.59	232.82± 23.38	
	t-dependent	0.000*	0.01*	
Lactic Acid	Pre-exercise	8.23 ±0.27	8.23 ± 0.46	0.000*
	Post-exercise	6.34 ± 0.31	7.82 ± 0.21	
	t-dependent	0.000*	0.02*	

Explanation: * = significance ($p \leq 0.05$), SD : standard deviation.

Glutathione peroxides of the 200 meter runners increased significantly ($p \leq 0.05$) in the two groups in the pre-exercise, compared with that in the post-exercise; in the WT±SM group and in the WT-SM group (Table 1). The increase in glutathione peroxides in the 200 meter runners in the post-exercise in WT±SM group was higher than that WT-SM group. The

percentage of the increase in glutathione peroxides content of the 200 meter runners. However, there was no significant disparity ($p \geq 0.05$) in the increase in glutathione peroxides content in the 200 meter runners in the post-exercise between WT+SM group and WT-SM group (Table 1).

Lactic acid content in the 200 meter runners decreased significantly ($p \leq 0.05$) in the two groups in pre-exercise, compared with that in the post-exercise: in the WT+SM group was and the WT-SM (Table 1). The decrease in lactic acid content in the 200 meter runners in the post-exercise in the WT+SM group was higher than that in the WT-SM group at the percentage of the decrease in the lactic acid content in the 200 meter runners. There was significant disparity ($p \leq 0.05$) in the decrease of lactic acid content in the 200 meter runners between the WT+SM group and the WT-SM group (Table 1).

- I have fixed it and attached in the article

Comment 4: Provide the reference no 28 after 27 in text and, also mention the reference no 26 citation after 25 references within the text.

- I have fixed it and attached in the article

Comment 5: Mention in significance statement that how the current study is beneficial to researchers in future.

SIGNIFICANCE STATEMENTS

This research showed that there is significant impact of combination of weight training with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage. This research provide scientific proof of the features of the combination of weight training with sport massage to improve the 200m runner's performance. Future research that needs to be done is to know the most effective type of weight training and type of sport massage to eliminate lactic acid and increase glutathione peroxide for a successful 200m runner achievement.

- I have fixed it and attached in the article





THE
Character Building
UNIVERSITY

Final Decision: Accepted After Minor Revision

Reference your article entitled “The Influence of Combination of Weight Training with Sport Massage on Glutathione Peroxides, and Lactic Acid” submitted for publication to Journal of Medical Sciences.

The current article is considered for further evaluation processing after addressing the following queries. My decision is based on the following reason(s):

Comment 1: Remove the reference from the objective of study within the objective of study.

The two researches above indicate two different results about the influence of Sport Massage on athletes’ physical condition. Therefore, this research was conducted because up to the present there has been no research which is concerned with the influence of Sport Massage on the performance of 200 meter runners that is measured with the duration of 200 meter race.

- I have fixed it and attached in the article

Comment 2: Check the running title “Weight training and sports massage collective effect on GP and LA” and do the modifications if required.

Influence of Combination of Weight Training with Sport Massage on 200 meter Runners

- I have fixed it and attached in the article

Comment 3: When the table 1 and fig 1and 2 are representing the similar data then remove one of them as per journal format.

The research subjects in the two groups, based on age were 18.65 years old, based on body weight were 61.65, based on height were 167.8, based on IMT were 22 based on systole blood pressure were 119.5, based on diastole blood pressure were 78., and based on pulse while in rest were 64.8. All in all, they showed no significant disparity ($p \geq 0.05$); therefore, all of them were homogeneous; they had the same characteristics, either physically or physiologically.

Table 1. Mean value of glutathione peroxides and lactic acid content in the 200 m runners

Variables		WT+SM	WT-SM	p
		Mean±SD	Mean±SD	
Glutathione Peroxides	Pre-exercise	218.35± 10.31	220.17 ±12.66	0.155
	Post-exercise	248.57 ± 16.59	232.82± 23.38	
	t-dependent	0.000*	0.01*	
Lactic Acid	Pre-exercise	8.23 ±0.27	8.23 ± 0.46	0.000*
	Post-exercise	6.34 ± 0.31	7.82 ± 0.21	
	t-dependent	0.000*	0.02*	

Explanation: * = significance ($p \leq 0.05$), SD : standard deviation.

Glutathione peroxides of the 200 meter runners increased significantly ($p \leq 0.05$) in the two groups in the pre-exercise, compared with that in the post-exercise; in the WT±SM group and in the WT-SM group (Table 1). The increase in glutathione peroxides in the 200 meter runners in the post-exercise in WT±SM group was higher than that WT-SM group. The

percentage of the increase in glutathione peroxides content of the 200 meter runners. However, there was no significant disparity ($p \geq 0.05$) in the increase in glutathione peroxides content in the 200 meter runners in the post-exercise between WT+SM group and WT-SM group (Table 1).

Lactic acid content in the 200 meter runners decreased significantly ($p \leq 0.05$) in the two groups in pre-exercise, compared with that in the post-exercise: in the WT+SM group was and the WT-SM (Table 1). The decrease in lactic acid content in the 200 meter runners in the post-exercise in the WT+SM group was higher than that in the WT-SM group at the percentage of the decrease in the lactic acid content in the 200 meter runners. There was significant disparity ($p \leq 0.05$) in the decrease of lactic acid content in the 200 meter runners between the WT+SM group and the WT-SM group (Table 1).

- I have fixed it and attached in the article

Comment 4: Provide the reference no 28 after 27 in text and, also mention the reference no 26 citation after 25 references within the text.

- I have fixed it and attached in the article

Comment 5: Mention in significance statement that how the current study is beneficial to researchers in future.

SIGNIFICANCE STATEMENTS

This research showed that there is significant impact of combination of weight training with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage. This research provide scientific proof of the features of the combination of weight training with sport massage to improve the 200m runner's performance. Future research that needs to be done is to know the most effective type of weight training and type of sport massage to eliminate lactic acid and increase glutathione peroxide for a successful 200m runner achievement.

- I have fixed it and attached in the article



Influence of Combination of Weight Training with Sport Massage on 200 meter Runners

¹Novita Sari Harahap, ²Aznan Lelo, ³Ambrosius Purba, ⁴Diky Setya Diningrat

¹The Faculty of Sports Science, Medan State University (Universitas Negeri Medan), Medan 20221, North Sumatra, Indonesia,

²Department of Pharmacology, the Faculty of Medicine, University of Sumatera Utara, Medan, North Sumatra, Indonesia

³Department of Physiology, the Faculty of Medicine, Padjajaran University, Bandung, West Java, Indonesia

⁴Department of Biology, the Faculty of Mathematic and Natural Sciences, Medan State University (Universitas Negeri Medan), Medan 20221, North Sumatra, Indonesia,

Background and Objective: Explosive power movement in a 200 meter race will potentially yield free radicals. The energy source of a 200 meter runner, based on his movement, comes from anaerobic predominant metabolism will also potentially yield high lactic acid. The increase in lactic acid causes the decrease in pH which tends to suffer from acidosis so that the establishment of ATP is hampered and cause pain in muscles. This condition can influence the performance of the 200 meter runners. The objective of the research was to find out the influence of combination of weight training with sport massage, on the change in glutathione peroxides and lactic acid elimination in the 200 meter runners. **Materials and Methods:** The research used experimental method with pre and post test design. The research subjects were 20 students of Faculty of Sports Science Medan State University who had not been trained as 200 meter runners and had fulfilled the inclusive and exclusive criteria, taken by using random sampling technique. They were divided into 2 (two) groups: WT+SM (Combination of weight training with sport massage) group and WT-SM (weight training without sport massage). The treatment was performed in 8 weeks with the frequency of exercise 3 times a week. **Results:** The research showed that there was significant influence of the treatment ($p \leq 0.05$) in the pre and post treatment in the WT+SM group and in the WT-SM group on the mean value of increase in glutathione peroxides content and the mean value of decrease in lactic acid content in 200 meter runners. There was no significant disparity ($p \geq 0.05$) of the increase in glutathione peroxides in 200 meter runners in the post treatment done by WT+SM group and by WT-SM group. There was significant disparity ($p \leq 0.5$) in the mean value of decrease in lactic acid content in the 200 meter runners in the post treatment done by WT+SM group and by WT-SM group. **Conclusion:** Combination weight training with sport massage was better in increasing glutathione peroxides content and decreasing lactic acid content in the 200 meter runners than that of only weight training.

Keywords: Weight Training, Sport Massage, Glutathione Peroxides, Lactic Acid, 200 meter Runner

Corresponding Author: Diky Setya Diningrat, Department of Biology, Faculty of Mathematics and Natural Sciences, Medan State University (Universitas Negeri Medan), 20221 Medan, North Sumatra, Indonesia Tel: +6181361362400 LiveDNA: 62.20093

Competing Interest: The authors have declared that no competing interest exists.

Name of the author and e-mail ID	Types of contribution
Dr. Novita Sari Harahap	Dr. Novita Sari Harahap researched on measuring antioxidant glutathione peroxides content and Accutrend Plus to measure lactic acid content and also measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure.
Dr. Aznan Lelo	Dr. Ambrosius Purba researched on data analysis on measuring antioxidant glutathione peroxides content and Accutrend Plus to measure lactic acid content
Dr. Ambrosius Purba	Dr. Ambrosius Purba researched on data analysis on measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure.
Dr. Diky Setya Diningrat	Dr. Diky Setya Diningrat was advisor for statistical data analysis and ELISA.

INTRODUCTION

Weight training is a kind of exercise which is known as resistance exercise. The positive effects of this kind of exercise include improvement in physical condition, fitness, and performance; therefore, it is usually recommended because it can decrease the risk for injury, and it functions as prophylaxis, compared with the other exercises^{1,2}. Every physical exercise like weight training has the potency for the incidence of free radicals which can bring about the incidence of oxidative stress. Our body has a number of limited antioxidants, while the activity of free radicals can reach 20% during the training. Therefore, the body has to have its own defense to protect from the attack of free radicals by performing regular exercises. The activity of antioxidant depends upon the intensity and the duration of exercises^{3,4}. Even though exercise can increase antioxidant defense system, a long period and heavy weight training can cause imbalance between free radical production and antioxidants which is called oxidative stress⁵.

The mechanism of the establishment of free radicals in 200 meter runners begins from muscle contraction during exclusive power movement. In the incidence of explosive power, ischemia will occur in muscles and be followed by muscle relaxation alternately. This muscle movement will cause blood flow to be hampered because of muscle contraction, and when the muscles are relaxed blood will flow rapidly. Ischemic reperfusion can cause the release of one electron of respiration chains so that free radicals are established^{6,7,8}. The establishment of free radicals in 200 meter runners can disturb mitochondria and they can even damage mitochondria. The disturbance of mitochondrion function which will cause the establishment of ATP, needed by 200 meter runners, is not fulfilled so that it causes fatigue easily. Therefore, the body has to have defense system; namely, antioxidants in order to protect from the attack of free radicals⁵.

Regular and rated standardized exercises like weight training should be done in order to decrease the negative impact of oxidative stress because of the increase in free radical production. Weight training is intended to increase physical condition of the 200 meter runners. The measurement of antioxidant enzyme content like glutathione peroxides (GPx) should also

be done in order to find out whether there is the increase in oxidative stress or not. In the use of energy in the 200 meter runners during the explosive power movement, most of the predominant source of energy is anaerobic lactic acid which potentially yields high lactic acid⁹. Physiologically, the increase in lactic acid content will decrease pH in body fluid. The decrease in pH can cause disturbance in the activity of oxidative enzyme in muscle cells so that the capacity of metabolism to produce ATP which is needed during muscle contraction is disturbed^{9,10}.

Recovery should be done immediately after doing the exercises in order to eliminate lactic acid in muscles by conducting sport massage. Immediate recovery after doing exercise is the most crucial factor in increasing an athlete's performance^{11,12}. Researches on the effect of sport massage therapy are relatively limited, and even some of them have different results about the influence of sport massage. For example, in the research conducted by Hemming et al (2000) on 8 amateur boxers; the treatment group was provided by sport massage within 20 minutes by using effleurage and petrissage techniques on the whole body, while the other group was provided with passive-relaxing treatment. The result showed that there was no significant disparity in lactic acid content and the performance of the boxers who were given sport massage treatment from those who were given passive-relaxing treatment¹³. Another controversial research result was conducted by Wiltshire et al (2005) on untrained individuals; it revealed that sport massage performed during the recovery in decreasing lactic acid content in blood was better than that with passive-relaxing treatment after 200 meter running activity¹⁴.

The two researches above indicate two different results about the influence of Sport Massage on athletes' physical condition. Therefore, this research was conducted because up to the present there has been no research which is concerned with the influence of Sport Massage on the performance of 200 meter runners that is measured with the duration of 200 meter race.

MATERIALS AND METHODS

Subjects: This study was not commercial research but part of doctoral dissertation. This research project was conducted from July 2016 to November 2016 in Pharmacology laboratory Faculty of Medicine North Sumatera University and Sport Science Laboratory of Medan State University. The research subjects were 20 students of Sports Science Study Program, UNIMED, with the criteria of males, 18 to 19 years old, untrained as 200 meter runners, physically healthy, willing to be the research subjects, and active in participating in the whole series of research procedure. They were divided randomly into two groups with 10 participants in the combination of weight training with sport massage (WT+SM) group, and 10 participants in weight training without sport massage (WT-SM) group.

Study Design: The research used experimental study method with Experimental Randomized Pre and Post test group design.

Materials and Device: Materials used in this research were vena blood to analyze GPx content and lactic acid content. The device used in the research was GPx kit to measure antioxidant glutathione peroxides content and Accutrend Plus to measure lactic acid content.

Research Preparation: This research had been approved by the Research Ethics Committee of the Faculty of Medicine, USU, in its Letter No. 270/KOMET/FKUSU/2016 after the research subjects had understood the objective of the research and its benefit. They were then willing to fill out their personal data and informed consent forms and to sign them. The measurement of physiological characteristics included body weight, body height, IMT, pulse while in rest, and blood pressure. Schedule for exercises was determined to be in 8 weeks with the frequency of

exercises 3 times a week. Exercise was performed in the afternoon, starting from 4:00 PM until 5:00 PM. The research procedure was done by professionals according to their fields. The subjects' blood and lactic acid content were taken by 3 medical analysts. Weight training program was trained by certified trainers who were competent as athletic trainers, while sport massage was done by 12 certified masseurs. During the process of the research, the subjects were not allowed to consume antioxidant substances like vitamin C and vitamin E or other supplements. They were also disallowed to do other physical activities except training program in the scope of the research. There would be no physical exercise for them, and they had to take enough rest.

Weight Training Procedure: The schedule for training was held in 8 weeks with frequency of exercise 3 times a week in the afternoon. Physical exercise was in the form of weight training plus technical training and sprint running tactics. The weight training was as follows: leg press: 8-12 RM, 3 sets; *squat jump*: 12-15 RM, 3 sets; *leg curl*: 15-25 RM, 3 sets; *sit ups*: 15-25 RM, sets; and Proprioceptive Neuromuscular Facilitation(PNF)exercise was done 3 times a week.

Sport Massage Treatment Procedure: Each time the exercise was done, WT+SM group was given specific sport massage to the 200 meter runners; it was partial massage in the lower legs by using effleurage technique. The method was by doing rubbing movement in the body, using the whole palms and fingers stuck on the lower leg areas; the movement was from the lower parts toward the upper ones. The duration of sport massage was 12 minutes (6 minutes from the right lower part of the legs, and the other 6 minutes from the left lower part of the legs)

Measuring Glutathione Peroxides and Lactic Acid: Pre-Exercise Program. Two days before measuring the Glutathione Peroxides and Lactic Acid, the research subjects were suggested that they not consume any drugs or other supplements; they were not allowed to do any physical exercise, and they had to take enough rest. They began to do warming-up in 15 minutes. After that, they did 200 meter race by recording their travel time. After they got to the finish line, ten minutes later, their blood was taken from vena mediana cubiti for assaying their Glutathione Peroxides and Lactic Acid content. The assaying of Glutathione Peroxides content was based on ELISA method with GPx Qayee-Bio Kit, stated in ng/ml unit, and spectrophotometer UV gauge with wavelength of 450 nm. The assaying of lactic acid content was done by using Accutrend Plus device with milimoles per liter (mmol/L) unit. By the end of the training program, the measurement of Glutathione Peroxides and Lactic Acid was the same as what had been done in sport massage (WT-SM).

Statistical analysis: All data were statistically analyzed with SPSS software (version 16). One-way analysis of variance (ANOVA) was used to study significant difference between means and significance level at $p = 0.05$.

RESULTS

The research subjects in the two groups, based on age were 18.65 years old, based on body weight were 61.65, based on height were 167.8, based on IMT were 22 based on systole blood pressure were 119.5, based on diastole blood pressure were 78., and based on pulse while in rest were 64.8. All in all, they showed no significant disparity ($p \geq 0.05$); therefore, all of them were homogeneous; they had the same characteristics, either physically or physiologically.

Table 1. Mean value of glutathione peroxides and lactic acid content in the 200 m runners

Variables	WT+SM	WT-SM	p
	Mean±SD	Mean±SD	

Glutathione Peroxides	Pre-exercise	218.35± 10.31	220.17 ±12.66	0.155
	Post-exercise	248.57 ± 16.59	232.82± 23.38	
	t-dependent	0.000*	0.01*	
Lactic Acid	Pre-exercise	8.23 ±0.27	8.23 ± 0.46	0.000*
	Post-exercise	6.34 ± 0.31	7.82 ± 0.21	
	t-dependent	0.000*	0.02*	

Explanation: * = significance ($p \leq 0.05$), SD : standard deviation.

Glutathione peroxides of the 200 meter runners increased significantly ($p \leq 0.05$) in the two groups in the pre-exercise, compared with that in the post-exercise; in the WT±SM group and in the WT-SM group (Table 1). The increase in glutathione peroxides in the 200 meter runners in the post-exercise in WT±SM group was higher than that WT-SM group. The percentage of the increase in glutathione peroxides content of the 200 meter runners. However, there was no significant disparity ($p \geq 0.05$) in the increase in glutathione peroxides content in the 200 meter runners in the post-exercise between WT+SM group and WT-SM group (Table 1).

Lactic acid content in the 200 meter runners decreased significantly ($p \leq 0.05$) in the two groups in pre-exercise, compared with that in the post-exercise: in the WT+SM group was and the WT-SM (Table 1). The decrease in lactic acid content in the 200 meter runners in the post-exercise in the WT+SM group was higher than that in the WT-SM group at the percentage of the decrease in the lactic acid content in the 200 meter runners. There was significant disparity ($p \leq 0.05$) in the decrease of lactic acid content in the 200 meter runners between the WT+SM group and the WT-SM group (Table 1).

DISCUSSION

Disparity in glutathione peroxides content in the 200 meter runners: Weight training is a type of sport which is intended to develop strength which uses gravitation against the gravity which is yielded by muscles through concentric or eccentric contraction. In this exercise, body muscles undergo contraction by using body weight or other devices to stimulate muscle growth/work strength, and resistance and by targeting certain muscle groups and types of movement. The exercise program can give benefit as maximal as possible and the risk as minimal as possible if it fulfills the training formulation which is called FITT (Frequency of Training, Intensity of Training, Time Spent, Type of Exercise)¹⁵.

Standardized exercise which is measured, guided, and programmed well can increase the physical condition of the 200 meter runners so that free radicals which exist as the result of the high intensity of physical exercise can be eliminated by the runners who have good physical condition. If they have good physical condition, their capability of producing endogenous antioxidants like Glutathione Peroxides will be better than the 200 meter runners who do not have good physical condition⁴. Weight training program is more focused on anaerobic exercise so that it will potentially yield relatively more lactic acid and can cause purine catabolism to become xanthenes and veins. This indicates the transition and acute vein deoxygenating which has the same problem with ischemic reperfusion. Purine catabolism and ischemic reperfusion will activate xanthenes oxides system which, in turn, establishes free radicals (O_2)^{4,16,17,18}. The research which was done by Best, et. al. (2002) on 100 meter swimmers revealed that there was the increase in oxidative stress biomarker like TBARS (4.1±0.7 versus 4.9±1.1), *Creatinine Kinase* (206±170.7 versus 244±176.9) and glutathione peroxides (0.52±0.06 versus 0.62±0.05). This research concluded that anaerobic exercise could cause oxidative stress; this increase in oxidative stress could hamper the establishment of Glutathione Peroxides and increased the use of endogenous antioxidants like Glutathione peroxides¹⁹.

The result of this research was in accordance with the research done by Groussard, et. al. (2003); it was aimed to determine the effect of implementing the combination of exercise with massage regularly on oxidant and antioxidant content. In their research, the research subjects were 25 people who did not do their sport regularly, and they did not receive any sport massage. They were randomly divided into three groups: control group (CG, n=9), exercise group (EG, n=8), and massage and exercise group (MEG, n=8). The result of the research showed that there was significant decrease ($p < 0.05$) in MDA content in the EG and MEG groups. It was also found that there was significant increase ($p < 0.05$) in glutathione peroxides in the EG and MEG groups, and there was significant increase ($p < 0.05$) in SOD content in the EG and MEG groups, compared with that in CG group. This indicated that physical activity which is done regularly and combined with massage can cause positive effect on the increase in antioxidants and endogen – GPx and SOD²⁰.

The result of this research was also in accordance with another research done by Azizbeigi et al. (2013); it was aimed to find out the influence of progressive resistance exercise on oxidative weight and the activity of enzymatic antioxidant in erythrocyte¹⁴. The research subjects were 20 men who were divided into two groups: progressive resistance exercise group with moderate intensity (50%) within 8 weeks and control group. The result of the research showed that progressive resistance exercise with moderate intensity significantly increased the activity of SOD erythrocyte ($p=0.014$), decreased MDA concentration ($p=0.030$), and GPx content tended to increase. The role of sport massage in the WT+SM group could give positive effect on the increase in GPx content even though statistically there was no significant disparity from WT-SM group, but there was the disparity in the increase in WT+SM group.

Disparity in lactic acid content in the 200 m runners: Exercise in WT+SM group was more significant in decreasing lactic acid in the 200 meter runners than that in the WT-SM group. This indicated that LPK and sport massage had significant influence on the decrease in lactic acid content in the 200 meter runners. The increase in lactic acid content is more significant in the WT+SM because weight training with standardized exercise which is measured, regular, and programmed well can increase tolerance against lactic acid and increase the capacity of bicarbonate buffer and phosphate in muscles. Besides that, sport massage can be efficient to eliminate lactic acid in the 200 meter runners²¹.

200 meter race athletics is the category of sprint and most of its predominant energy source is lactacyd anaerobic lactacyd which will yield relatively high lactic acid. The result of measuring lactic acid in the pre-exercise program in all treatment groups indicated the average increase of 8.23 mmol/L measured 10 minutes in the post 200 meter race. The increase in lactic acid was caused by the incidence of hypoxia in muscle tissues so that anaerobic metabolism which yields lactic acid occurs. The increase in lactic acid is in accordance with the result of the research done by Kawczyński et al., (2014) which found that there was the increase in lactic acid content in the 100 meter runners at the mean value of 9.51 mmol/L which was measured in the 10th minute in the post 100 meter race²². The higher the physical activity is, the more increasing the need for oxygen. The need for oxygen can be increased by increasing the function of heart and lungs. When physical activity is high while oxygen intake from heart and lungs is not optimal, anaerobic metabolism exist to fulfill the need for energy. This condition increases lactic acid content in blood and muscles. The increase in lactic acid can change reactive oxygen compound to be more reactive so that the physical condition of the 200 meter runners will not maximal¹⁹.

Physiologically, the increase in lactic acid content will decrease pH in the body liquid. The decrease in pH will affect the establishment of ATP which is needed during muscle contraction. Muscle contraction occurs because nervous impulse will reach axon terminal, followed by the release of acetylcholine (Ach) which immediately potentially stimulates muscle action in the

membrane of muscles sarcolemma. Potential action in muscles spreads through tubules T and causes reticulum sarcoplasm to release calcium ion to sarcoplasm. When there was the increase in lactic acid, the releasing process of acetylcholine and calcium ion from reticulum sarcoplasm will be disturbed. In consequence, there is no calcium bond with troponin. This bond will open tropomyosin which covers cross-bridge receptor. The opening of cross-bridge receptor from tropomyosin bond causes the actin to shift and the caput myosin to be stuck; in this process, ATP is needed²³. The increase in lactic acid will disturb the ATP splitting metabolism to be the energy.

Calcium ion low content in cytosol will decrease the muscle contraction power (leg muscle strength, abdominal and leg muscle resistance, leg muscle explosive power, running speed, and reaction speed). During the resting period, calcium ion enters again into reticulum sarcoplasm, and tropomyosin will close cross-bridge receptor again so the bond of actins and myosin is released and contraction will not occur. Physical activity will cause the bond of actins and myosin to become smooth so that muscle contraction highly depends on the higher or lower content of lactic acid in the 3200 meter runners²³.

Sport massage given in the pre-exercise to the WT+SM group in the 200 meter runners was an active recovery. It can cause muscles to be relaxed and regain the condition of blood vessel which used to be pinched by muscle contraction. Optimal relaxation can increase the backflow of vena blood in the lower muscle part of legs in bringing optimally lactic acid to the hearts of the 200 meter runners; therefore, it could increase the oxygen of muscle tissues and expedite the elimination of lactic acid²⁴. The increase in tissue oxygenation will increase the conversion of lactic acid in heart through the Cori cycle to pyruvate acid which, in turn, will be changed to glucose so that it will expedite the supply for energy and expedite recovery⁹.

Physiologically, massage is proved to be able to decrease heartbeat, to increase blood pressure, to increase blood circulation and lymph, to decrease muscle tension, too increase the range of joint movement, and to decrease pain. Physiology is widely used by athletes, either for their physical performance or for prevention, therapy, and rehabilitation for injuries and the side effect of sport²⁵. Sport massage on the body will give physiological effect such as increasing blood flow and lymphatic flow, stimulating nervous system, and increasing vena blood backflow. The increase in this vena blood backflow will efficiently return blood to the heart and drain off lactic acid which is deposited in muscles, and expedite the elimination of lactic acid in blood and muscles. It has been proved that effleurage technique is more effective in decreasing lactic acid deposit in muscle cells than that with moderate intensity sport^{24,26}.

This glucose can be reused as metabolism in active muscles or kept in muscles through the process of gluconeogenesis to become glycogen and can be used as an energy source. Pyruvate acid will enter mitochondria when the exercise is done below the lactate threshold (LT) and the establishment of energy is processed anaerobically, while in the exercise which is done over the lactate threshold (LT) like conventional exercise (heavy intensity exercise), pyruvate acid will be changed to lactic acid because the supply of oxygen is not enough to enter mitochondria²⁷. In a research done by Groussard, et. al., (2003) by using Wingate test, the research subjects pedaled bicycles within 30 seconds. Wingate test would stimulate anaerobic glycolysis metabolism by activating purine catabolism and lactic acid production. The data showed that anaerobic exercise by using Wingate test is the potential factor which causes oxidative stress. It is relatively the same as its movement pattern to increase anaerobic glycolysis in the 200 meter runners^{20,27}.

According to Best et al., (2008), an athlete can carry on his exercise in relatively high intensity in the condition of lactic acid of 6-7 mmol/L¹⁹. Therefore, the research of this research with the mean value of lactic acid content in the 200 meter runners in the LPK+SM of 6.34

mmol/L showed that the level of fatigue was still low so the the performance in the WT+SM group was better than that in the WT-SM group. The research done by Weerapong et al (2005) on individuals who were not trained found that sport massage during the recovery period could decrease lactic acid content in blood more than passive rest after the activity in 200 meter race^{25,28}. The research done by Monedero and Donne (2000) also found that sport massage could increase the elimination of lactic acid during the recovery period after doing high intensity exercise²⁸.

Pyne, et al., (2005) pointed out that sport massage done within 20 minutes could cause the decrease in lactic acid content in blood of 36.21%, and sport massage done within 45 minutes could decrease lactic acid in blood of 72.4%²⁷. The research done by Hanon, et al., (2010) was aimed to evaluate the influence of three methods for recovery – massage, active and passive recovery on the change in lactic acid concentration in blood, and the time of performance in swimming athletes²⁹. The research subjects were 17 swimming athletes. The result of the research showed that there was no disparity between active recovery and massage, and massage was more effective that passive recovery to eliminate lactic acid in blood. The research result was also in accordance with the research done by Devlin, et al., (2014)³⁰. It stated that active recovery after heavy exercise would eliminate more quickly the accumulation of lactic acid in blood than passive recovery.

The explanation above indicates that in regular physical exercise and measured standardized exercise the body will be adjusted so that the capacity to work in higher intensity will be able to use lactic acid during the exercise^{31,32}. This will cause more lactic acid to enter mitochondria and to be used for aerobic metabolism so that it can decrease the amount of lactic acid released to the blood flow. The combination with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage.

CONCLUSION

Combination of weight training with sport massage was better in increasing glutathione peroxides and decreasing lactic acid content in the 200 meter runners than weight training only.

SIGNIFICANCE STATEMENTS

This research showed that there is significant impact of combination of weight training with sport massage can increase the elimination of lactic acid more quickly than that of passive recovery, exercise without sport massage. This research provide scientific proof of the features of the combination of weight training with sport massage to improve the 200m runner's performance. Future research that needs to be done is to know the most effective type of weight training and type of sport massage to eliminate lactic acid and increase glutathione peroxide for a successful 200m runner achievement.

ACKNOWLEDGMENT

The authors would like to thank Ministry of research and higher education Republic Indonesia for providing Research Grants (Doctoral Research Grant, grant no. 038A/UN33.8/LL/2017.). The authors would also like to thank Medan State University (Universitas Negeri Medan) and Faculty of Medicine, North Sumatra University for providing the necessary facilities.

REFERENCES

- [1] Westcott, W., 2009. ACSM strength training guidelines: Role in body composition and health enhancement. *ACSM's Health & Fitness Journal*, 13(4), 14-22.

- [2] Novatchkov, H. and Baca, A., 2013. Artificial intelligence in sports on the example of weight training. *Journal of sports science & medicine*, 12(1), p.27.
- [3] Gomez-Cabrera, M.C., Domenech, E. and Viña, J., 2008. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radical Biology and Medicine*, 44(2), 126-131.
- [4] El Abed, K., Rebai, H., Bloomer, R.J., Trabelsi, K., Masmoudi, L., Zbidi, A., Sahnoun, Z., Hakim, A. and Tabka, Z., 2011. Antioxidant status and oxidative stress at rest and in response to acute exercise in judokas and sedentary men. *The Journal of Strength & Conditioning Research*, 25(9), 2400-2409.
- [5] Escribano, B.M., Tunez, I., Requena, F., Rubio, M.D., De Miguel, R., Montilla, P., Tovar, P. and Agüera, E.I., 2010. Effects of an aerobic training program on oxidative stress biomarkers in bulls. *Veterinari Medicina*, 55(9), 422-8.
- [6] Kostaropoulos, I.A., Nikolaidis, M.G., Jamurtas, A.Z. and Ikonou, G.V., 2006. Comparison of the blood redox status between long-distance and short-distance runners. *Physiological research*, 55(6), p.611.
- [7] Güzel, N.A., Hazar, S. and Erbas, D., 2007. Effects of different resistance exercise protocols on nitric oxide, lipid peroxidation and creatine kinase activity in sedentary males. *Journal of sports science & medicine*, 6(4), p.417.
- [8] Sahlin, K., Shabalina, I.G., Mattsson, C.M., Bakkman, L., Fernström, M., Rozhdestvenskaya, Z., Enqvist, J.K., Nedergaard, J., Ekblom, B. and Tonkonogi, M., 2010. Ultraendurance exercise increases the production of reactive oxygen species in isolated mitochondria from human skeletal muscle. *Journal of applied physiology*, 108(4), 780-787.
- [9] Astrand, P.O., Rodahl, K. (2003) *Textbook of Work Physiology, Physiological Base of Exercise*. New York: McGraw-Hill.
- [10] Hall, J.E., 2015. *Guyton and Hall textbook of medical physiology e-Book*. Elsevier Health Sciences.
- [11] Wiltshire, E.V., Poitras, V., Pak, M., Hong, T., Rayner, J. and Tschakovsky, M.E., 2010. Massage impairs postexercise muscle blood flow and "lactic acid" removal. *Medicine & Science in Sports & Exercise*, 42(6), 1062-1071.
- [12] Pinar, S., Kaya, F., Bicer, B., Erzeybek, M.S. and Cotuk, H.B., 2012. Different recovery methods and muscle performance after exhausting exercise: comparison of the effects of electrical muscle stimulation and massage. *Biology of sport*, 29(4), p.269.
- [13] Hemmings, B., Smith, M., Graydon, J. and Dyson, R., 2000. Effects of massage on physiological restoration, perceived recovery, and repeated sports performance. *British journal of sports medicine*, 34(2), 109-114.
- [14] Azizbeigi, K., Azarbayjani, M.A., Peeri, M., Agha-Alinejad, H. and Stannard, S., 2013. The effect of progressive resistance training on oxidative stress and antioxidant enzyme activity in erythrocytes in untrained men. *International journal of sport nutrition and exercise metabolism*, 23(3), 230-238.
- [15] Loturco, I., Ugrinowitsch, C., Roschel, H., Tricoli, V. and González-Badillo, J.J., 2013. Training at the optimum power zone produces similar performance improvements to traditional strength training. *Journal of sports science & medicine*, 12(1), p.109.
- [16] Valado, A., Pereira, L. and Paula, C., 2007. Effect of the intense anaerobic exercise on nitric oxide and malondialdehyde in studies of oxidative stress. *International Journal of Biology and Biomedical Engineering*, 1(1), 32-26.
- [17] Fisher-Wellman, K. and Bloomer, R.J., 2009. Acute exercise and oxidative stress: a 30 year history. *Dynamic medicine*, 8(1), p.1.

- [18] George, B.O. and Osharechiren, O.I., 2009. Oxidative stress and antioxidant status in sportsmen two hours after strenuous exercise and in sedentary control subjects. *African Journal of Biotechnology*, 8(3).
- [19] Best, T.M., Hunter, R., Wilcox, A. and Haq, F., 2008. Effectiveness of sports massage for recovery of skeletal muscle from strenuous exercise. *Clinical Journal of Sport Medicine*, 18(5), 446-460.
- [20] Groussard, C., Rannou-Bekono, F., Machefer, G., Chevanne, M., Vincent, S., Sergent, O., Cillard, J. and Gratas-Delamarche, A., 2003. Changes in blood lipid peroxidation markers and antioxidants after a single sprint anaerobic exercise. *European journal of applied physiology*, 89(1), 14-20.
- [21] Mika, A., Mika, P., Fernhall, B. and Unnithan, V.B., 2007. Comparison of recovery strategies on muscle performance after fatiguing exercise. *American journal of physical medicine & rehabilitation*, 86(6), 474-481.
- [22] Kawczyński, A., Kobiałka, K., Dariusz, M., Paweł, C., Adam, M., Adam, Z. and Jan, C., 2015. Blood lactate concentrations in elite Polish 100 m sprinters. *International Journal of Performance Analysis in Sport*, 15(1), 391-396.
- [23] McArdle, W.D., Katch, F.I. and Katch, V.L., 2010. *Exercise physiology: nutrition, energy, and human performance*. Lippincott Williams & Wilkins.
- [24] Cafarelli, E. and Flint, F., 1992. The role of massage in preparation for and recovery from exercise. *Sports medicine*, 14(1), 1-9.
- [25] Weerapong, P., Hume, P.A. and Kolt, G.S., 2005. The mechanisms of massage and effects on performance, muscle recovery and injury prevention. *Sports medicine*, 35(3), 235-256.
- [26] Goats, G.C., 1994. Massage--the scientific basis of an ancient art: Part 2. Physiological and therapeutic effects. *British journal of sports medicine*, 28(3), 153-156.
- [27] Pyne, D.B., Lee, H.A.M.I.L.T.O.N. and Swanwick, K.M., 2001. Monitoring the lactate threshold in world-ranked swimmers. *Medicine and Science in Sports and Exercise*, 33(2), 291-297.
- [28] Monedero, J. and Donne, B., 2000. Effect of recovery interventions on lactate removal and subsequent performance. *International journal of sports medicine*, 21(08), 593-597.
- [29] Hanon, C., Lepretre, P.M., Bishop, D. and Thomas, C., 2010. Oxygen uptake and blood metabolic responses to a 400-m run. *European journal of applied physiology*, 109(2), 233-240.
- [30] Devlin, J., Paton, B., Poole, L., Sun, W., Ferguson, C., Wilson, J. and Kemi, O.J., 2014. Blood lactate clearance after maximal exercise depends on active recovery intensity. *Journal of Sports Medicine and Physical Fitness*, 54(3), 271-278.
- [31] Leprêtre, P.M., Lopes, P., Thomas, C. and Hanon, C., 2012. Changes in cardiac tone regulation with fatigue after supra-maximal running exercise. *The Scientific World Journal*, 2012.
- [32] Ali, S.R., Koushkie, M.J., Asadmanesh, A. and Salesi, M., 2012. Influence of massage, active and passive recovery on swimming performance and blood lactate. *The Journal of sports medicine and physical fitness*, 52(2), 122-127.