



Andaliman

by Ciciek Suryani

THE
Character Building
UNIVERSITY

Submission date: 30-Nov-2021 06:37AM (UTC+0700)

Submission ID: 1715722084

File name: om_North_Sumatra_Province_of_Indonesia_Using_Maturase_K_Gene.pdf (888.03K)

Word count: 5291

Character count: 26701



DNA Barcoding of Andaliman (*Zanthoxylum acanthopodium* DC) from North Sumatra Province of Indonesia using *maturase K* Gene

Authors:

Cicik Suriani, Eko Prasetya*, Tri Harsono, Johannes Manurung, Hary Prakasa, Dina Handayani, Miftahul Jannah and Yuanita Rachmawati

*Correspondence: ekoprasetya.biologi@gmail.com

8

DOI: <https://doi.org/10.21315/tlsr2021.32.2.2>

Highlights

- cpDNA *maturase K* gene marker sequence can be used as DNA barcodes to identify *Zanthoxylum acanthopodium*.
- Based on the cpDNA *maturase K* gene sequence on *Zanthoxylum acanthopodium* had very high homology (97.2%).
- The genetic distance between *Zanthoxylum acanthopodium* using the cpDNA *maturase K* gene sequence is very small (<1%).

THE
Character Building
UNIVERSITY

TLSR, 32(2), 2021

© Penerbit Universiti Sains Malaysia, 2021

DNA Barcoding of Andaliman (*Zanthoxylum acanthopodium* DC) from North Sumatra Province of Indonesia Using *Maturase K* Gene

¹Cicik Suriani, ¹Eko Prasetya*, ¹Tri Harsono, ¹Johannes Manurung, ¹Hary Prakasa, ¹Dina Handayani, ²Miftahul Jannah and ³Yuanita Rachmawati

⁶
¹Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Medan, Jl. Willem Iskandar, Pasar V, Medan Estate, Medan, 20221, North Sumatra, Indonesia

²³
²Department of Biology, Faculty of Sciences and Technology, Islamic University of Assyafiyah, Jl. Jatiwaringin Raya, No 12, East Jakarta, 17411 Indonesia

³Department of Biology, Faculty of Sciences and Technology, Islamic State University Sunan Ampel Surabaya, Jl. Ahmad Yani 117 Surabaya, East Java, 60237, Indonesia

³
Publication date: 29 June 2021

To cite this article: Cicik Suriani, Eko Prasetya, Tri Harsono, Johannes Manurung, Hary Prakasa, Dina Handayani, Miftahul Jannah and Yuanita Rachmawati. (2021). DNA barcoding of Andaliman (*Zanthoxylum acanthopodium* DC) from North Sumatra Province of Indonesia using *maturase K* gene. *Tropical Life Sciences Research* 32(2): 15–28. <https://doi.org/10.21315/tlsr2021.32.2.2>

To link to this article: <https://doi.org/10.21315/tlsr2021.32.2.2>

¹⁰
Abstract: Andaliman (*Zanthoxylum acanthopodium* DC) is a native plant of North Sumatra province. *Zanthoxylum acanthopodium* is a member of *Rutaceae* family widely found in northern Sumatra, Indonesia. The aim of this study was to barcode *Z. acanthopodium* in North Sumatra province, Indonesia based on cpDNA *maturase K* (*matK*). Samples were collected in seven localities across six regions of North Sumatra province. Phylogenetic analysis was conducted using Maximum likelihood method. The results of phylogenetic analysis indicate that *Z. acanthopodium* is a monophyletic group that is derived from a common ancestor. The results of the phylogenetic tree construction show that there is a grouping of accession between *Z. acanthopodium* species separated from other species in the *Zanthoxylum* genus as well as those of the *Rutaceae* family. The results showed that cpDNA *matK* marker can effectively be used as DNA barcoding to identify *Z. acanthopodium*.

Keywords: *Z. acanthopodium*, *Maturase K*, North Sumatra, DNA Barcoding

¹⁰ INTRODUCTION

Andaliman (*Zanthoxylum acanthopodium* DC) is a native plant of North Sumatra province in Indonesia (Suryanto *et al.* 2004). *Zanthoxylum acanthopodium* is a member of the *Rutaceae* family that is widely spread throughout the northern part of Sumatra, Indonesia (Siregar 2003). This fruit is commonly used as

*Corresponding author. ekoprasetya.biologi@gmail.com

spice in traditional Batak cuisine (Kristanty & Suriawati 2015). Spice made of *Z. acanthopodium* is known by the name of "batak pepper" since it is prominent in traditional Batak cuisine (Hidayah 2015).

Zanthoxylum is a member of the Rutaceae (Pani 1993) family which consists of around 200 species spread across the region of Central Asia and North America (Hartley 1966) with natural characteristics of being thorny bushes, small trees with bushy and branched leaves and thorny stems, producing edible fruits with strong-smelling leaves (Chyau *et al.* 1996). *Z. acanthopodium* is known for its distinct citrus-like flavour and spiciness with unique taste, rendering numbness and sharp pain on one's tongue (Wijaya 2000). This plant has also been used to preserve foods such as raw fish and tofu (Parhusip *et al.* 1999).

Zanthoxylum is a complex genus with many different species and is not well-studied (Arun & Paridhavi 2012). Species in the *Zanthoxylum* genus have many similarities in its visible morphological characters, making it difficult to distinguish species within this genus (Gupta & Mandi 2013). Therefore, DNA sequence analysis is required to identify species within the *Zanthoxylum* genus. Research using molecular markers was performed to facilitate identification using morphological markers due to its higher stability (Yunus 2007) and resistance to environmental differences and robust plant life, resulting in more accurate data (Julisaniah *et al.* 2008).

Research on identification of *Z. acanthopodium* using DNA barcoding from chloroplast DNA (cpDNA) is still rare. Feng *et al.* (2015) analysed genetic variations and relationships between species in the *Zanthoxylum* genus using sequence-related amplified polymorphism (SRAP) marker. Putri *et al.* (2016) and Sembiring *et al.* (2015) analysed the genetic diversity of *Z. acanthopodium* using the Random Amplified Polymorphic DNA (RAPD) marker. Gupta and Mandi (2013) established DNA fingerprints using Amplified Fragment Length Polymorphism (AFLP) markers to differentiate *Z. acanthopodium* from *Z. oxyphyllum*.

Chloroplast DNA can be used to reveal diversity and trace evolutionary family tree of *Z. acanthopodium*. Chloroplast DNA has been widely used for phylogenetic studies of various plants such as *Bouea* (Harsono *et al.* 2017), Sedoideae subfamily (Lim & Choi 2018), *Solanum* (Olmstead & Palmer 1997), and Aurantioideae subfamily (Bayer *et al.* 2009). Chloroplast DNA is used as a barcoding because it is easy to purify with a very conservative character and low evolution rate, so it can be used for phylogenetic analysis between taxa in flowering plant families (Clegg & Durbin 1990; Tsumura *et al.* 1996).

Chloroplast DNA is a double-chain DNA of circular shape (Didriksen 2010) which consists of various genes such as *rbcl*, *trnL-F* and *matK* (Kress & Erickson 2007; Kalangi *et al.* 2014). The Consortium for the Barcode of Life (CBOL) recommended *rbcl* and *matK* as the standard barcode (Hollingworth *et al.* 2009). The *matK* gene is more commonly used in various researches compared to *rbcl* due to its specific level of accuracy at species level (Yu *et al.* 2011). Maturation of *matK* is a gene present in the chloroplast and located between exons 5' and 3' of *trnK* and lysine-tRNA (Enan & Ahmed 2014). The *matK* gene has been widely used as barcode in angiospermae plants (Yu *et al.* 2011). In the present paper, we

use the cpDNA *maturase K* marker to identify *Z. acanthopodium* and evaluated its use as DNA barcoding marker.

MATERIALS AND METHODS

Plant Samples

Z. acanthopodium plant sample is obtained from various regions in North Sumatra province which are represented by six regions namely South Tapanuli, North Tapanuli, Humbang Hasudutan, Dairi, Simalungun and Toba Samosir (Fig. 1). The samples used in this research are fresh leaf samples obtained by field exploration. The outgroup used to compare the barcoding sequences obtained were *Citrus x paradisi* and *Melicope glabra*, while the ingroup used were several species from the genus *Zanthoxylum* (Table 1).



Figure 1: Sampling locations in North Sumatra, Indonesia at seven points in six regencies. (Source: Google Map and processed using ArcGis software).

Table 1: Sample of *Z. acanthopodium* and *outgroup* species of the *Zanthoxylum* genus and from Rutaceae family.

No	Species	Accession Number	Origin
1	<i>Zanthoxylum holtzianum</i> (outgroup)	JX518057.1	South Africa
2	<i>Z. capense</i> (outgroup)	JQ025016.1	South Africa
3	<i>Z. davyi</i> (outgroup)	JX517950.1	South Africa
4	<i>Z. avicennae</i> (outgroup)	HQ415288.1	China
5	<i>Z. nitidum</i> (outgroup)	AB925028.1	Cambodia
6	<i>Z. fagara</i> (outgroup)	JQ589092.1	Costa Rica
7	<i>Z. acuminatum</i> (outgroup)	JQ589550.1	Costa Rica
8	<i>Z. delagoense</i> (outgroup)	KX146326.1	Mozambique
9	<i>Z. punctatum</i> (outgroup)	KJ012828.1	Puerto Rico
10	<i>Z. monophyllum</i> (outgroup)	KJ012827.1	Puerto Rico
11	<i>Z. flavum</i> (outgroup)	KJ012826.1	Puerto Rico
12	<i>Citrus x paradisi</i> (outgroup)	JN315360.1	India
13	<i>Melicope glabra</i> (outgroup)	KJ709002.1	Singapore
14	<i>Z. acanthopodium</i>	Sample	Indonesia
15	<i>Z. acanthopodium</i>	Sample	Indonesia
16	<i>Z. acanthopodium</i>	Sample	Indonesia
17	<i>Z. acanthopodium</i>	Sample	Indonesia
18	<i>Z. acanthopodium</i>	Sample	Indonesia
19	<i>Z. acanthopodium</i>	Sample	Indonesia
20	<i>Z. acanthopodium</i>	Sample	Indonesia

DNA Isolation and Amplification of cDNA *matK* sequence

The DNA isolation was carried out by following the procedure of the GeneJet Plant Genomic DNA Purification Kit (Thermo Fisher Scientific, Waltham, MA, USA). A total of 100 mg of *Z. acanthopodium* leaf sample was crushed with an additional Lysis Buffer A of 350 μ L. The solution is homogenised with the addition of 50 μ L Lysis Buffer B and 20 μ L of RNase A and incubated at 65°C for 10 min. Subsequently, the solution was added with 130 μ L of precipitation solution and centrifuged to isolate the supernatant. The supernatant in turn was added with 400 μ L of Plant gDNA Binding Solution and 400 μ L of 96% ethanol. The solution is then transferred into a purification column and centrifuged at 8,000 rpm for 1 min. The purification process was carried out using Wash Buffer I and Wash Buffer II in the purification column. The genomic DNA elution was performed with the addition

of 100 μL of Elution Buffer, followed by centrifugation at 10,000 rpm for 1 min. Purified DNA is then stored at -20°C .

The *matK* sequence is amplified using *matK*-F 5'-ACC CAG TCC ATC TGG AAA TCT TGG TTC-3' and *matK*-R 5'-CGT ACA GTA CTT TTG TGT TTA CGA G-3' primers (Ki-Joong Kim, School of Life Sciences and Biotechnology, Korea University, Korea, unpublished) with total reaction volume of 25 μL [2.5 μL of DNA template; 2.5 μL of *matK*-F primer; 2.5 μL of *matK*-R primer; 5 μL of distilled water, and 12.5 μL of PCR mix (MyTaq HS Red Mix (Bioline, USA))] mixture with a final concentration of template DNA of 50 ng. Amplification of cpDNA *matK* sequence with a predenaturation condition of 5 min at 97°C , followed by 40 cycles under denaturation reaction conditions 94°C for 5 min, annealing at 52°C , and extension at 72°C for 1 min, then the PCR process terminated with post-extension at 72°C for 5 min. PCR products were visualised using agarose gel plus 5 μL of SYBR® Safe DNA Gel Stain (Invitrogen, USA). Results PCR products that show positive results (DNA bands are clearly visible) will be sent to the First Base DNA Sequencing Service in Singapore for sequencing.

Barcoding and Phylogenetic Analysis

The result of *matK* sequencing was analysed using Bioedit 7.0.1 (Hall 1999) program to determine consensus sequence. Phylogenetic tree, nucleotide composition, and genetic distance were carried out using MEGA (Molecular Evolutionary Genetic Analysis) version 7 (Kumar *et al.* 2016) program based on alignment of sequence data. The method used for analysis is Maximum Likelihood with 1,000 bootstrap replicates.

RESULTS

Amplification of *matK* gene from the chloroplast of *Z. acanthopodium* genome was successful. The result of PCR visualised using agarose shows a single band, which means that the *matK* sequence has been successfully amplified (Fig. 2).

The result of *matK* gene sequence alignment shows that the *matK* gene sequence in *Z. acanthopodium* consists of 850 characters. Based on the data, 827 of the characters are conservative sequences, two are potentially informative parsimony characters, and five are variable sites. The result of alignment showed that the *matK* gene sequence on *Z. acanthopodium* had very high homology level (97.3%). The base frequency of the *matK* gene sequence on *Z. acanthopodium* is 35.16% (T), 19.11% (C), 27.34% (A), and 18.35% (G). This sequence is rich in T/A (62.47%), whereas G/C content is 37.48% (Table 2).




Figure 2: Visualisation of PCR results of *matK* gene sequences with agarose: (3) South Tapanuli; (4) North Tapanuli; (5) Humbang Hasudutan; (11) Toba Samosir 1; (6) Dairi; (9) Simalungun; (10) Toba Samosir 2.

Table 2: Composition of nucleotides, A/T content, and G/C content in the *matK* gene sequence on *Z. acanthopodium*.

Species	Composition (%)				Total	Content (%)	
	T(U)	C	A	G		A/T	G/C
<i>Z. acanthopodium</i> (Humbang Hasudutan)	34.9	19.4	27.1	18.6	850	62	38
<i>Z. acanthopodium</i> (Dairi)	35.4	19.2	27.1	18.3	804	62.5	37.5
<i>Z. acanthopodium</i> (Simalungun)	34.8	18.7	28.1	18.4	787	62.9	37.1
<i>Z. acanthopodium</i> (Tapanuli Selatan)	35.4	19.2	27	18.4	808	62.4	37.6
<i>Z. acanthopodium</i> (Tapanuli Utara)	35.1	19.3	27.2	18.4	794	62.3	37.7
<i>Z. acanthopodium</i> (Toba Samosir 1)	35.3	19.1	27.4	18.1	827	62.7	37.2
<i>Z. acanthopodium</i> (Toba Samosir 2)	35.2	18.9	27.5	18.3	829	62.7	37.2
Average	35.16	19.11	27.34	18.36	814.14	62.5	37.45

4 The phylogenetic tree presented in Fig. 3 was constructed using 1 Maximum Likelihood and 1000x bootstrap methods. This method is used to identify differences in genetic distance and analyse similarity between samples.




Figure 3: The phylogenetic tree of *matK* sequence from *Z. acanthopodium* and outgroup (species in the *Zanthoxylum* genus and Rutaceae family) as a result of reconstruction using the Maximum Likelihood method based on the kimura-2-parameter model. Branching is analysed with a bootstrap value of 1000x.

There are total of seven accessions of *Z. acanthopodium* clustered on the same branch and separated from other species of the *Zanthoxylum* genus and those from the Rutaceae family. Analysis of phylogenetic trees based on the *matK* sequence shows that the *Zanthoxylum* genus originated from one common ancestor. The results of this analysis also show that the *matK* sequence can be used as DNA Barcoding on *Z. acanthopodium*. It also indicates that the genetic distance between *Z. acanthopodium* species is very low compared to that between species in the *Zanthoxylum* genus or Rutaceae family. The lowest genetic distance between *Z. acanthopodium* is found in *Z. acanthopodium* from Dairi, Tapanuli Selatan and Toba Samosir 2. The genetic distance between the lowest *Zanthoxylum* genus is between *Z. capense* and *Z. davyi* (0.001) while the highest genetic distance is between *Z. nitidum* with *Z. capense* (0.023). Outgroup species from the Rutaceae family other than the *Zanthoxylum* genus showed significant differences (see Table 3).

Table 3: Genetic distance of Andaliman (*Z. acanthopodium*) with species in *Zanthoxylum* genus and Rutaceae family.

No	Species	1	2	3	4	5	6	7	8	9	10
1	<i>Citrus x paradisi</i> (JN315360.1)	0									
2	<i>Melicope vitiflora</i> (KM894552.1)	0.095	0								
3	<i>Z. acuminatum</i> (JQ589550.1)	0.054	0.066	0							
4	<i>Z. avicennae</i> (HQ415288.1)	0.054	0.068	0.012	0						
5	<i>Z. capense</i> (JQ025016.1)	0.057	0.070	0.018	0.021	0					
6	<i>Z. davyi</i> (JX517950.1)	0.055	0.071	0.016	0.019	0.001	0				
7	<i>Z. delagoense</i> (KX146326.1)	0.052	0.063	0.013	0.016	0.010	0.009	0			
8	<i>Z. fagara</i> (JQ589092.1)	0.055	0.068	0.007	0.013	0.019	0.018	0.015	0		
9	<i>Z. flavum</i> (KJ012826.1)	0.055	0.068	0.010	0.013	0.019	0.018	0.015	0.012	0	
10	<i>Z. holtzianum</i> (JX518057.1)	0.050	0.067	0.012	0.015	0.006	0.004	0.004	0.013	0.013	0
11	<i>Z. monophyllum</i> (KJ012827.1)	0.055	0.066	0.010	0.016	0.022	0.021	0.018	0.012	0.012	0.016
12	<i>Z. nitidum</i> (AB925028.1)	0.055	0.071	0.013	0.010	0.023	0.021	0.018	0.015	0.015	0.016
13	<i>Z. punctatum</i> (KJ012828.1)	0.054	0.070	0.004	0.016	0.022	0.021	0.018	0.012	0.015	0.016
14	<i>Z. acanthopodium</i> (Humbang Hasudutan)	0.055	0.068	0.007	0.013	0.018	0.016	0.015	0.009	0.012	0.013
15	<i>Z. acanthopodium</i> (Dairi)	0.054	0.067	0.009	0.015	0.016	0.015	0.013	0.010	0.013	0.012
16	<i>Z. acanthopodium</i> (Simalungun)	0.054	0.070	0.009	0.015	0.016	0.015	0.013	0.010	0.013	0.012
17	<i>Z. acanthopodium</i> (Tapanuli Selatan)	0.054	0.067	0.009	0.015	0.016	0.015	0.013	0.010	0.013	0.012
18	<i>Z. acanthopodium</i> (Tapanuli Utara)	0.057	0.070	0.009	0.015	0.016	0.015	0.016	0.010	0.013	0.015
19	<i>Z. acanthopodium</i> (Toba Samosir 1)	0.057	0.070	0.009	0.012	0.019	0.018	0.016	0.010	0.013	0.015
20	<i>Z. acanthopodium</i> (Toba Samosir 2)	0.054	0.067	0.009	0.015	0.016	0.015	0.013	0.010	0.013	0.012

(continued on next page)

Table 3: (continued)

No	Species	11	12	13	14	15	16	17	18	19	20
11	<i>Z. monophyllum</i> (KJ012827.1)	0									
12	<i>Z. nitidum</i> (AB925028.1)	0.018	0								
13	<i>Z. punctatum</i> (KJ012828.1)	0.015	0.018	0							
14	<i>Z. acanthopodium</i> (Humbang Hasudutan)	0.012	0.015	0.012	0						
15	<i>Z. acanthopodium</i> (Dairi)	0.013	0.016	0.013	0.001	0					
16	<i>Z. acanthopodium</i> (Simalungun)	0.013	0.016	0.013	0.001	0.003	0				
17	<i>Z. acanthopodium</i> (Tapanuli Selatan)	0.013	0.016	0.013	0.001	0.000	0.003	0			
18	<i>Z. acanthopodium</i> (Tapanuli Utara)	0.013	0.016	0.013	0.001	0.003	0.003	0.003	0		
19	<i>Z. acanthopodium</i> (Toba Samosir 1)	0.013	0.013	0.013	0.001	0.003	0.003	0.003	0.003	0	
20	<i>Z. acanthopodium</i> (Toba Samosir 2)	0.013	0.016	0.013	0.001	0.000	0.003	0.000	0.003	0.003	0

DISCUSSION

Various molecular markers have been developed for the purpose of identification of plant species. One of the most recommended molecular markers for identification purposes is DNA barcoding. One of the gene sequences contained chloroplast DNA, the *matK* is a common barcode used in species identification. The *matK* gene is recommended by *The Consortium for the Barcode of Life* (Hollingsworth et al. 2009).

Andaliman (*Z. acanthopodium* DC) is a commonly found plant in the northern part of Sumatra and has an important role in the customs and culture of Batak tribe. The fruit of this plant is used in a variety of traditional Batak cuisines in the North Sumatra province, Indonesia (Kristanty & Suriawati 2015). Research on this plant focuses on secondary metabolites with benefits in health and food (Li et al. 2012).

The *matK-F* and *matK-R* primers used in this study successfully amplified the *matK* gene sequence at a length of 850 bp. Previous study indicated that this primer can amplify various species of *Angiospermae* trees at amplification length of 830 bp–857 bp (Tosh et al. 2016). This primer has also successfully amplified the genus *Abelmoschus* (Fattah et al. 2014), *Ficus* (Moraceae) (Li et al. 2012), *Sycygium* (Tallei et al. 2016), Liliaceae (Ma et al. 2014) and Vitaceae (Habib et al. 2017). The results showed that the sequence of *matK* gene on *Z. acanthopodium* had high homology level (97.3%). This value is higher than the homology level of 14 species of Anacardiaceae in the ITS core genome area of 75% (Hidayat et al. 2011), *trnL-F* sequence on *Bouea macrophylla* (97.26%) (Harsono et al. 2017), and sequences of *matK* genes in species within the Fabaceae family (96%) (Gao et al. 2011). This value is lower than the homology level of *trnL-F* gene sequence on *Bouea oppositifolia* (97.48%) (Harsono et al. 2017).

Table 3 shows that the genetic distance in the *matK* gene sequence on *Z. acanthopodium* is very small (<1%). This shows that the *matK* gene sequence has very high conserved. Variation of order of sequences in cpDNA are generally caused by a single nucleotide mutation that has occurred over a very long period (Fitmawati & Hartana 2010; Borsch et al. 2003). Changes in the nucleotide sequence in the chloroplast genome are able to provide important information about the species evolution because the chloroplast genome is inherited maternally where small changes take place over a very long period of time (Hancock 2003), whereas in the DNA core changes occur due to recombination of both parentals.

The result of phylogenetic tree analysis in Fig. 3 shows that phylogenetic tree produced is monophyletic with three main groups. The first group is those of *Z. acanthopodium* species consisting of seven accessions. The second group consists of species belonging to the genus *Zanthoxylum* in addition to *acanthopodium* species. The third group is an outgroup group that all species in the *Zanthoxylum* genus derived from *Citrus x paradisi* and *Melicope vitiflora* belonging to the Rutaceae family. This is consistent with the statement of Taberlet et al. (1991) which states that the chloroplast genome is best used for inter-species kinship analysis but is less suitable in interspecies kinship analysis.

4 The variety indicated by cpDNA markers is relatively different from the diversity shown by morphological markers. The patterns emerging from cpDNA markers do not necessarily correlate with patterns generated from morphological markers, and vice versa. This is possible because the expression at the morphological level is the result of recombination of two parents and environmental factors. In addition, the gene sequences located on chloroplast DNA experience a lower rate of evolution than that of core DNA (Taberlet *et al.* 1991). The noncoding area has a high mutation rate, making the variations appear more and more informative when compared with the coding area (Taberlet *et al.* 1991; Hamilton M B. (1999).

CONCLUSION

5 Based on the results of research in this study, it can be concluded that the cpDNA *matK* marker can be amplified by the length of 850 bp on *Z. acanthopodium*. cpDNA *matK* marker can be used as DNA barcoding to identify *Z. acanthopodium*. cpDNA *matK* markers can also be used to separate *Z. acanthopodium* from other species of the genus *Zanthoxylum* and separate from outgroups of the Rutaceae family (*Citrus x paradisi* and *Melicope vitiflora*).

12 ACKNOWLEDGEMENTS

This research was supported by Ministry of Research, Technology and Higher Education of the Republic of Indonesia through Higher Education Basic Research Grants 2019 (No. 7/E/KPT/2019).

REFERENCES

- Arun K and Paridhavi M. (2012). An ethno botanical phytochemical and pharmacological utilization of widely distributed species *Zanthoxylum*: A comprehensive overview. *International Journal of Pharmaceutical Invention* 2(1): 24–35.
- Bayer R J, Mabberley D J, Morton C, Miller C H, Sharma I K, Pfeil B E, Rich S, Hitchcock R and Sykes S. (2009). A molecular phylogeny of the orange subfamily (Rutaceae: Aurantioideae) using nine cpDNA sequences. *American Journal of Botany* 96(3): 668–685. <https://doi.org/10.3732/ajb.0800341>
- Borsch T, Hilu K W, Quandt D, Wilde V, Neinhuis C and Barthlott W. (2003). Noncoding plastid *trnT-trnF* sequences reveal a well resolved phylogeny of basal angiosperms. *Journal of Evolutionary Biology* 16(4): 558–576. <https://doi.org/10.1046/j.1420-9101.2003.00577.x>
- Chyau C C, Mau J L and Wu C M. (1996). Characteristics of the steam-distilled oil and carbon dioxide extract of *Zanthoxylum simulans* fruits. *Journal of Agricultural and Food Chemistry* 44(4): 1096–1099. <https://doi.org/10.1021/jf950577d>

- Clegg M T and Durbin M L. (1990). Molecular approaches to the study of plant biosystematics. *Australian Systematic Botany* 3(1): 1–8. <https://doi.org/10.1071/SB9900001>
- Didriksen A. (2010). Molecular biology and physiology of isolated chloroplasts from the algae *Vaucheria*. Master's thesis, Universitetet i Tromsø.
- Enan M R and Ahmed A. (2014). DNA barcoding based on plastid matK and RNA polymerase for assessing the genetic identity of date (*Phoenix dactylifera* L.) cultivars. *Genetics and Molecular Research* 13(2): 3527–3536. <https://doi.org/10.4238/2014.February.14.2>
- Fattah Y R, Kamu V S, Runtuwene M R and Momuat L I. (2014). Identifikasi barcode tumbuhan gedi merah (*Abelmoschus manihot* L.) dan gedi hijau (*abelmoschus moschatus*) berdasarkan gen matK. *Jurnal MIPA Unsrat Online* 3(2): 120–124. <https://doi.org/10.35799/jm.3.2.2014.5863>
- Feng S, Yang T, Liu Z, Chen L, Hou N, Wang Y and Wei A. (2015). Genetic diversity and relationships of wild and cultivated *Zanthoxylum* germplasm based on sequence-related amplified polymorphism (SRAP) markers. *Genetic Resources and Crop Evolution* 62(8): 1193–1204. <https://doi.org/10.1007/s10722-015-0222-x>
- Fitmawati F and Hartana A. (2010). Phylogenetic study of *Mangifera laurina* and its related species using cpDNA *trnL-F* spacer markers. *Hayati Journal of Biosciences* 17(1): 9–14. <https://doi.org/10.4308/hjb.17.1.9>
- Gao T, Sun Z, Yao H, Song J, Zhu Y, Ma X and Chen S. (2011). Identification of Fabaceae plants using the DNA barcode matK. *Planta Medica* 77(1): 92–94. <https://doi.org/10.1055/s-0030-1250050>
- Gupta D D, and Mandi S S. (2013). Species specific AFLP markers for authentication of *Zanthoxylum acanthopodium* & *Zanthoxylum oxyphyllum*. *Journal of Medicinal Plants Studies* 1(6): 1–9.
- Habib S, Dang V C, Ickert-Bond S M, Zhang J L, Lu L M, Wen J and Chen Z D. (2017). Robust phylogeny of *Tetrastigma* (Vitaceae) based on ten plastid DNA regions: Implications for infrageneric classification and seed character evolution. *Frontiers in Plant Science* 8: 590. <https://doi.org/10.3389/fpls.2017.00590>
- Hall T A. (1999). BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41(41): 95–98.
- Hamilton M B. (1999). Four primer pairs for the amplification of chloroplast intergenic regions with intraspecific variation. *Molecular Ecology* 8(3): 521–523.
- Hancock J F. (2003). *Plant evolution and the origin of crop species*. Second Edition. Wallingford, UK: CABI Publishing.
- Harsono T, Pasaribu N, Sobir S, Fitmawati F and Prasetya E. (2017). Phylogenetic analysis of Indonesian gandaria (*Bouea*) using molecular markers of cpDNA *trnL-F* intergenic spacer. *Biodiversitas Journal of Biological Diversity* 18(1): 51–57. <https://doi.org/10.13057/biodiv/d180108>
- Hartley T G. (1966). A revision of the Malesian species of *Zanthoxylum* (Rutaceae). *Journal of the Arnold Arboretum* 47(3): 171–221. <https://www.jstor.org/stable/43781566>
- Hidayah Z. (2015). *Ensiklopedi suka bangsa Indonesia*. Indonesia: Yayasan Pustaka Obor Indonesia.
- Hidayat T, Pancoro A and Kusumawaty D. (2011). Utility of matK gene as DNA barcode to assess evolutionary relationship of important tropical forest tree genus *Mangifera* (Anacardiaceae) in Indonesia and Thailand. *BIOTROPIA: The Southeast Asian Journal of Tropical Biology* 18(2): 74–80. <https://doi.org/10.11598/btb.2011.18.2.41>

- Hollingsworth M L, Clark A A, Forrest L L, Richardson J, Pennington R T, Long D G, Cowan R, Chase M W, Gaudeul M, & Hollingsworth P M. (2009). Selecting barcoding loci for plants: Evaluation of seven candidate loci with species-level sampling in three divergent groups of land plants. *Molecular Ecology Resources* 9(2): 439–457. <https://doi.org/10.1111/j.1755-0998.2008.02439.x>
- Julisaniah N I, Sulistyowati L, and Sugiharto A N. (2008). Analisis kekerabatan mentimun (*Cucumis sativus* L.) menggunakan metode RAPD-PCR dan isozim. *Biodiversitas* 9(2): 99–102. <https://doi.org/10.13057/biodiv/d090205>
- Tsumura Y, Kawahara T, Wickneswari R and Yoshimura K. (1996). Molecular phylogeny of Dipterocarpaceae in Southeast Asia using RFLP of PCR-amplified chloroplast genes. *Theoretical and Applied Genetics* 93(1–2): 22–29. <https://doi.org/10.1007/BF00225722>
- Kalangi C, Kamu V S, and Kumaunang M. (2014). Barcode DNA tanaman leilem (*Clerodendrum minahassae* L.) berdasarkan gen matK. *Jurnal MIPA Unsrat Online* 3(2): 108–112. <https://doi.org/10.35799/jm.3.2.2014.5861>
- Kress W J and Erickson D L. (2007). A two-locus global DNA barcode for land plants: The coding rbcL gene complements the non-coding trnH-psbA spacer region. *PLoS one* 2(6): e508. <https://doi.org/10.1371/journal.pone.0000508>
- Kristanty R E and Suriawati J. (2015). The Indonesian *Zanthoxylum acanthopodium* DC.: Chemical and biological values. *International Journal of PharmTech Research* 8(6): 313–321.
- Kumar S, Stecher G and Tamura K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33(7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Li H Q, Chen J Y, Wang S, and Xiong S Z. (2012). Evaluation of six candidate DNA barcoding loci in *Ficus* (Moraceae) of China. *Molecular Ecology Resources* 12(5): 783–790. <https://doi.org/10.1111/j.1755-0998.2012.03147.x>
- Lim M S and Choi S H. (2018). Estimation of phylogeny of nineteen Sedoideae species cultivated in Korea inferred from chloroplast DNA analysis. *The Horticulture Journal* 87(1): 132–139. <https://doi.org/10.2503/hortj.OKD-087>
- Ma H L, Zhu Z B, Zhang X M, Miao Y Y and Guo Q S. (2014). Species identification of the medicinal plant *Tulipa edulis* (Liliaceae) by DNA barcode marker. *Biochemical Systematics and Ecology* 55: 362–368. <https://doi.org/10.1016/j.bse.2014.03.038>
- Olmstead R G and Palmer J D. (1997). Implications for the phylogeny, classification, and biogeography of *Solanum* from cpDNA restriction site variation. *Systematic Botany* 22(1): 19–29. <https://doi.org/10.2307/2419675>
- Parhusip A J N, Sibuea P dan Tarigan A. (1999). Studi tentang aktivitas antimikroba alami pada andaliman. *Seminar Nasional Teknologi Pangan*. Jakarta, 12–13 October.
- Pirani J R. (1993). A new species and a new combination in *Zanthoxylum* (Rutaceae) from Brazil. *Brittonia* 45(2): 154–158. <https://doi.org/10.2307/2807498>
- Putri L A, Sinaga A O, Sembiring I M and Sinaga A. (2016). Assessing genetic diversity of Indonesian Andaliman (*Zanthoxylum acanthopodium* DC.) germplasm based on 7 RAPD markers. *Proceeding International Conference on Global Resource Conservation*. Universitas Brawijaya, Malang, September.
- Sembiring I M S, Putri L A and Setiada H. (2015). The application of five RAPD primers for genetic diversity analysis of North Sumatera's andaliman (*Zanthoxylum acanthopodium* DC.). *Jurnal Agroekoteknologi* 4(1): 1748–1755.
- Siregar B L. (2003). Andaliman (*Zanthoxylum acanthopodium* DC.) di Sumatera Utara: Deskripsi dan perkecambahannya. *Hayati Journal Bioscience* 10(1): 38–40.

Cicik Suriani et al.

- Suryanto E, Sastrohamidjojo H, Raharjo S and Tranggongo. (2004). Antiradical activity of andaliman (*Zanthoxylum acanthopodium* DC.) fruit extract. *Indonesian Food and Nutrition Progress* 11(1): 15–19. <https://doi.org/10.22146/jifnp.26>
- Taberlet P, Gielly L, Pautou G and Bouvet J. (1991). Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Molecular Biology* 17(5): 1105–1109. <https://doi.org/10.1007/BF00037152>
- Tallei T E, Irawan P D and Kolondam B J. (2016). DNA barcoding analysis of matK gene of some syzygium species. In: *Bioinformatics Workshop*. Al Azhar University Jakarta, 13–15 September 2016.
- Tosh J, James K, Rumsey F, Crookshank A, Dyer R and Hopkins D. (2016). Is DNA barcoding child's play? Science education and the utility of DNA barcoding for the discrimination of UK tree species. *Botanical Journal of the Linnean Society* 181(4): 711–722. <https://doi.org/10.1111/boj.12449>
- Wijaya C H. (2000). Isolasi dan identifikasi senyawa trigeminal aktif buah andaliman (*Zanthoxylum acanthopodium* DC). *Hayati Journal Bioscience* 7: 91–95.
- Yu J, Xue J H and Zhou S L. (2011). New universal matK primers for DNA barcoding angiosperms. *Journal of Systematics and Evolution* 49(3): 176–181. <https://doi.org/10.1111/j.1759-6831.2011.00134.x>
- Yunus A. (2007). Identifikasi keragaman genetik jarak pagar (*Jatropha curcas* L.) di Jawa Tengah berdasarkan penanda Isoenzym. *Biodiversitas* 8(3): 249–252. <https://doi.org/10.13057/biodiv/d080317>

UNIMED
UNIVERSITY
THE Character Building

Andaliman

ORIGINALITY REPORT

22%

SIMILARITY INDEX

20%

INTERNET SOURCES

10%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	digilib.unimed.ac.id Internet Source	5%
2	C Suriani, E Prasetya, T Harsono, D Handayani. "Habitat characteristics of Andaliman (<i>Zanthoxylumacanthopodium</i> DC) in North Sumatra using a GIS (Geographical Information System) approach", <i>Journal of Physics: Conference Series</i> , 2019 Publication	1%
3	web.usm.my Internet Source	1%
4	Submitted to Universitas Riau Student Paper	1%
5	jurnal.unimed.ac.id Internet Source	1%
6	doaj.org Internet Source	1%
7	Carmen Garcia-Davila, David Aldana Gomero, Jean-François Renno, Rossana Diaz Soria et al. "Molecular evidence for three	1%

genetic species of Dipteryx in the Peruvian Amazon", Genetica, 2019

Publication

8	jps.usm.my Internet Source	1 %
9	biodiversitas.mipa.uns.ac.id Internet Source	1 %
10	B L Siregar, L A M Siregar, T C Nisa, L A P Putri. "Flower morphology and inflorescence of Andaliman (<i>Zanthoxylum acanthopodium</i> DC.) from Dairi, North Sumatera", IOP Conference Series: Earth and Environmental Science, 2019 Publication	1 %
11	www.veterinaryworld.org Internet Source	1 %
12	www.microbiologyjournal.org Internet Source	1 %
13	academic.oup.com Internet Source	<1 %
14	www.plantsjournal.com Internet Source	<1 %
15	S Lailiyah, T Nusantara, C Sa'dijah, E B Irawan, Kusaeri, A H Asyhar. "Structuring students' analogical reasoning in solving algebra problem", IOP Conference Series: Materials Science and Engineering, 2018 Publication	<1 %

16	dspace.cas.upm.edu.ph Internet Source	<1 %
17	www.ccrjournal.com Internet Source	<1 %
18	Submitted to Anglia Ruskin University Student Paper	<1 %
19	hdl.handle.net Internet Source	<1 %
20	smujo.id Internet Source	<1 %
21	bmcgenomics.biomedcentral.com Internet Source	<1 %
22	res.mdpi.com Internet Source	<1 %
23	journal.uin-alauddin.ac.id Internet Source	<1 %
24	"Advances in Food Biotechnology", Wiley, 2015 Publication	<1 %
25	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
26	www.ukessays.com Internet Source	<1 %
27	E. V. Zhokhova, A. V. Rodionov, M. N. Povydysh, M. Yu. Goncharov, Ya. A.	<1 %

Protasova, G. P. Yakovlev. "Current State and Prospects of DNA Barcoding and DNA Fingerprinting in the Analysis of the Quality of Plant Raw Materials and Plant-Derived Drugs", *Biology Bulletin Reviews*, 2019

Publication

28

www.biorxiv.org

Internet Source

<1 %

29

www.nrcresearchpress.com

Internet Source

<1 %

30

docslide.us

Internet Source

<1 %

31

scholar.ufs.ac.za:8080

Internet Source

<1 %

32

tsukuba.repo.nii.ac.jp

Internet Source

<1 %

33

www.smujo.id

Internet Source

<1 %

34

E H Frederick, M T Sibero, A P Wijaya, E Syafitri et al. "Preliminary Evaluation of Anti Fish Pathogenic Bacteria and Metabolite Profile of Andaliman Fruit (*Zanthoxylum acanthopodium* DC.) Ethanol Extract", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

<1 %

35

researcharchive.lincoln.ac.nz

Internet Source

<1 %

- 36 www.freepatentsonline.com <1 %
Internet Source
-
- 37 www.plosone.org <1 %
Internet Source
-
- 38 www.thermoscientificbio.com <1 %
Internet Source
-
- 39 "The Date Palm Genome, Vol. 1", Springer
Science and Business Media LLC, 2021 <1 %
Publication
-

Exclude quotes On

Exclude matches Off

Exclude bibliography On

UNIVERSITAS NEGERI
MEDAN
UNIMED

THE
Character Building
UNIVERSITY