

Study of Thermodynamic and  
Structure of Cu(II) 1,2,4 H-  
Triazole Complex using The ab  
Initio Method

*by Asep Wahyu Nugraha*

THE  
*Character Building*  
UNIVERSITY

---

**Submission date:** 15-Mar-2022 03:33PM (UTC+0700)

**Submission ID:** 1784752097

**File name:** 8\_eai.18-10-2018.2287369\_-\_2019.pdf (454.08K)

**Word count:** 2858

**Character count:** 14401

# Study of Thermodynamic and Structure of Cu(II) 1,2,4 H-Triazole Complex using The ab Initio Method

Asep Wahyu Nugraha<sup>1</sup>, Zainuddin Muchtar<sup>1</sup>, Junifa Laila Sihombing<sup>1</sup>, Iis Siti Jahro<sup>2</sup>  
{aw.nugraha@unimed.ac.id}

<sup>1</sup>Physical Chemistry Division Faculty of Mathematics and Natural Sciences, Universitas Negeri Medan


<sup>2</sup>Inorganic Chemistry Division Faculty of Mathematics and Natural Sciences, Universitas Negeri Medan

**Abstract.** The complex of Cu(II) 1,2,4 H-triazole is one complex compound having a polymeric structure. The objective of this study determines the difference in energy formation and structure data of Cu(II) 1,2,4 H-triazole complex. The computational study used the Hartree-Fock method and the basis set 3-21G and 6-31G(d). The complex was studied of  $[\text{Cu}_2(\text{trz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$ . The result of a computational study with function/ basis set UHF/ 6-31G(d) shows the distance between Cu(II) ions for complexes with deprotonated ligands of 3,433Å, while complexes with undepronated ligands are 3,551Å. Bond length of Cu-N in the complex with deprotonated ligands are 1.953Å - 2.167Å, whereas for the complex with undepronated ligands are 2,061Å - 2,123Å. The difference of energy from a computational study using the function/ basis set UHF/ 6-31G(d) for the  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  complex is -3693.43 KJ/ Mol and the  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complex is -1666.01 KJ/ Mol.

**Keywords:** Cu(II) 1,2,4 H-triazole, the computational study, the structure data, and difference of energy

## 1 Introduction

The ion of Cu(II) has nine electrons in orbitals of d ( $d^9$ ) formed the octahedral structure in the complex compounds. The Cu(II) 1,2,4 H-triazole complex was formed the covalent coordination bonds with the octahedral structure. The ligand of 1,2,4 H-triazole (Htrz) is an intermediate field ligand. The atom of H, which is bonded the N4 atom from the Htrz ligand can be released to form a  $\text{trz}^{-1}$  ion. The structure of the Htrz and  $\text{trz}^{-1}$  ligand rings are presented in Figure 1.



**Fig 1.** The structure of the ligand rings of (a) Htrz and (b)  $\text{trz}^{-1}$

The Cu(II) ions with Htrz ligand have a polymeric structure in which between the Cu(II) ions is connected with the Htrz ligand bridge. The structure of the Cu(II) 1,2,4 H-triazole

(Cu(II)-Htrz) complex is similar to Fe(II) 1,2,4 H-triazole. The structure of the Cu(II) 1,2,4 H-triazole complex is shown in Figure 2.




Fig. 2 Polymeric complex structure of Cu(II)-Htrz.

Experimental studies of Cu(II)-Htrz complexes or derivatives have been carried out by various methods. Analysis of the single crystal structure of the complex  $[\text{Cu}(\text{hyetrz})_3](\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$  (hyetrz) 4-(2-hydroxyethyl)-1,2,4-triazole was carried out using Enraf-Nonius CAD4 diffractometer and EXAFS13 spectrometer. The final coordinate data and isotropic thermal parameters are visualized by CRYSTAL MAKER to get the crystal structure of the complex. The crystal structure obtained shows between Cu(II) ions are connected by three bridges  $\text{N}_1$ ,  $\text{N}_2$ -Htrz with a distance between Cu1-Cu2 of  $6.8530(8)\text{\AA}$  and Cu2-Cu3 of  $3.8293(2)\text{\AA}$ . Complex forms a chain with an angle between pairs of Cu(II) ions between Cu1-Cu2 and Cu2-Cu3 at  $175.18(2)^\circ$ . The data show the complex forms a chain that only slightly deviates from the linearly shaped chain (Garcia et al, 1997). The structural characteristics of  $[\text{Fe}(\text{NH}_2\text{trz})_3](\text{NO}_3)_2$  (1) and  $[\text{Cu}(\text{NH}_2\text{trz})_3](\text{NO}_3)_2 \cdot \text{H}_2\text{O}$  (2) complexes have been observed using SEM and XRPD. A single crystal complex (2) has been observed using XRD, lead to the complex structural parameter data is obtained (Dirtu et al., 2010). Both complex crystal form shows similarities based on SEM images with greater magnification.

An experimental study of the Cu(II)-Htrz complex was carried out by several previous researchers. The computational chemistry methods could predict the crystal structure of Cu(II)-Htrz complex. The data are compared with structural data has been obtained experimentally. The study determined differences energy data of Cu(II)-Htrz complex too. This study was used in the ab initio method with UHF functions and basis sets of 3-21G and 6-31G(d).

Determination of energy with the completion of the Schrödinger equation is carried out using analytic and semi-empirical methods (Hückel method) for a set of molecular orbitals obtained in the form of the variations of energy. This energy can be integrated as an orbital for each of these energies (one electron formalism). The wave functions for many electrons are a result of the Hartree (Hartree Product) wave function as stated in equation 1.

$$H = \sum_{i=1}^N h_i \quad (1)$$

where  $h_i$  is a wave function for each electron. An interaction between nuclei is considered zero because the core is considered motionless (Born Oppenheimer Approximation).

$$h_i = -\frac{1}{2} \nabla_i^2 - \sum_{k=1}^M \frac{Z_k}{r_{ik}} \quad (2)$$

where M represents the total number of cores

Slater proposed the addition of the Hartree SCF with Slater Determinant of the wave function. The Fock operator for one electron defined for each electron  $i$  is:

$$f_i = -\frac{1}{2}\nabla_i^2 - \sum_k \frac{Z_k}{r_{ik}} + V_i^{HF}\{j\} \quad (3)$$

The Hartree-Fock assumption states that each electron is in a fixed field produced by the other electron densities (Cramer, 2004).

Determination of structure, vibration frequency, relative energy, and dipole moments are calculated using basis sets 3-21G almost the same compared to the results using a basis set 6-21G (Binkley et al., 1980). Comparison of geometry parameters, normal vibration frequency modes, and dipole moments from computational using basis sets 3-21G and 3-21G\* according to the experimental data (Pietro et al., 1982). Determination of bond length using a basis set 6-21G is more in line with experimental results than using a basis set 3-21G (Gordon et al., 1982). The determination of energy using 6-31G\* is closer to the experimental results than the determination of energy using 3-21G\* (Schleyer et al., 1984). The calculated activation energy with MP4 (SDTQ)/ 6-31G\*// MP2/ 6-31G\* is in accordance with experimental data. One point calculation at MP2 / 6-31G\* level with geometry optimization at 3-21G gives an energy value that is in accordance with the experimental data (Spellmeyer et al., 1987).

The geometry optimization at Hartree-Fock (RHF) level with basis set of 3-21G and 6-31+G\* for the complex compounds with alkali metals. The results of this study obtained that the calculation results using a basis set 6-31+G\* are more suitable with experimental data than basis set 3-21G (Glendening et al., 1994). Results of a study of various compounds using a Local density functional (LDF) theory with a basis set of STO-3G and 3-21G obtained suitability with experimental data with an average deviation of 0.026Å. Specifically for the compound Fe(CO)<sub>5</sub> basis set 3-21G more in line with the experimental results compared to the STO-3G basis set. Determination of frequency the LDF results are better than the results of Hartree-Fock (Sosa et al., 1992). The results of the geometry optimization show that the basis set 6-31G(d) is in accordance with the experiment compared to STO-3G, 6-31G, 6-31+G, and 6-31+G(d) (Foresman et al., 1991). Based on comparisons of computational results with experimental data show that the basis set 3-21G and 6-31G(d) is quite good to be used in the determination of structure and energy. Prediction of the structure and the difference of energy [Fe<sub>4</sub>(Htrz)<sub>8</sub>(trz)<sub>4</sub>]<sup>4+</sup> and [Fe<sub>4</sub>(Htrz)<sub>12</sub>]<sup>8+</sup> complexes used the function/basis set TPSSH/ TZVP. The results showed that the energy difference in the formation of the [Fe<sub>4</sub>(Htrz)<sub>8</sub>(trz)<sub>4</sub>]<sup>4+</sup> was -9285.974 KJ/ Mol and [Fe<sub>4</sub>(Htrz)<sub>12</sub>]<sup>8+</sup> of -3501.534 KJ/ Mol (Nugraha et al., 2015).

## 2 Computational Method

The computational calculations to determine the structure and the difference of energy using the UHF function and the basis set of 3-21G and 6-31G(d). The software used is NWChem 6.6 (Valiev et al., 2010) for computational chemistry calculations, Jmol (<http://www.jmol.org/>) and Avogadro (Hanwell et al., 2012) for visualization structure complexes. The data obtained from computational calculations are used to determine the difference in energy complexes (Ochterski, 2000).

### 2.1 Determination of complex structured data.

Based on the structured from the geometry optimization results obtained the bond length and bond angle data for the Cu(II)-Htrz complex.

### 2.2 Determination of complex energy difference.

The difference in energy complex is obtained energy data from computational chemistry calculations.

11

### 3 Results and Discussion

#### 3.1 Determination of Complex Structure

The result of geometry optimization with function/ basis set UHF/ 6-31G(d) shows that the Cu(II)-Htrz complex has a polymeric structure and between Cu(II) ions connected by three Htrz ligand rings. Structural visualization of the results of geometry optimization of the complexes  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  are shown in Figure 3.




Fig. 3. Structure from geometry optimization results of the (a)  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and (b)  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes.

The parameter's structure of the Cu(II)-Htrz complex were studied data on the distance between Cu(II) ions and the bond length of Cu-N. The distance between Cu(II) ions as a result of geometry optimization with functions/basis set UHF/ 3-21G for the complex with deprotonated ligands is 3.635Å, while complex with undepronated ligands of 3.194Å. The distance between Cu(II) ions as a result of geometry optimization with function /basis set UHF/ 6-31G(d) for the complex with deprotonated ligands is 3.433Å, while complex with undepronated ligands is 3.551Å. The data of distance between Cu(II) ions of  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes result of computational calculations with UHF functions and 3-21G and 6-31G(d) basis sets stated in Table 1.

Table 1 The distance between Cu(II) ions result in geometry optimization

Functions/ basis sets	The distance between Cu(II) ions (Å)	
	$[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$	$[\text{Cu}_2(\text{Htrz})_6]^{4+}$
UHF/ 3-21G	3.635	3.194

The bond length of Cu-N on deprotonated complexes are shorter than undepronated complexes. For the deprotonated complexes, the bond length of Cu-N on the deprotonated ring is shorter than the undepronated ring. The computational study with the function/ basis set UHF/ 3-21G shows that the bond length of Cu-N in the complexes with the deprotonated ligand are 1.953Å - 2.087Å, while the undepronated ligand is 1.885Å - 2.109Å. The computational calculation of the function/ basis set UHF/ 6-31G(d) shows that the bond length of Cu-N in the complexes with a deprotonated ligand is 1.953Å - 2.167Å, while the undepronated ligand is 2.063Å - 2.123Å. Bond length of Cu-N in the  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes result of the computational calculations with UHF functions and the basis sets 3-21G and 6-31G(d) presented in Table 2.

**Table 2** The bond length of Cu-N as a result of geometry optimization

Functions/ basis sets	The bond length of Cu-N (Å)		
	$[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$		$[\text{Cu}_2(\text{Htrz})_6]^{4+}$
	The deprotonated ring	The undepronated ring	
UHF/ 3-21G	1.953 – 1.967	2.035 - 2.087	1.885 – 2.109
UHF/ 6-31G(d)	1.953 – 2.002	2.077 – 2.167	2.063 – 2.123

### 3.2 Determination of complex energy

Determination energy of Cu(II)-Htrz complex was carried out on complexes with one deprotonated ligand and undepronated ligands for the UHF functions and 3-21G and 6-31G(d) basis sets. The data on energy difference of  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes result of computational calculations on UHF functions and 3-21G and 6-31G(d) basis sets are presented in Table 3.

**Table 3** The difference energy of  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes with the UHF function and 3-21G and 6-31G(d) basis sets.

The complexes	Functions / Basis Sets	The amount of Energy (Ht)				The Energy Difference	
		Cu	Htrz	trz	Complexes	(Ht)	(kJ/mol)
$[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$	UHF/ 3-21G	-1630.11	-239.36	-238.79	-4696.88	-1.65	-4338.32
	UHF/ 6-31G(d)	-1637.79	-240.73	-240.17	-4720.25	-1.41	-3692.43
$[\text{Cu}_2(\text{Htrz})_6]^{4+}$	UHF/ 3-21G	-1630.11	-239.36		-4697.18	-0.81	-2130.70
	UHF/ 6-31G(d)	-1637.79	-240.73		-4720.58	-0.63	-1666.01

THE  
*Character Building*  
UNIVERSITY



#### 4 Conclusion

The computational calculation to function/ basis set UHF/ 6-31G(d) shows the distance between Cu(II) ions in the complexes with deprotonated ligands of 3.433 Å, while undepronated ligand is 3.551 Å. The bond length of the Cu-N in the complexes with a depronated ligand is 1.953 Å – 2.167 Å, whereas for the complexes with undepronated ligand are 2.063 Å – 2.123 Å. The difference energy from computational chemistry calculations for  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  and  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complexes are -3693.43 KJ/ Mol; and -1666.01 KJ/ Mol.  $[\text{Cu}_2(\text{Htrz})_4(\text{trz})_2]^{2+}$  complex is more stable than the  $[\text{Cu}_2(\text{Htrz})_6]^{4+}$  complex.

#### 8:knowledgements

This work was supported by State University of Medan, The Ministry of Research, Technology and Higher Education the Republic of Indonesia by the KDBK Research Grant (Project No. 282/UN33.8/PL/2018).

#### References

- [1] Garcia Yann, Petra J. van Koningsbruggen, Georges Bravic, Philippe Guionneau, Daniel Chasseau, Giovanni Luca Cascarano, Jacques Moscovici, Katia Lambert, Alain Michalowicz, and Olivier Kahn, (1997): Synthesis, Crystal Structure, EXAFS, and Magnetic Properties of catena-Poly[μ-tris(4-(2-hydroxyethyl)-1,2,4-triazole- $N^1,N^2$ )copper(II)] diperchlorate trihydrate: Relevance with the structure of the iron(II) 1,2,4-triazole spin transition molecular materials, *Inorg. Chem*, **36**, 6357 – 6365.
- [2] Dirtu, Marinela M, Christine Neuhausen, Anil D Naik, Aurelian Rotaru, Leonard Spinu, and Yann Garcia (2010): Insights into the Origin of Cooperative Effects in the Spin Transition of  $[\text{Fe}(\text{NH}_2\text{trz})_3](\text{NO}_3)_2$ : the Role of Supramolecular Interactions Evidenced in the Crystal Structure of  $[\text{Cu}(\text{NH}_2\text{trz})_3](\text{NO}_3)_2 \cdot \text{H}_2\text{O}$ , *Inorganic Chemistry*, **49**, 5723-5736.
- [3] Cramer, Christopher J., (2004): Essentials of Computational Chemistry Theories and Models, second edition, John Willey & Sons Ltd, The Atrium Southern Gate Chichester West Sussex, England
- [4] Binkley, J. Stephen, John A. Pople, and Warren J. Hehre. (1980). Self-consistent Molecular Orbital Methods. 2 1. Small Split-Valence Basis Sets for First-Row Elements. *Journal of the American Chemical Society*, **102**, 3, 939-947
- [5] Pietro, W. J. , M. M. Francl, W. J. Hehre, D. J. DeFrees, J. A. Pople, and J. S. Binkley, (1982). Self-consistent Molecular Orbital Methods. 24. Supplemented Small Split-Valence Basis Sets for Second-Row Elements. *J. Am. Chem. Soc.*, **104**, 19, 5039-5048
- [6] Gordon, Mark S., J. Stephen Binkley, John A. Pople, William J. Pietro, and Warren J. Hehre. (1982). Self-consistent Molecular-Orbital Methods. 22. Small Split-Valence Basis Sets for Second-Row Elements. *J. Am. Chem. Soc.*, **104**, 10, 2797-2803.
- [7] Schleyer, Paul von Rague, Timothy Clark, Alexander J. Kos, Gunther W. Spitznagel, Cornelia Rohde, Dorit Arad,la K. N. Houk, and Nelson G. Rondan. (1984). Structures and Stabilities of a-Hetero-Substituted Organolithium and Organosodium Compounds. Energetic Unimportance of Second-Row d-Orbital Effects. *J. Am. Chem. Soc.*, **106**, 22, 6468 – 6475.
- [8] Spellmeyer, David C. and K. N. Houk, (1987), Transition Structures of Pericyclic Reactions. Electron Correlation and Basis Set Effects on the Transition Structure and Activation Energy of the Electrocyclization of Cyclobutene to Butadiene, *J. Am. Chem. Soc.*, **110**, 11, 3412-3416
- [9] Glendening, Eric D., David Feller, and Mark A. Thompson, (1994), An ab Initio Investigation of the Structure and Alkali Metal Cation Selectivity of 18-Crown-6, *J. Am. Chem. Soc.*, **116**, 23, 10657-10669.

- [10] Sosa, Carlos, Jan Andzelm, Brad C. Elkin, Erich Wimmer, Kerwin D. Dobbs, and David A. D. (1992); A Local Density Functional Study of the Structure and Vibrational Frequencies of Molecular Transition-Metal Compounds, *J. Phys. Chem.*, **96**, 16, 6630-6636
- [11] Foresman, James B., Martin Head-Gordon, John A. Pople, and Michael J. Frisch (1991), Toward a Systematic Molecular Orbital Theory for Excited States, *J. Phys. Chem.*, **96**, 1, 135-149
- [12] Nugraha, Asep Wahyu, Djulia Onggo, and Muhamad A Martoprawiro, (2015), Computational study of structure and stability of polymeric complexes of  $Fe_4(Htrz)_8(trz)_4^{4+}$  and  $[Fe_4(Htrz)_{12}]^{8+}$ , *Procedia Chemistry*, **17**, 9 – 15
- [13] Valiev, M., E.J. Bylaska, N. Govind, K. Kowalski, T.P. Straatsma, H.J.J. van Dam, D. Wang, J. Nieplocha, E. Apra, T.L. Windus, W.A. de Jong "NWChem: a comprehensive and scalable open-source solution for large scale molecular simulations" *Comput. Phys. Commun.* **181**, 1477 (2010) doi:10.1016/j.cpc.2010.04.018
- [14] Jmol: an open-source Java viewer for chemical structures in 3D. <http://www.jmol.org/>
- [15] Hanwell M. D., D. E. Curtis, D. C. Lonie, et al., Avogadro, *Journal of Cheminformatics*, **4**, (17) (2012).
- [16] Ochterski, Joseph W., (2000): *Thermochemistry in Gaussian*, [help@gaussian.com](mailto:help@gaussian.com)



THE  
*Character Building*  
UNIVERSITY



# Study of Thermodynamic and Structure of Cu(II) 1,2,4 H-Triazole Complex using The ab Initio Method

## ORIGINALITY REPORT

**22%**  
SIMILARITY INDEX

**11%**  
INTERNET SOURCES

**21%**  
PUBLICATIONS

**1%**  
STUDENT PAPERS

## PRIMARY SOURCES

- 1** A W Nugraha, I S Jahro, D Onggo, M A Martoprawiro. " Study of thermodynamics and complex structures on spin transitions of [Fe (Htrz) (trz) ]Cl and [Fe (Htrz) (trz) ]Cl complexes using the Density Functional Theory method ", Journal of Physics: Conference Series, 2020  
Publication **8%**
- 2** [s3-eu-west-1.amazonaws.com](https://s3-eu-west-1.amazonaws.com)  
Internet Source **4%**
- 3** Nugraha, Asep Wahyu, Djulia Onggo, and Muhamad A. Martoprawiro. "Computational Study of Structure and Stability of Polymeric Complexes of [Fe<sub>4</sub>(Htrz)<sub>8</sub>(trz)<sub>4</sub>]<sup>4+</sup> and [Fe<sub>4</sub>(Htrz)<sub>12</sub>]<sup>8+</sup>", Procedia Chemistry, 2015.  
Publication **2%**
- 4** Marinela M. Dîrtu, Christine Neuhausen, Anil D. Naik, Aurelian Rotaru, Leonard Spinu, Yann Garcia. " Insights into the Origin of Cooperative Effects in the Spin Transition of

[Fe(NH trz) ](NO ) : the Role of Supramolecular Interactions Evidenced in the Crystal Structure of [Cu(NH trz) ](NO ) ·H O ", Inorganic Chemistry, 2010

Publication

---

5	<a href="http://manualzz.com">manualzz.com</a> Internet Source	1 %
6	Yann Garcia, Petra J. van Koningsbruggen, Georges Bravic, Philippe Guionneau et al. " Synthesis, Crystal Structure, EXAFS, and Magnetic Properties of -Poly[ $\mu$ -tris(4-(2-hydroxyethyl)-1,2,4-triazole- , )copper(II)] Diperchlorate Trihydrate: Relevance with the Structure of the Iron(II) 1,2,4-Triazole Spin Transition Molecular Materials ", Inorganic Chemistry, 1997 Publication	1 %
7	<a href="http://digilib.unimed.ac.id">digilib.unimed.ac.id</a> Internet Source	1 %
8	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	1 %
9	<a href="http://docplayer.net">docplayer.net</a> Internet Source	<1 %
10	<a href="http://jptc.srbiau.ac.ir">jptc.srbiau.ac.ir</a> Internet Source	<1 %

---

11 Hong Zhou. "A tetranuclear copper(II) complex: crystal structure, assembly, EPR, electrochemistry and magnetism", Journal of Coordination Chemistry, 09/16/2008

Publication

<1 %

12 Fatma Kandemirli. " Synthesis and Theoretical Study of Vanadium Oxotrichloride Complex of Hexachlorotriphosphazene and Investigations of Some Simple Cyclic Triphosphazenes ", Phosphorus, Sulfur, and Silicon and the Related Elements, 2003

Publication

<1 %

13 Handbook of Computational Chemistry, 2012.

Publication

<1 %

14 Ping. Du, David A. Hrovat, Weston Thatcher. Borden. "Ab initio calculations on the ring opening of cyclopropane radical cation. Trimethylene radical cation is not a stable intermediate", Journal of the American Chemical Society, 2002

Publication

<1 %

15 [www.pubs.acs.org](http://www.pubs.acs.org)

Internet Source

<1 %

16 Geanina Marin. "Structural Diversity in Metal–Organic Frameworks Derived from Binuclear Alkoxo-Bridged Copper(II) Nodes

<1 %

17

Michael Kruppa, Burkhard König. "Reversible  
Coordinative Bonds in Molecular Recognition",  
Chemical Reviews, 2006

Publication

<1 %

18

Vaiana, L.. "Binuclear Co(II), Ni(II), Cu(II) and  
Zn(II) complexes with Schiff-bases derived  
from crown ether platforms: Rare examples of  
ether oxygen atoms bridging metal centers",  
Polyhedron, 20100728

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On