SAJ27-19929812.pdf

by Ahmad Imran

Submission date: 16-May-2023 01:59PM (UTC+0700)

Submission ID: 2094444662

File name: SAJ27-19929812.pdf (180.14K)

Word count: 5209

Character count: 26768

The Effect of Quick Strength Training on the Agility and Leg Power of Futsal Junior Athletes

Imran Akhmad^{1,*}, Rahma Dewi¹, Suprayitno², Galih Priyambada³

¹Department of Sport Coaching Education, Faculty of Sport Science, Universitas Negeri Medan, Medan, North Sumatra, 20221
Indonesia

²Department of Physical Education Health and Recreation, Faculty of Sports Sciences, Universitas Negeri Medan, Medan, North Sumatra, 20221 Indonesia

³Sports Education Study Program, Faculty of Teacher Training in Education, Universitas Muhammadiyah Kalimantan Timur, Indonesia

Received November 1, 2022; Revised March 20, 2023; Accepted April 7, 2023

Cite This Paper in the Following Citation Styles

(a): [1] Imran Akhmad, Rahma Dewi, Suprayitno Galih Priyambada, "The Effect of Quick Strength Training on the Agility and Leg Power of Futsal Junior Athletes," International Journal of Human Movement and Sports Sciences, Vol. 11, No. 2, pp. 477 - 483, 2023. DOI: 10.13189/saj.2023.110227.

(b): Imran Akhmad, Rahma Dewi, Suprayitno, Good Priyambada (2023). The Effect of Quick Strength Training on the Agility and Leg Power of Futsal Junior Athletes. International Journal of Human Movement and Sports Sciences, 11(2), 477 - 483. DOI: 10.13189/saj.2023.110227.

Copyright©2023 by authors, all rights reserved. Authors agree that this article remains permanently open access under the terms of the Creative Commons Attribution License 4.0 International License

Abstract This research aimed to examine the effect of Quick Strength Training (QST) on agility and leg power. It used experimental methods with a pre-test-post-test design. A purposive sampling technique was used with several criteria. Before the treatment was executed, the 11 participants in each group were tested for their agility with the side step test and leg power with vertical jump. The 11 participants in the experimental group performed QST training program for 42 days or six weeks and the focus was carried out with an intensity of three times a week, then in the other control group, there were 11 participants, as a CON, who did not do any physical activities for 6 weeks. After six weeks of treatment, the 22 participants performed side step and vertical jump as a post-test. There was an increase in the average of their agility; CON test before treatment and test after treatment QST test before treatment and test after treatment (p=0.007). The increase in leg strength in participants is also a record for a result; CON test before treatment and test after treatment and QST pre-test with post-test (p=0.000). Agility test before treatment has similarities between CON and QST (p=0.171), but there was a post-test agility difference between CON and QST (p=0.538). There was also a difference in the pre-test leg power between CON and QST (p=0.061), and there was a post-test leg power difference between CON and QST (p=0.001). The results showed that

QST increased futsal athletes' agility and leg power.

Keywords Quick Strength, Agility, Leg Power

1. Introduction

Futsal is a sport that has the same characteristics as football. The only differences are the size of the field, the type and size of the ball, game duration and number of players. Because the field size is smaller, the intensity of the game is very high. Futsal athletes need to have the ability to quickly dribble and change directions to stay in control of the ball. The game requires the athletes to move fast and change direction precisely, which is characterized as agility [1], [2]. Aside from agility, the athletes should have leg power to run fast and kick. Performing such agility optimally with leg power is determined by the athletes' good physical condition. It requires good quality of the local muscles of the lo through strength training. Strength training is a method of exercise that can improve several physical components, including agility and leg power. Strength training and power training can be integrated to improve muscle strength, power and speed [4].

Lower muscle strength training, for instance, has a significant effect on the agility, leg power and neuromuscular adaptation of football athletes [3]. Furthermore, core strength training can increase the power, speed and agility of football athletes [5]. The core muscle part has a dominant role for soccer athletes and has an important role in the components of human motion [6]. These studies become the basis that strength training can increase agility and leg power. The use of weight training methods that are often done by trainers today tends to use classic methods. In a similar vein, circuit methods are also often done. However, these methods tend to be at the stage of anatomical adaptation aimed at stimulating muscles to receive weights. Strength training circuit, on the other hand, has a more significant effect than plyometric agility [7].

The main objectives of physical preparatory training are to increase athletes' functional potential and develop their bio-motor abilities to the highest standards [8]. Speed is a complex ability because in general, speed is an ability that allows a soccer player to move as quickly as possible at the level of specific resistance [9]. Agility is an important quality in most field sports. In badminton, for example, agility demonstrates the ability to move the legs quickly and precisely is part of the abilities that can be used in the ability to move [10]. It is the ability to maintain and control a correct body position while quickly changing direction [11], [12]. Acceleration is a part that cannot be taken lightly, and has an important position and can be collaborated with footsteps and speed [13], [14]. movement speed, acceleration, agility, and dexterity skills are some of the important parts in the components of the game of football that must always be considered [14]. Findings suggested that the peak of muscle strength was of great significance in the human lower body in soccer games, success in the game is supported by components of physical movement which include the lower extremities [15]. Due to the shared characteristics between futsal and soccer, futsal then can also be speed, acceleration and agility are part of the group and category in question.

Previous related studies have investigated the effect of Quick Strength Training (QST) on agility, particularly of football athletes [16]. Other studies focused on the analysis of movement ability, speed and agility have a dominant position on motion activity in games and matches [15], [17]. A programmed quick strength training was shown to be effective in increasing agility. This was demonstrated in a study in which the experimental group performed quick strength exercises for 8 weeks, giving a greater impact compared to the group conditioned as a control on football athletes with measurements on the agility component [18], [19]. Similarly, a short-term agility training program (3 weeks duration) improved agility test results among young professional soccer players [20]. Other studies revealed that both squat and plyometric trainings were necessary for improving hip and thigh power production as measured by vertical jump [21]. To sum up, these studies exhibited that the components of agility included balance, coordination, power and speed [22].

This study focused on junior futsal players who often struggled to improve their agility. Improvement in their agility is necessary due to a variety of reasons. The players have to be able to gain acceleration and to be accurate in passing and shooting, all of which require leg power. To improve their performance, accurate training programs with the right number of sessions and quality training are crucial. The success of such programs is significantly influenced by the right types and methods of strength training, such as Quick Strength Training (QST).

The literature has indicated that the use of circuit method in QST and its effect on agility and 143 power is still under exploration. Thus, this research aimed to discover the effect of Quick Strength Training (QST) on futsal athletes' agility and leg power. It put the emphasis on the implementation of programmable QST with circuit method as the training method to improve their agility and leg

2. Materials and Methods

This study aimed to discover the effect of Quick Strength Training (QST) training on agility (AG) and Leg Power (LG) of futsal athletes aged between 17 and 19 years old. This study used experimental methods with a pre-test-posttest design. The 22 futsal athletes formed into two conditions, where the first condition is called an experiment with a total of 11 people, while the other 11 athletes were put into a control group. Before the treatment, both groups performed initially an AG test using sidestep test and an LG test using Vertical Jump test.

The experimental group normally went under a 6-week QST for 18 meetings. The frequency of the training was 3 times a week every Monday, Wednesday and Friday, while the group did the training every other day After 18 exercises, the final test with the same test parameters was carried out.

Participant

The population in this study consisted of all members of Futsal Academy Ligasu-ABAA of 48 athletes. A purposive sampling technique with several criteria was used. The criteria were as follows: (1) male, (2) aged from 17-19 years, (3) Training duration of at least 1 year, and (4) willingness to be included as research participants with parents' consents related to COVID-19 situation. Twenty-two athletes met the criteria and became the sample. The characteristics of the 22 athletes that were randomly selected as the object of this study were: 18.05 \pm 0.84 years of age; 63.18 ± 5.36 kg of body weight; and 168.22 ± 5.14 cm of height. All participants were controlled during the treatment by maintaining a balanced nutrition, prohibiting any use of supplements, and keeping them from getting injured during treatment so that the test was not interrupted. All participants received the same

QST program during the training period. To measure their agility, the side step test was used [23], and to measure their leg power, the vertical jump test was used [24].

Sidestep Test Implementation

Participants stood in the middle of the field with 1 meter wide and the point of 60 cm. Participants performed activities starting from the middle point and did a sideways right-left jumping movement with a distance of 30 cm for 1 minute. Score was recorded based on the correct number of returns obtained by the participants.

Vertical Jump Test Implementation

Standing on the side wall to reach the specified place without jumping, then followed by a jump as high as possible to get optimal results, the distance from the initial achievement without a prefix and the second with a jump is the result that is recorded.

Experimental design

The 22 are irregularly divided into two conditions called groups. The first was the QST experimental group consisting of 11 participants with age = 18.09 ± 0.83 ; body weight = 64.82 ± 4.69 kg; height = 168.82 ± 4.77 cm). The second group was the control groups (CON) consisting of

also 11 participants with age = 18 ± 0.89); body weight = 61.55 ± 5.70 kg; height = 167.64 ± 5.66 cm. Before the treatment was conducted, all of the participants were tested for their agility using the side step test as well as their leg power using the vertical jump test. The 11 participants in the experimental group performed QST-Circuit Training (CT) program three meetings in one week and carried out for six weeks. Design of the QST CT program is explained in table 1 and figure 1.

CT program consisted of 8 stations, namely; (1) Alternate Side Lunges (ASL), (2) Explosive Step Up (ESU), (3) Quick Step Side (QSS), (4) Scissors Jump (SJ), (5) Lateral Step In-Out (LSIO), (6) Quick Squat Jump (QSJ), (7) Side to Side Jump (SSJ), and (8) High Knee Run (HKR). The other 11 participants in the control group (CON) did not carry out activities or treatment as in the experimental group, but did physical activities such as the usual training model given. in this context, there is no manipulation of new training models. Afterwards, the 22 participants performed side step test and vertical jump test as a post-test. The data analysis technique used SPSS program and previously it is 3 pecessary to carry out normality and homogeneity tests with a significant level of p<0.05.

Table 1. The 6-Week QST-CT program

Week	Variable	QST/Number of Station (1-8)	
	Intensity	80%	
	Number of repetitions	8 sec	
1-2	Number of set	4	
	Interval (set/Station)	90-120 sec/30 sec	
	Intensity	90%	
	Number of repetitions	8 sec	
3-4	Number of set	5	
	Interval (set/Station)	90-120 sec/30 sec	
	Intensity	100%	
	Number of repetitions	8 sec	
5-6	Number of set	5-6	
	Interval (set/Station)	90-120 sec/30 sec	



Figure 1. Implementation of QST-CT

3. Result

The agility (AG) pre-test data showed the average score of the control group; QST (47.00±4.54); (46.36±4.15) with a score ranging from 42-55;40-53. The description of the pre-test leg power (LP) data showed the average score of the control group; QST of (48.00±6.05 cm); (53.36±5.59 cm) with a score ranging from 41-60 cm; 47-65 cm. A description of the pre-test data is shown in table 2.

Table 3 shows the results of tests carried out before treatment on AG. There is no difference or has the same results in the average value of the test that was carried out before the intermediate treatment CON (47.00±4.54) and QST (46.36±4.15) (p=0.171). However, there was a difference in the post-test average score of CON (48.18±4.31) and QST (48.91±4.25) (0.538) on AG. For the results of the AG on the CON pre-test mean \pm SD and post-test mean \pm SD, the results obtained are (p = 0.007). Furthermore, for the results of the AG on the QST pre-test mean \pm SD and post-test mean \pm SD and post-test mean \pm SD, the results obtained were (p = 0.000).

There was a difference in the pre-test average score between CON (48.00±6.08) and QST (53.36±6.56)

(p=0,061). Furthermore, there was a difference in the post-test average score of CON (50.55 \pm 6,67) and QST (62.55 \pm 7.48) (p=0,002 on LP. For the results of the LP on the CON pre-test mean \pm SD and post-test mean \pm SD, the results obtained are (p = 0.000). Furthermore, for the results of the AG on the QST pre-test mean \pm SD and post-test mean \pm SD and post-test mean \pm SD, the results obtained were (p = 0.000).

Based on the results of calculations using the ShapiroWilk test, it can be concluded that there is significance in the LP pre-test score data (sig = 0.125) and LP post-test (sig = 0.202). Thus, it can be concluded that the data is a normally distributed sample on the grounds that the significance level is more than 0.05. Based on the results of the calculation of the homogeneity test (Levene test), it is obtained that the significance value on the average LP test before treatment and test after treatment data is 0.733, provided that probability and significance of more than 0.05, it can be said that the population has the same variance. Furthermore, the mean difference test was carried out paired 10 pple t test interpretation I based on the pair 1 output, the Sig. (2-tailed) of 0.000 < 0.005, it can be concluded that there is a difference in the average pre-test LP and post-test LP.

Session Monday Tuesday Wednesday Thursday Saturday Sunday Advance Technique Afternoon (4 Basic QST QST Day of rest QST Days of rest - 6 FM) Technique /Games

Table 2. Conditioning Program- an Overview of In-season Training (6-week QST intervention)

Table 3. Data description of the pre-test

AG	CON	QST	P-value	
pre-test mean±SD	47.00±4.54	±4.54 46.36±4.15		
post-test mean±SD	48.18±4.31	48.91±4.25 0,53		
P-value	0,007	0,000		
LP	CON	QST	P-value	
pre-test mean±SD	48,00±6,08	53,36±6,56 0,061		
post-test mean±SD	50,55±5,67	62,55±7,48 0,002		
P-value	0,000	0,000		

4. Discussion

The first findings showed that there was an increase in AG after implementing QST CT for 6 weeks. Improved QST exercise results on AG were shown in the pre-test average scores of the pre-test of 47.00±4.54 with the st-test 48.18±4.15 and tested a significant level of -3.36 with a significant level of p<0.05. The findings of this study provided information on the rapid rhythmic QST CT that stimulated rapid muscle contraction in various positions, both in direction changes and body position. Many strength training methods are proven to improve the neuromuscular work system for young futsal athletes [25] both separately and in the combination of weight training with plyometric [26]. The positive response of QST exercise also had an effect on increasing AG after performing 8 weeks of exercise [18]. This study was slightly different from other studies, as it conducted the application of quick strength exercises with the circuit method for 6 weeks. However, QST can also be done by implementing a circuit method.

A decrease in lactic acid levels and an increase in bicarbonate and muscle phosphate buffers can be done by doing disabled and optimal exercises [27]. The combined exercise took a form of QST with the characteristics of weight strength training with fast rhythm. The 6-week QST CT program directly affected the improvement of the futsal athletes' agility [28]. This was in accordance with the results of several studies. They showed that plyometric exercises increased LP, while QST significantly improved soccer athletes' vertical jump and agility [29]. Other studies also revealed that the resistance training increased sets with 4-8 repetitions for 6 weeks was agility [30], and fast rhythm strength training with a variety of movements increased agility quickly [31].

These results became the basis for the implementation

of the fast motion programmatic circuit strength exercise. The second finding of this study suggested that there was a significant effect of QST on LP. This is evident from the mean difference test Interpretation of Paired Samp 10 test 1 based on pair 1 output, the Sig value is obtained. (2-tailed) of 0.000 <0.005, it can be concluded that there is a difference in the average pre-test LP and post-test LP. This research is 3 imilar to our previous research that investigated Speed, Agility and Quickness (SAQ) circuit training to improve kick speed and agility of taekwondo junior athletes [32].

This was seen in the results from the data analysis demonstrating that the average score of the pre-test was 53.36±6.56 while that of the post-test was 62.55±7.48 and tested a significance level of -6.53 with a significant level of p<0.05. The findings in this study are in line with the findings of previous studies [33]. Plyometric training programs had a more positive effect, as compared to a typical training program in terms of increased performance and strength [34], [35]. QST circuit rhythm fast was also a part of plyometric exercises and increased leg power [36], [37]. Resistance training method will be a correction and proven to be no better than power-band resistance training, it has a significant effect on the football player group in the context of physical performance [38].

Although the results showed that the exercise combining strength with speed increased agility and leg power, it was not common for all athletes. Age changes affected the content of the exercise due to changes in the physiological characteristics of the athletes. Therefore, it is necessary to adjust the contents of the training program, especially the volume, intensity, and type of exercise based on the initial condition [39]. QST with low-medium intensity (80%-100% 1 RM), 2-3.

Possible to increase leg power [40]. It is important for everyone to understand the possible advantages and

^{*}QST: Quick Strength Training

disadvantages of several methods of resistance and training. research in this regard has been widely carried out to improve performance and of course as part of efforts to minimize the risk of sports injuries [41].

5. Conclusions

The results of this study showed that QST increased agility and leg power. Fast rhythm strength exercises with a variety of movements improved agility quickly. In addition, QST fast rhythm circuit that is also a part of plyometric exercises increased leg power. Therefore, the QST-CT should be applied during special preparation stages for athletes. QST forms should be adapted to the characteristics of futsal sports.

Although this study showed a positive effect of QST-CT on agility and leg power, there may still be other influencing factors. This is because the sample is limited to only 22 people at the Futsal academy Ligasu-ABAA. Thus, it is necessary to conduct further research by other researchers with a larger sample and other variables.

Acknowledgements

We would like to thank Ligasu-ABAA Futsal Academy Manager, Reza Pranopik, for giving permission and supporting this research. We would also like to extend our appreciation to coaches Bima S, Agung P, and Aan Deki P. who have helped this project as well as the process of retrieving research data. Finally, our gratitude goes to all athletes who participated in the experiments for 6 weeks.

REFERENCES

- T. O. Bompa and G. G. Haff, Periodization: Theory and Methodology of Training. 2009.
- [2] I. Akhmad, A. Supriadi, R. Dewi, and Y. Swara, "The Influence of SAQ Training on Speed and Agility for Futsal Young Athletes on X-Trail 14 Futsal Academy," International Journal of Science and Research, vol. 8, no. 12, pp. 933–936, 2019, doi: 10.21275/ART20203413.
- [3] M. Hammami, Y. Negra, F. Billaut, S. Hermassi, R. J. Shephard, and M. S. Chelly, "Effects of lower-limb strength training on agility, repeated sprinting with changes of direction, leg peak power, and neuromuscular adaptations of soccer players," Journal of Strength and Conditioning Research, vol. 32, no. 1, pp. 37–47, 2018, doi: 10.1519/JSC.0000000000001813.
- [4] D. G. Behm et al., "Effectiveness of traditional strength vs. power training on muscle strength, power and speed with youth: A systematic review and meta-analysis," Frontiers in Physiology, vol. 8, no. JUN, 2017, doi: 10.3389/fphys.2017.00423.

- 5] D. W. Wondale, B. Akhtar, and H. T. Debele, "Effects of core strength training on power, speed and agility on soccer players in Digotsion Town; Northwestern Ethiopia," vol. 7, no. 10, pp. 341–349, 2021.
- [6] D. W. Wondale, B. Akhtar, and H. T. Debele, "Effects of core strength training on power, speed and agility on soccer players in Digotsion Town; Northwestern Ethiopia," International Journal of Innovative Research in Technology, vol. 7, no. 10, pp. 341–349, 2021.
- [7] N. Mohanta, S. Kalra, and S. Pawaria, "A Comparative Study of Circuit Training and Plyometric Training on Strength, Speed and Agility in State Level Lawn Tennis Players," Journal of Clinical and Diagnostic Research, no. February, 2019, doi: 10.7860/jcdr/2019/42431.13348.
- [8] K. Azmi and N. W. Kusnanik, "Effect of Exercise Program Speed, Agility, and Quickness (SAQ) in Improving Speed, Agility, and Acceleration," Journal of Physics: Conference Series, vol. 947, no. 1, p. 012043, Jan. 2018.
- [9] M. Jovanovic, G. Sporis, D. Omrcen, and F. Fiorentini, "Effects of speed, agility, quickness training method on power performance in elite soccer players," Journal of Strength and Conditioning Research, vol. 25, no. 5, pp. 1285-92, 2011. doi: 10.1519/JSC.0b013e3181d67c65.
- [10] D. H. T. A. H. H. Frederick, M.F.A1, 2, K. H., A. H Omar2, 3, and 3 Kamaruzaman Soeed2, 3 and Izwyn Zulkapri2, "Badminton: Specific Movemen Agility Testing System," Journal of Sports Sciences, vol. 27, no. 14, pp. 1591–1599, 2009, doi: 10.1080/02640410903352907.
- [11] C. H. Ooi et al., "Physiological characteristics of elite and sub-elite badminton players," Journal of Sports Sciences, vol. 27, no. 14, pp. 1591–1599, 2009.
- [12] G. Sporis, I. Jukic, L. Milanovic, and V. Vucetic, "Reliability and factorial validity of agility test for soccer players," Journal of Strength and Conditioning Research, vol. 24, no. 3, pp. 679–686, 2010.
- [13] K. Adams, J. P. O'Shea, K. L. O'Shea, and M. Climstein, "The effect of six weeks of squat, plyometric and squat-plyometric training on power production," Journal of Strength and Conditioning Research, vol. 6, no. 1, pp. 36– 41, 1992.
- [14] I. Akhmad and B. S. Hasibuan, "Contribution of SAQ Exercises and Plyometric Exercises Against Smash in Princess Volleyball Games," vol. 23, no. UnICoSS 2019, pp. 201–204, 2020.
- [15] G. Cometti, N. A. Maffiuletti, M. Pousson, J. C. Chatard, and N. Maffulli, "Isokinetic strength and anaerobic power of elite, subelite and amateur French soccer players," International Journal of Sports Medicine, vol. 22, no. 1, pp. 45–51, 2001.
- [16] U. Devrim and K. Erdem, "Evaluation of the Effects of Core-Quick Strength and Core-Plyometric Studies on Balance, Agility and Strength Traits of Volleyball Players," Asian Journal of Education and Training, vol. 5, no. 3, pp. 482–487, 2019, doi: 10.20448/journal.522.2019.53.482.48
- [17] J. Bangsbo, M. Mohr, and P. Krustrup, "Physical and metabolic demands of training and match-play in the elite football player," Journal of Sports Sciences, vol. 24, no. 7, pp. 665–674, 2006.

- [18] M. Alp and B. Baydemir, "The effects of quick strength training on agility performance in soccer," Universal Journal of Educational Research, vol. 7, no. 4, pp. 1001– 1006, 2019.
- [19] H. Jullien, C. C. Bisch, N. L. T, and C. Manouvrier, "Does A Short Period Of Lower Limb Strength Training Improve Performance In Field-Based Tests Of Running And Agility In Young Professional Soccer Players?," Journal of Strength and Conditioning Research, vol. 22, pp. 404–411, 2008.
- [20] H. Jullien, C. Bisch, N. Largouét, C. Manouvrier, C. J. Carlentg, and V. Amiard, "Does a short period of lower limb strength training improve performance in field-based tests of running and agility in young professional soccer players?," Journal of Strength and Conditioning Research, vol. 22, no. 2, pp. 404–411, 2008, doi: 10.1519/JSC.0b013e31816601e5.
- [21] A. Supriadi, I. Akhmad, and R. Dewi, "Design and Development of 'Speed Light' Saq Lighting and Training Equipment Digital Lighting Digitalization Based on Infra Red Motion Sensor," International Journal of Science and Research (IJSR), vol. 7, no. 9, pp. 1194–1196, 2018.
- [22] M. Rogers, N. Tamulevicius, S. Semple, and Z. Krkeljas, "Efficacy of home-based kinesthesia, balance & agility exercise training among.pdf," J Sports Sci Med, vol. 85, no. December, pp. 751–758, 2012.
- [23] Brian T. McCormick, "The Reliability and Validity of Various Lateral Side-Step Tests," IJASS (International Journal of Applied Sports Sciences), vol. 26, no. 2, pp. 67– 75, 2014.
- [24] P. De Salles, F. Vasconcellos, G. De Salles, R. Fonseca, and E. Dantas, "Validity and reproducibility of the sargent jump test in the assessment of explosive strength in soccer players," Journal of Human Kinetics, vol. 33, no. 1, pp. 115–121, 2012.
- [25] M. G. Miller, J. J. Herniman, M. D. Ricard, C. C. Cheatham, and T. J. Michael, "The effects of a 6-week plyometric training program on agility," Journal of Sports Science and Medicine, vol. 5, no. 3, pp. 459–465, 2006.
- [26] M. S. Chelly, M. Fathloun, N. Cherif, M. Ben Amar, Z. Tabka, and E. Van Praagh, "Effects of a back squat training program on leg power.jump, and sprint performances in junior soccer players," Journal of Strength and Conditioning Research, vol. 23, no. 8, pp. 2241–2249, 2009.
- [27] N. S. 6 rahap, S. Suprayitno, N. Simatupang, and R. M. Sari, "Lactic acid level in soccer who consume red dragon fruit juice and regular exercise," Journal of Physics: Conference Series, vol. 1811, no. 1, p. 12059, 2021, doi: 10.1088/1742-6596/1811/1/012059.
- [28] B. R. Ronnestad, N. H. Kvamme, A. Sunde, and T. Raastad, "Short-term effects of strength and plyometric training on sprint and jump performance in professional soccer players," Journal of Strength and Conditioning Research, vol. 22, no. 3, pp. 773–780, 2008.
- [29] M. Christou, I. Smilios, K. Sotiropoulos, K. Volaklis, T. Pilianidis, and S. P. Tokmakidis, "Effects of resistance training on the physical capacities of adolescent soccer players," Journal of Strength and Conditioning Research,

- vol. 20, no. 4, pp. 783-791, 2006.
- [30] K. E. T. Homas and D. U. F. Rench, "The Effect Of Two Plyometric Training Techniques On Muscular Power And Agility In Youth Soccer Players," Journal of Strength and Conditioning Research, vol. 23, pp. 332–335, 2009.
- [31] S. A. I. Shalfawi, T. Haugen, T. A. Jakobsen, E. Enoksen, and E. Tonnessen, "The effect of combined resisted agility and repeated sprint training vs. strength training on female elite soccer players," Journal of Strength and Conditioning Research, vol. 27, no. 11, pp. 2966–2972, 2013.
- [32] I. Akhmad, T. Nugraha, and P. Sembiring, "Speed, Agility, and Quickness (SAQ) training of the circuit system: How does it affect kick speed and agility of junior taekwondo athletes?," Journal Sport Area, vol. 6, no. 2, pp. 175–182, 2021, doi: 10.25299/sportarea.2021.vol6(2).6433.
- [33] M. S. Chelly, M. A. Ghenem, K. Abid, S. Hermassi, Z. Tabka, and R. J. Shephard, "Effects of in-season short-term plyometric training program on leg power, jump-and sprint performance of soccer players," Journal of Strength and Conditioning Research, vol. 24, no. 10, pp. 2670–2676, 2010.
- [34] S. Harmandeep, K. Satinder, R. Amita, and S. Anupriya, "Effects of Six-Week Plyometrics on Vertical Jumping Ability of Volleyball Players," Research Journal of Physical Education Sciences, vol. 3, no. 4, pp. 1–4, 2015.
- [35] L. Czerwosz et al., "T Raining on F Unctional P Erformance in," European journal of medical research, vol. 14, no. 3, pp. 652–657, 2002.
- [36] E. T. Katushabe and M. Kramer, "Effects of combined 7 wer band resistance training on sprint speed, agility, vertical jump height, and strength in collegiate soccer players," International Journal of Exercise Science, vol. 13, no. 4, pp. 950–963, 2020.
- [37] C. E. Anderson, G. A. Sforzo, and J. A. Sigg, "The Effects Of Combining Elastic And Free Weight Resistance On Strength And Power In Athletes Corey," Journal of Strength and Conditioning Research, vol. 22, no. 2, pp. 567–574, 2008.
- [38] K. Wirth, H. Hartmnn, A. Sander, and M. Kiner, "The Impact Of Back Squat And Leg-Press Exercises On Maximal Strength And Speed-Strength Parameters," vol. 30, no. 5, pp. 11–13, 2016.
- [39] D. Rodri'Guez-7sell, F. F.-M. Rquez, and R. Mora-Custodio, "Effect Of High-Speed Strength Training On Physical Performance In Young Soccer Players Of Different Ages," Journal ofStrength and Conditioning Research, vol. 31, pp. 2498–2508, 2017.
- [40] L. Branquinho, R. Ferraz, P. D. Mendes, J. Petricia, J. Serrano, and M. C. Marques, "The effect of an in-season 8-week plyometric training programme followed by a detraining period on explosive skills in competitive junior soccer players," Montenegrin Journal of Sports Science and Medicine, vol. 9, no. 1, pp. 33–40, 2020.
- [41] E. T. Katushabe and M. Kramer, "Effects of combined 7 wer band resistance training on sprint speed, agility, vertical jump height, and strength in collegiate soccer players," International Journal of Exercise Science, vol. 13, no. 4, pp. 950–963, 2020.

SAJ2	27-199298	12.pui				
ORIGINALITY REPORT						
SIMILA	2% ARITY INDEX	11% INTERNET SOURCES	12% PUBLICATIONS	% STUDENT PAPERS		
PRIMAR	Y SOURCES					
1	digilib.ur	nimed.ac.id		2%		
2	rie.binac Internet Sourc	darma.ac.id		1 %		
3	journal.u Internet Sourc			1 %		
4	Aaron Uthoff, Jon Oliver, John Cronin, Craig Harrison, Paul Winwood. "Sprint-Specific Training in Youth", Journal of Strength and Conditioning Research, 2020 Publication					
5	Islahuzzaman Nuryadin, Rumi Iqbal Doewes. "The Effect of Playing Approach Learning Models on Football Dribbling Skills", Proceedings of the 4th International Conference on Learning Innovation and Quality Education, 2020 Publication					

6 www.coursehero.com
Internet Source

1 %

training improves kicking motion motor performance", Physical Therapy in Sport, 2014

Publication

14

worldwidescience.org

Internet Source

%

Exclude quotes Off
Exclude bibliography Off

Exclude matches

< 1%