


Volume 10, Number 2
August 2021
ISSN (Print): 2089-7790
ISSN (Online): 2502-6194

DEPIK

Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan
(Journal of Aquatic, Coastal and Fishery Sciences)



DEPIK

Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan (Journal of Aquatic, Coastal and Fishery Sciences)

ISSN (Print): 2089-7790; ISSN (Online): 2502-6194

Publisher : Faculty of Marine and Fisheries, Universitas Syiah Kuala, Indonesia
Editor in Chief : Ichsan Setiawan, M.Si, Universitas Syiah Kuala (Oceanology/Oceanography)
Managing Editor : Dr. Nur Fadli, M.Sc, Universitas Syiah Kuala (Marine Biology)
Associate Editors : Haekal Azief Haridhi, S.Kel, M.Sc, Ph.D, Universitas Syiah Kuala (Earth Science)
Dr. Yudi Haditiar, S.Kel, Universitas Syiah Kuala (Marine Sciences)
Dedi F. Putra, M.Sc, Universitas Syiah Kuala (Aquatic Parasitology)
Adrian Damora, M.Si, Universitas Syiah Kuala (Fisheries Management)
Djamani Rianjuanda, M.Si, Universitas Syiah Kuala (Capture Fisheries)
Teuku Haris Iqbal, M.Sc, Universitas Syiah Kuala (Marine Technology)

Editor Board :

Prof. Dr. Zainal Abidin Muchlisin, Ichthyology, Universitas Syiah Kuala, Indonesia
Prof. Dr. Syamsul Rizal, Physical Oceanography, Universitas Syiah Kuala, Indonesia
Prof. Dr. Musri Musman, Natural Product Chemistry, Universitas Syiah Kuala, Indonesia
Prof. Dr. Adlim, M. Sc, Fish Processing and Nano Particles, Universitas Syiah Kuala, Indonesia
Prof. Dr. Mohd Nor Siti-Azizah, Fish Genetic, Universiti Sains Malaysia, Malaysia
Prof. Dr. Ahmad J. Nayaya, Biological Sciences, Abubakar Tafawa Balewa University, Nigeria
Prof. Dr. Muhammad Ali Sarong, Aquatic Ecology, Universitas Syiah Kuala, Indonesia
Dr. Eriyusni Eriyusni, Fish Reproduction, Universitas Sumatera Utara, Indonesia
Dr. Indra Suharman, Fish Nutrition, University Riau, Indonesia
Dr. Agung Setia Batubara, Aquaculture, Universitas Negeri Medan, Indonesia

Graphical Design : Firman M. Nur, S.Si

Finance & Circulation : Muhammad Saumi, A.Md

Publisher Head : Doktorandus Muhammad, M.Si

Depik Ilmu-Ilmu Perairan, Pesisir dan Perikanan (Journal of Aquatic, Coastal and Fishery Sciences) is indexed by



Mailing Address:

Faculty of Marine and Fisheries, Universitas Syiah Kuala
Kopelma Darussalam, Banda Aceh 23111, Aceh Province, Indonesia

Website: <http://jurnal.unsyiah.ac.id/depik>

Principal Contact: +62-(0)852-2018-9228, email: ichsansetiawan@unsyiah.ac.id

Support Contact: +62-(0)813-6003-4100, email: nurfadli@unsyiah.ac.id



Table of Contents

ARTICLES

FRONT MATTER PDF

...

In vivo test of *Litopenaeus vannamei* infected by *Vibrio* using *Moringa oleifera* leaf extract PDF
Iko Imelda Arisa, Cut Mutia, Sri Agustina, Nurfadillah Nurfadillah, Sofyatuddin Karina 86-90

[doi>](#) 10.13170/depik.10.2.17510

Determination of potential fisheries areas based on trophic status (Case study in Situ Gede, Tasikmalaya) PDF
Sevi Sawestri, Ni Komang Suryati, Dina Muthmainnah 91-97

[doi>](#) 10.13170/depik.10.2.20177

Survival and growth of *Pangasianodon hypophthalmus* cultured under controlled photoperiod PDF
Cyntia Uli Artha Sihombing, Muhammad Fauzi, Windarti Windarti 98-102

[doi>](#) 10.13170/depik.10.2.20053

Effect of transplantation media on *Pocillopora* coral growth rate at TWAL Pulau Weh PDF
Chitra Octavina, Maria Ulfah, Adrian Damora, Zulkarnain Jalil, Nanda Muhammad Razi, Muhammad Agustiar, Puad Batari Harahap, Nurul Najmi, Samsul Bahri, Munandar Munandar, Shan-Yin Vanson Liu 103-106

[doi>](#) 10.13170/depik.10.2.19222

Practical application of sea urchin shell flour supplementation as a stimulant moulting in *vannamei* shrimp PDF
Heriansah Heriansah, Nursyahrhan Nursyahrhan, Nursidi Nursidi, Nur Fajriani Nursida, Najamuddin Najamuddin 107-114

[doi>](#) 10.13170/depik.10.2.19456

Condition of coral reef in Batee Island waters, Peukan Bada Sub-District, Aceh Besar PDF
Maria Ulfah, Saiful Mahlil, Muhammad Nasir, Sayyid Afdhal El Rahimi, Syahrul Purnawan, Muhammad Rizki Fazillah 115-119

[doi>](#) 10.13170/depik.10.2.19995

Post feeding protection from Bilimbi to Motile *Aeromonas* Septicemia disease in African Catfish PDF
Sefti Heza Dwinanti, Gusti Setiawan, Ade Dewi Sasanti, Madyasta Anggana Rarassari 120-124

[doi>](#) 10.13170/depik.10.2.20445

Analysis of catches data collection system at Fish Landing Base (PPI) of Dumai PDF
Ratu Sari Mardiah, Sri Yenica Roza, Edy Miswar 125-129

[doi>](#) 10.13170/depik.10.2.18565

Observation and numerical modeling of physical oceanography in the Balikpapan Bay, East Kalimantan: Preliminary results PDF
130-135

ABOUT THE JOURNAL:

[Aims and Scope](#)

[Editorial Team](#)

[Reviewer Team](#)

FOR AUTHORS:

[Guideline for Author](#)

[Publication Ethics](#)

[Google Scholar Citation](#)

[Scopus Citation](#)

[Indexed By](#)

[Article Publication Charges](#)

ONLINE SUBMISSION:

[SUBMIT](#)

PAPER TEMPLATE:



Visitors

	ID 282,163		PL 61
	US 16,113		IR 61
	MY 915		PK 53
	JP 807		NG 46
	IN 802		MX 44
	SG 765		PT 38
	CN 575		IL 38
	NL 377		BD 35
	GB 333		CZ 35
	RU 300		SE 30
	TH 293		IT 29
	DE 281		CO 28
	KR 273		KH 26
	AU 234		RO 25
	CA 162		BE 24
	TW 159		EG 24
	PH 148		SA 23
	FR 148		ES 22
	TR 147		FI 21
	ZA 116		PE 19
	TL 113		GR 18
	BR 108		CL 17
	HK 97		UA 14
	IE 71		BN 13
	VN 65		CH 13

Pageviews: 856,044
Flags Collected: 130



Mutiara Rachmat Putri, Iwan P Anwar, Zetsaona Sihotang, Lamona I. Bernawis, Agus Setiawan, Muhammad Riza, Idris Mandang, Willem M. Tatipatta

[doi> 10.13170/depik.10.2.19259](https://doi.org/10.13170/depik.10.2.19259)

Heavy metal content in pumpkin pond of Kutaraja Fishing Port Banda Aceh

PDF

Muhammad Muhammad, Shelly Anggi Pratiwi, Thaib Rizwan, Zulkarnain Jalil

136-142

[doi> 10.13170/depik.10.2.20943](https://doi.org/10.13170/depik.10.2.20943)

Anesthetic effectiveness of tekelan leaves on tinfoil barb fish in closed transportation

PDF

Eva Ayuzar, Rachmawati Rusydi, Hayatun Muna, Muliani Muliani

143-150

[doi> 10.13170/depik.10.2.18519](https://doi.org/10.13170/depik.10.2.18519)

Characteristics of catfish oil, red palm oil and shark liver oil as functional foods

PDF

Ulil Amri, Andarini Diharmi, Mery Sukmiwati

151-160

[doi> 10.13170/depik.10.2.19131](https://doi.org/10.13170/depik.10.2.19131)

Analysis of the utilization of landsat 8 oli imagery for mapping the distribution of coral reefs in Pulau Weh Sabang

PDF

M Taufik Rahmadi, Sugiharto Sugiharto, Mbina Pinem, Ayu Suciani

161-166

[doi> 10.13170/depik.10.2.20966](https://doi.org/10.13170/depik.10.2.20966)

Fishing season of large pelagic fish in Idi Rayeuk waters, East Aceh, Indonesia

PDF

M.A. Chaliluddin, Rizka Alfita, Thaib Rizwan, Rahmat Rizqi, Rosi Rahayu, Sayyid Afdhal El Rahimi, Ichsan Rusydi

167-173

[doi> 10.13170/depik.10.2.21203](https://doi.org/10.13170/depik.10.2.21203)

The diversity of plankton as bioindicators in Kakap River Estuary, West Kalimantan

PDF

Widadi Padmarsari Soetignya, Patrisia Marniati, Mardan Adjijaya, Yunita Magrima Anzani

174-179

[doi> 10.13170/depik.10.2.21303](https://doi.org/10.13170/depik.10.2.21303)

Feed efficiency with the addition of probiotics for asang fish fry, *Osteochilus vittatus*, CV

PDF

Usman Bulanin, Yuneidi Basri, Sri Puja Lestari

180-185

[doi> 10.13170/depik.10.2.20168](https://doi.org/10.13170/depik.10.2.20168)

Grouper DNA barcoding studies in Indonesia: A short review

PDF

Nanda Muhammad Razi, Zainal A. Muchlisin, Siti Maulida, Mutia Ramadhaniaty, Firman M. Nur, Adrian Damora, Sumarni Laila Buang Manalu, Nur Fadli

186-193

[doi> 10.13170/depik.10.2.21255](https://doi.org/10.13170/depik.10.2.21255)

Evaluating the addition of starry triggerfish (*Abalistes stellaris*) bone charcoal as a feed supplement to the growth performance and intestinal villi length of Nile tilapia (*Oreochromis niloticus*)

PDF

194-200

Azwar Thaib, Lia Handayani, Ardila Hanum, Nurhayati Nurhayati, Faisal Syahputra

[doi> 10.13170/depik.10.2.20367](https://doi.org/10.13170/depik.10.2.20367)

BACK MATTER

PDF

...

00637803

[View My Stats](#)

CITATIONS AND H-INDEX OF DEPIK IN GOOGLE SCHOLAR: [\(Click Here\)](#)

DEPIK JURNAL ILMU ILMU PERAIRAN, PESISIR, DAN PERIKANAN IS INDEXED BY

DOAJ DIRECTORY OF OPEN ACCESS JOURNALS

 GARUDA GARSA RUJUKAN DIGITAL



 Crossref

 SINTA Science and Technology Index

SINTA - 2



OPEN JOURNAL SYSTEMS

User

Username

Password

Remember me



Better Writing Made Easy





Analysis of the utilization of landsat 8 oli imagery for mapping the distribution of coral reefs in Pulau Weh Sabang

M Taufik Rahmadi^{1*}, Sugiharto Sugiharto¹, Mbina Pinem¹, Ayu Suciani²

¹Jurusan Pendidikan Geografi, Fakultas Ilmu Sosial, Universitas Negeri Medan, Indonesia.

²Program Studi Pendidikan Geografi, Fakultas Keguruan dan Ilmu Pendidikan, Universitas Samudra.

ARTICLE INFO

ABSTRACT

Keywords:

Landsat 8 oli
Mapping
Distribution
Coral reef
Weh Island

DOI: [10.13170/depik.10.2.20966](https://doi.org/10.13170/depik.10.2.20966)

As one of the largest archipelagic countries globally, Indonesia has diverse natural resources, one of which is the coral reef ecosystem. Coral reef ecosystems are spread across almost all Indonesian waters, and Pulau Weh Sabang is one distribution area. This study aims to determine the distribution of coral reef ecosystems and test landsat 8 oli imagery accuracy in mapping coral reef ecosystems. The method used in this research is the nearest neighbour algorithm object-based classification method. The results showed that the coral reef ecosystem in Pulau Weh Sabang was divided into two classes: a healthy coral reef ecosystem class with 277.38 hectares and a medium condition coral reef ecosystem class with an area of 710.01 Ha.

Introduction

Indonesia is one of the countries that have abundant natural resources. As one of the largest archipelagic countries and significant natural resources, the Indonesian archipelago has various marine ecosystems, both those that live in shallow seas and those that live in the deep sea. Dahuri (2003) suggests that coastal ecosystems' characteristics can be classified into artificial ecosystems and natural ecosystems. One of the natural ecosystems located in the coastal zone is the coral reef ecosystem. Koroy (2020) argues that coral reefs have distributed throughout Indonesia's territorial waters with varying conditions, both in type and quantity. According to Tangke (2010), coastal ecosystems generally consist of three components: sea, coral reefs, and mangroves.

Nybakken (2001) suggests that coral reef ecosystems are shallow marine organisms whose organic productivity is very high compared to other organisms and is typically found in tropical areas. The increased productivity of coral reef ecosystems is because the coral reef ecosystem has various benefits

for the surrounding ecosystem and human life, such as fish spawning grounds, habitat and food sources, coastal protection, and carbon sinks. The coral reef ecosystem also has economic benefits as a marine tourism attraction. The existence of coral reef ecosystems is spread across almost all coastal areas of Indonesia with various conditions and characteristics. According to Hadi *et al.* (2018), based on observations at 1067 points, Indonesia's coral reefs are divided into several conditions: poor condition by 36.18%, moderate condition by 34.3%, and good condition 22.96%, and excellent condition by 6.56%. Muhsoni (2011) stated that human activities threaten 88% of coral reefs threatening biological and economic values that are very important for human life. 50% of the endangered coral reefs are at very high threat levels, only 12% are at low threat levels. According to Lutfi and Anugrah (2017), the condition of coral reefs is currently experiencing damage and decline caused by various activities, including bombing fish, fishing using toxic materials, illegal trading of ornamental corals, and climate change. Several studies on the damage to

* Corresponding author.

Email address: taufikrahmadi@unimed.ac.id

coral reef ecosystems due to human activities, among others, were put forward by Uar et al. (2016); several human activities that damage coral reefs are fishing using fish bombs, arrows, nets, traps, taking coral for decoration and building materials. Meanwhile, research conducted by Jubaedah and Anas (2019) stated that marine tourism in the Nusa Penida marine conservation area impacted reducing the scope of coral reefs by 4%.

One of the areas in Indonesia that has a coral reef ecosystem is Pulau Weh Sabang. Pulau Weh is a marine conservation area managed by the Traditional Laot Institution system following the Sabang Mayor Number 729 of 2010 concerning the Sabang Island Coastal Marine Protected Area's Reservation. According to Hastuty (2014), the eastern coastal area of Pulau Weh is unique in its management system, namely using local wisdom (customary law), which had been implemented long before this area was designated as a conservation area.

Coral reef ecosystems must be well managed, and following the conditions of life that can be tolerated by coral reef ecosystems such as salinity levels, water temperature, pH, and human activity factors, this is because coral reef ecosystems have very little growth every year (Dahuri et al., 1996). The low growth rate of coral reef ecosystems and the high human need for coral reef ecosystems today and in the future requires a method that is relevant, effective, and efficient in managing coral reef ecosystems so that that coral reef ecosystems can be managed sustainable, one of which is using remote sensing technology.

The development of remote sensing technology to manage coral reef ecosystems is inseparable from technology's rapid growth. Currently, remote sensing technology for coral reef ecosystems varies significantly with the availability of low-resolution imagery to high-resolution imagery. Coral reef management can be carried out sustainably. According to Sutanto (1992), remote sensing images have advantages, including pictures depicting objects, areas, and symptoms on the earth's surface following the shape of the original object. The use of remote sensing technology is very effective in various aspects such as cost efficiency, time, and human resources to monitor changes and management of natural resources, incredibly natural resources in shallow waters and coastal areas (Mumby et al. 1999; Green et al. 2000; Maeder et al. 2002; Lillesand et al. 2004; Zhigang et al. 2008; Tamondong et al. 2013 in Fahriansyah et al. 2017).

Research on the use of remote sensing images with various resolutions for shallow water ecosystems has been carried out before, including

Hafizt et al. (2017) with the title study of the LANDSAT 8 OLI image classification method for benthic habitat mapping in the Padaido Islands, Papua with image accuracy results of 47.57% for multispectral classification and 36.17% for the object-based category. Nababan et al. (2018), with the title research object-based benthic habitat mapping using sentinel images in the waters of Wangi-Wangi Island, Wakatobi Regency with the accuracy of sentinel-2 images using object-based classification, which has an accuracy value of 60.4% for 12 benthic habitat classes and 64.1% for nine benthic habitat classes.

Based on previous research, this study aims to map the distribution of coral reef ecosystems on Weh Sabang Island and test the accuracy of LANDSAT 8 OLI (Operational Land Imager) imagery in mapping the distribution of coral reefs. This study produces data that can be used as a reference for the management of coral reef ecosystems on Weh Sabang Island. The condition of the coral reef ecosystem on Weh Sabang Island can be preserved and can be used for economic development in the management of sustainable marine tourism. This research must consider that coral reefs will easily change due to various natural factors and human activity factors.

Materials and Methods

Location and time of research

This research was conducted from August - October 2020. Geographically, Weh Sabang Island is located between 95°12'00" - 95°16'00" BT and 5°51'00" - 5°55'00" LU. Weh Sabang Island is part of the Sabang City administration, with a land area of 121.7 km² and a water area of 920.05 km². From a geographical perspective of Indonesia, the Weh Sabang Island region is the westernmost administrative area and directly borders neighbour countries, namely Malaysia, Thailand, and India (Badan Pusat Statistik, 2019). The coast of Pulau Weh Sabang is a Marine Conservation Area which was determined based on the decision of the Minister of Marine Affairs and Fisheries of the Republic of Indonesia Number 57/KEPMEN-KP/2013. For more details, the research location can be seen in Figure 1.

Research tools and materials

The tools used in this study consisted of ENVI 5.1 software, e-Cognition 9.0 software, eutech instruments, handheld reference meters, Global Positioning System (GPS), and cameras. Meanwhile, the research materials used included LANDSAT 8 OLI (Operational Land Imager) imagery recorded on

June 25, 2020, a 1: 50,000 scale RBI map of Pulau Weh Sabang (Sheet number 0421-53 and 0421-54), and coral transect data from field surveys.

Research stages

The research stages consisted of literature studies, image data acquisition (USGS Earth Explore), digital image processing, coral reef distribution analysis, and field surveys. The field survey aims to directly see whether the condition of coral reefs follows the results of a tentative map and to determine the factors of damage to coral reefs caused by natural and human factors. The biological factor in this study is the condition of the waters of Weh Island, Sabang. The image processing stages used in the study consist of (1) image correction (geometric and radiometric), (2) masking and classification, and (3) field survey (based on the tentative map). For more details, the stages of the research can be seen in Figure 2.

This study uses image data recorded in June 2020. The LANDSAT 8 OLI image is an L1-T (level one terrain corrected) image that has been free from sensor errors, so there is no need for geometric corrections anymore. Meanwhile, the radiometric corrections carried out include converting the Digital Number (DN) value to the radian spectral value, converting the radian spectral value to the reflectance spectral value, Fast Line of sight Atmospheric Analysis of Spectral Hypercubes (FLAASH) correction, Sun glint correction, and Water Column correction. The technique used in image masking is a visual technique based on Pulau Weh Sabang's administrative area. Image classification used in this study is using the object-based segmentation and classification (OBIA) nearest neighbour algorithm. According to Simamora et al. (2015), OBIA classification is a classification approach with advantages over other classifications because OBIA classifies by considering the object's spatial aspects.

The field survey was conducted in August-October 2020 based on a tentative map of LANDSAT 8 OLI image data processing recorded in 2020. This research field survey used the photo transect method developed by Roelfsema and Phinn (2009). This method is effective because it records all objects at the sampling point in the form of photographs. Coral reef information will be obtained in qualitative and quantitative states, such as the percentage of cover, composition, and coral reef ecosystem conditions.




Figure 1. Research location map.




Figure 2. Framework for the stages of coral reef distribution research.

Results

Distribution of coral reefs

Mapping of coral reef ecosystem distribution is obtained based on the nearest neighbour algorithm object-based classification. The object-based classification process (OBIA) consists of image segmentation and image classification resulting from segmentation. Mapping the distribution of coral reef ecosystems in object-based classification begins with the image segmentation process and segmentation sampling for each object of the coral reef ecosystem.

The distribution of coral reefs was obtained by analyzing the results of image classification and field surveys. For more details, the distribution of the coral reef ecosystem on Pulau Weh Sabang in 2020 can be seen in Table 1 and Figure 3.

Table 1. Distribution of coral reef ecosystem extensions in Pulau Weh Sabang.

No	Coral Reef Class	Ecosystem Area (Ha)	Percentage (%)
1.	Healthy Coral Reef Ecosystem	277,38	28,10
2.	Medium Coral Reef Ecosystem	710,01	71,90
	Amount	987,39	100

Source: Data Processing (2020).




Figure 3. Map of distribution of coral reefs ecosystems in Pulau Weh Sabang 2020.

Image classification

The results of the object-based classification, the distribution of coral reef ecosystems on Weh Sabang Island, were only classified into two classes, namely the healthy condition coral reef ecosystem class with an area of 277.38 Ha (28.10%) and the coral reef ecosystem class in medium condition with an area of 710.01 Ha (71.90%). Based on image classification results, there are sand objects that dominate the distribution of coral reefs. The number of sand objects is due to damaged and dead coral reefs, so the classification results are detected as sand. For more details, the coral reefs on Pulau Weh Sabang can be seen in Figure 4 and 5.

Water conditions

Based on the Pulau Weh area's field survey data, the water temperature ranges from 29.6°C-30.9°C, the salinity level varies from 33.17‰-32.96‰, and the pH ranges from 8.44-8.56. Based on these data, Pulau Weh Sabang's waters have a temperature greater than 0.9°C and a pH level greater than 0.6 than the tolerance value that the coral reef ecosystem can accept. Changes in the composition of benthic

habitats can occur due to various sustainability factors, such as human activities and natural disasters. This is because the habitat benthic have different tolerance limits to sustainability factors. Some biological factors influence changes in the composition of the benthic habitat between temperature, salinity and other pH levels outside benthic habitat tolerance limits. While characteristics of human activity that damage in the form of underwater tourism activities that are not taking into account the conditions of benthic habitats and ship fuel disposal (Rahmadi, 2017)

Image accuracy test

The accuracy test is the final stage to determine the accuracy of the distribution map of coral reef ecosystems. The accuracy test is carried out on the corrected image using a field survey sample. In this study, the accuracy-test used an error matrix table (Confusion matrix). Based on image classification and data to the field, the accuracy-test results of the distribution mapping of coral reef ecosystems in Pulau Weh Sabang are 57%. The low accuracy of mapping the distribution of coral reef ecosystems occurs because of the low quality of images and many atmospheric disturbances.

Discussion

Based on data from the classification results of the image classification of the coral reef ecosystem class on Weh Sabang Island, the coral reef area is equal to 987.39 Ha which is divided into two categories, namely a healthy class coral reef ecosystem of 277.38 Ha and a medium-class coral reef ecosystem of 710.01 Ha. The low classification level of the coral reef ecosystem class is due to several factors, namely the quality of the image, which has many atmospheric effects, and the sunglint effect. In classifying the image, data can only detect two objects of coral reef class. According to Danoedoro (2012), the accuracy of satellite image data processing accuracy is determined by the image resolution used.

The results of previous research conducted by Souisa and Makailipessy (2017) on the waters of the Tayando District, the LANDSAT 8 OLI image was only able to detect the bottom substrate of the waters consisting of fine sand, sand mixed with dead coral debris, dead coral, coral reefs, and seagrass. In another study conducted by Fadhli and Pin (2018) in the waters of Karawang in the LANDSAT 8 OLI image, there is a difference between the results of the mapping and the field survey where in the field survey there are 11 clusters of coral reefs, while the mapping results only classify nine groups of coral reefs.



Figure 4. Condition of Healthy Coral Reef in Pulau Weh Sabang



Figure 5. Condition of Medium Coral Reefs in Pulau Weh Sabang

The coral reef ecosystem on Pulau Weh Sabang is scattered in almost all of Weh Island's water areas, with different conditions for each distribution. The distribution of coral reef ecosystems in a healthy state is dominant in the regions that are the center of coral reef conservation, namely, Rubiah Island waters, Gampong Iboih waters, Gampong Gapang waters, Kasih coastal waters, elephant tread coastal waters, and the southern part of Weh Island. Meanwhile, the distribution of coral reefs is almost evenly distributed throughout the waters of Pulau Weh Sabang. The implementation of the marine conservation area (Watershed Nature Reserve) of Pulau Weh Sabang was determined based on the Decree of the Minister of Marine Affairs and Fisheries of the Republic of Indonesia Number 57/KEPMEN-KP/2013.

According to [Rusman et al. \(2017\)](#), coral reefs on Weh Island, Sabang consist of 11 species, namely *Acropora Cytherea*, *Acropora digitifera*, *Acropora Humilis*, *Acropora Hyacinthus*, *Acropora Rosaria*, *Favites Abdita*, *Favites Acuticolis*, *Goniastrea Pectinata*, *Montipora SP*, *Pocillopora Verrucosa*, and *Porites australiensis*.

The coral reef is an ecosystem that has high productivity and is very sensitive to water conditions. Coral reef growth is determined by good water conditions and is below the tolerable limits of coral reefs. According to [Wisha et al. \(2019\)](#), Weh Island is one of the outer islands of the Indonesian Territory, which is directly adjacent to the Indian Ocean. Water circulation that occurs in the waters of Pulau Weh is influenced by processes that occur in the Indian Ocean, such as the monsoon cycle and the Indian Ocean Dipole, thus affecting sea surface temperature that occurs in the waters of Pulau Weh. [Marshall et al. \(2012\)](#) suggest that coral reefs have acceptable tolerance limits for their survival. If they are excessive

or deficient for an extended period, it will result in coral bleaching. According to the [Decree of the Minister of the Environment \(2014\)](#), coral reefs have temperature tolerance limits between 28°C-30°C, salinity ranges from 32-35‰, and pH ranges from 7-8.5. [Maududi and Lutfi \(2018\)](#) stated that the quality of the water parameters greatly determines the survival of biota in each marine ecosystem. Temperature changes can affect coral reefs' survival process that involves chemical, physical, and biological processes in water bodies. Changes in temperature of 1°C-2°C will put pressure on coral reefs to turn white for a long time.

Meanwhile, the high pH value in the waters of Pulau Weh Sabang is due to the high concentration of carbon dioxide in the atmosphere, where the carbon dioxide released by the sea returns to the atmosphere. The absorption of carbon dioxide in the oceans depends on its concentration in the atmosphere. It is closely related to sea surface temperature, currents, and the level of biological activity in the oceans ([Marshall et al., 2012](#)).

The accuracy test is the final stage to determine the accuracy of the distribution map of coral reef ecosystems. The accuracy test is carried out on the corrected image using a field survey sample. In this study, the accuracy-test used an error matrix table (confusion matrix). Based on image classification and data to the field, the accuracy-test results of the distribution mapping of coral reef ecosystems in Pulau Weh Sabang are 57%. The low level of accuracy of mapping the distribution of coral reef ecosystems occurs because of the many atmospheric disturbances. [Green et al. \(2000\)](#) suggest that remote sensing image data accuracy that can be used as a reference for further research is >60%. The low level

of accuracy obtained in the final results of this study is due to several things, namely the number of coral reef classes, the number of observation points made, and the unfavourable condition of the image data.

The LANDSAT 8 OLI imagery in this study has not detected coral reefs correctly and provides maximum classification results due to several things, namely poor image quality (numerous atmospheric effects during recording) and low resolution (30 meters).

Conclusion

Based on the results of the study, it can be concluded that the distribution of coral reef ecosystems in Pulau Weh Sabang is divided into two classes, namely a healthy coral reef ecosystem class of 277.38 Ha (28.10%) and a moderate condition coral reef ecosystem class of 710.01 Ha (71.90%). The accuracy test for LANDSAT 8 OLI imagery for mapping the distribution of coral reef ecosystems in Pulau Weh Sabang is 57%.

The results of this study still need further research to determine the level of accuracy of the LANDSAT 8 OLI image data in the development of coral reefs on Weh Sabang Island by using different methods and extraction of image data so that that image capabilities can be seen and provide additional accuracy and classification results. in the coral reef

References

Badan Pusat Statistik. 2019. Kota Sabang dalam angka tahun 2018. BPS, Kota Sabang.

Dahuri, R, J. Rais, S.P. Ginting, D.M. Sitepu. 1996. Pengelolaan sumber daya wilayah pesisir secara terpadu. PT. Pradnya Paramita, Jakarta.

Dahuri, R. 2003. Keanekaragaman hayati laut: aset pembangunan berkelanjutan Indonesia. Gramedia Pustaka Utama, Jakarta.

Danoedoro, P. 2012. Pengantar penginderaan jauh digital. Penerbit Andi, Yogyakarta.

Fadhli, R., T.G. Pin. 2018. Persebaran terumbu karang di wilayah perairan Karawang. *Jurnal Geografi Lingkungan Tropik*, 2(1): 38-51.

Fahriansyah, F., J.L. Gaol, J.P. Panjaitan. 2017. Pemetaan Geomorfologi Terumbu Karang Pulau Tunda Menggunakan Klasifikasi Berbasis Objek. *Jurnal Teknologi Perikanan dan Kelautan*, 8(2): 147-156.

Green, E., P. Mumby, A. Edwards, C. Clark. 2000. Remote sensing handbook for tropical coastal management. The United Nations Educational, Scientific and Cultural Organization, Paris.

Hadi, T.A, B.P. Giyanto. 2018. Status terumbu karang Indonesia 2018. Pusat Penelitian Oseanografi, Lembaga Ilmu Pengetahuan Indonesia, Jakarta.

Hafizt, M., M.Y. Iswari, B. Prayudha. 2017. Kajian metode klasifikasi citra landsat-8 untuk pemetaan habitat bentik di Kepulauan Padaido, Papua. *OLDI (Oseanologi dan Limnologi di Indonesia)*, 2(1): 1-13.

Hastuty, R., L. Adrianto. 2014. Tutupan karang dan komposisi ikan karang didalam dan luar kawasan konservasi pesisir timur Pulau Weh, Sabang. *DEPIK Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 3(2): 99-107.

Jubaedah, I, P. Anas. 2019. Dampak pariwisata bahari terhadap ekosistem terumbu karang di Perairan Nusa Penida, Bali. *Jurnal Penyuluhan Perikanan dan Kelautan*, 13(1): 59-75.

Koroy, K., D. Alwi, N.G. Paraisu. 2020. Pengaruh laju sedimentasi terhadap tutupan terumbu karang di perairan Kota Daruba, Kabupaten Pulau Morotai. *DEPIK Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 9(2): 193-199.

Luthfi, O.M, P.T. Anugrah. 2017. Distribusi karang keras (Scleractinia) sebagai penyusun utama ekosistem terumbu karang di Gosong Karang Pakiman, Pulau Bawean. *DEPIK Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 6(1): 9-22.

Marshall, N.J., D. A. Kleine, A.J. Dean. 2012. Coral watch: education, monitoring, and sustainability through citizen science. *Frontiers in Ecology and the Environment*, 10(6): 332-334.

Maududi, M.A, O.M. Luthfi. 2018. Tutupan makroalga pada terumbu karang di kawasan konservasi perairan (KKP) Nusa Penida, Bali. *DEPIK Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 7(1): 69-75.

Muhsoni, F.F. 2011. Pemetaan terumbu karang menggunakan Citra ALOS di Pulau Kangean Kabupaten Sumenep. *Embryo*, 8(1): 216-188.

Nababan, B, J.P. Panjaitan. 2018. Pemetaan habitat bentik berbasis objek menggunakan citra sentinel-2 di Perairan Pulau Wangi-Wangi Kabupaten Wakatobi. *Jurnal Ilmu dan Teknologi Kelautan Tropis*, 10(2), 381-396.

Nybakken, J.W. 2001. *Marine biology: an ecological approach* (Vol. 5). Benjamin Cummings, San Francisco.

Peraturan Menteri Lingkungan Hidup Republik Indonesia Nomor 5 Tahun 2014. Baku Mutu Air Limbah.

Rahmadi, M.T. 2017. Identifikasi faktor-faktor yang mempengaruhi perubahan habitat bentik di sebagian Pulau Weh tahun 2010 dan 2015. *Publikauma Jurnal Administrasi Publik Universitas Medan Area*, 5(2): 1-5.

Roelfsema, C.M, S.R, Phinn. 2009. A manual for conducting georeferenced photo transects surveys to assess the benthos of coral reef and seagrass habitats version 3.0. University of Queensland Brisbane, Australia.

Rusman, F, L. Utami, Y. Wahyuni, N. Falah. 2017. Jenis-jenis terumbu karang di Pulau Rubiah Kota Sabang. *Prosiding Seminar Nasional Biotik*, 5(1): 153-157.

Simamora, F.B, B. Sasmito, H. Haniah. 2015. Kajian metode segmentasi untuk identifikasi tutupan lahan dan luas bidang tanah menggunakan citra pada google earth (studi kasus: Kecamatan Tembalang, Semarang). *Jurnal Geodesi Undip*, 4(4): 43-51.

Souisa, F., M. M. Makailipessy. 2017. Pemetaan substrat dasar perairan dangkal di Kecamatan Tayando Kota Tual Menggunakan Citra Landsat 8. *Neritic*, 6(1): 1-6.

Sutanto. 1992. *Penginderaan Jauh*. Gadjah Mada Press, Yogyakarta.

Tangke, U. 2010. Ekosistem padang lamun (manfaat, fungsi dan rehabilitasi). *Agrikan: Jurnal Agribisnis Perikanan*, 3(1): 9-29.

Uar, N.D, S.H. Murti, S. Hadisusanto. 2016. Kerusakan Lingkungan akibat aktivitas manusia pada ekosistem terumbu karang. *Majalah Geografi Indonesia*, 30(1): 88-96.

Wisha, U.J, T.A, Tanto, N.N.H. Ridwan, R. Dhiuddin. 2019. Dampak fluktuasi suhu permukaan laut terhadap kematian karang di Perairan Pulau Weh, Indonesia. *Jurnal Kelautan Nasional*, 14(2): 103-112.

How to cite this paper:

Rahmadi, M.T., S. Sugiharto, M. Pinem, A. Suciani. 2021. Analysis of the utilization of landsat 8 oli imagery for mapping the distribution of coral reefs in Pulau Weh Sabang. *Depik Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 10(2): 161-166.