

KATA PENGANTAR

Pembaca yang budiman di seluruh tanah air, dengan mengucap syukur dan terima kasih kepada Tuhan Yang Maha Esa, rasa bangga dan dan Salam kimiawan, Jurnal Pendidikan Kimia Volume 7 Nomor 1 edisi April 2015 ini baik edisi cetak maupun on-line di internet, kami sajikan sebagai pembuka komunikasi kita di awal kwartal 2015. Edisi yang diterbitkan mengkombinasikan 13 buah artikel hasil penelitian dalam bidang kimia dan pembelajarannya, serta peneliti kimia dan terapannya khususnya dalam bidang kesehatan, lingkungan dan Biomedik.

Melalui kombinasi jenis penelitian yang disajikan dengan oriijinalitas yang dipercaya, begitupula penulisnya yang beragam profesi dan latar belakang instansi, bahkan berasal dari beberapa provinsi di Indonesia, jurnal yang kami hadirkan ini akan semakin menambah ketertarikan dan minat serta motivasi siswa, mahasiswa, dosen dan praktisi untuk berinspirasi betapa ilmu kimia itu sangat menyentuh kehidupan umat manusia.

Kami yakin bahwa Jurnal Pendidikan Kimia yang diterbitkan oleh Program Pascasarjana Universitas Negeri Medan ini, akan semakin bermutu seiring dengan telah dibukanya Program Doktor (S3) Pendidikan Kimia di UNIMED sejak tahun akademik 2014-2015. Keberadaan 3 program studi linier dan berjenjang (Sarjana **S₁**, Magister **S₂** akreditasi A BAN PT, dan Doktor **S₃**) Pendidikan Kimia dalam satu atap di UNIMED, dan diperolehnya Akreditasi Institusi AIPT UNIMED peringkat B (nilai 353) dari BAN PT Kemdikbud, akan menambah kepercayaan masyarakat luas kepada institusi ini..

Seperti biasanya, semua isi artikel dalam Jurnal Pendidikan Kimia Pascasarjana Unimed, telah dapat diakses secara on-line melalui internet dari situs <http://www.digilib.unimed.ac.id/journal>. Kami menyadari bahwa isi maupun tayangan artikel pada edisi ini masih banyak kekurangan dan kelemahannya, kami mohon maaf, kiranya pada penerbitan mendatang semakin bagus. Kami tetap berharap kiranya para mahasiswa, guru, dosen, peneliti dan pemerhati yang terkait dengan ilmu kimia, biomedik , bioteknologi, kesehatan dan lingkungan agar mengirimkan naskahnya untuk dipublikasi dalam Jurnal ini.

Medan, April 2015

Penyunting

Volume 7 Nomor 1 Edisi April 2015

Daftar Isi

- 1 Penilaian TNF-Alfa pada hati mencit jantan setelah pemberian ekstrak etanol Manggis *Garcinia mangostana L* dengan metode imunohistokimia
Nora Maulina
- 2 Pengembangan bahan ajar inovatif untuk siswa SMA Pokok Bahasan Struktur Atom dan Tabel Periodik Unsur
Devi Anriani Siregar; Iis Siti Jahro; Ramlan Silaban
- 3 Ikatan translokator protein dari *Rhodobacter sph* hemin dalam deterjen DDM dan DPC
Nora Susanti
- 4 Pengembangan bahan ajar inovatif rumus kimia dan persamaan reaksi berbasis model pembelajaran PBL
Ramlan Silaban; Saronom Silaban; Freddy T.M., Panggabean; Elsa Ginting
- 5 Analisis kadar merkuri Hg dalam urin penambang emas tradisional (Studi Kasus di Desa Panton Luas Kecamatan Sawang Tapak Tuan, Aceh)
Harianto Bangun; Gusbakti Rusip; Zul Alfian
- 6 Pengaruh penggunaan media animasi computer dan motivasi belajar terhadap prestasi belajar mahasiswa dalam perkuliahan materi reaksi redoks dan elektrokimia
Freddy T.M. Panggabean; Saronom Silaban
- 7 Isolasi dan karakterisasi senyawa metabolit sekunder fraksi etil asetat dari kulit batang tumbuhan *Bauhinia purpurea L*
Ardiansyah; Herdini; Abdullah
- 8 Perbedaan hasil belajar kimia siswa menggunakan model pembelajaran TGT (*Teams Games Tournament*) dengan media Mind-mapping dan molymood pada Pokok Bahasan Hidrokarbon
Ratna Sari Dewi
- 9 Analisis resiko, karakteristik sosial, ekonomi, perilaku dan faktor lingkungan terhadap malaria (Studi Kasus di Kecamatan Arongan Lambalek Kabupaten Aceh Barat)
Susy Sriwahyuni Sukiswo; Rinidar; Sugito
- 10 Pengembangan buku ajar kimia SMA/MA Kelas XI Semester II dalam upaya meningkatkan hasil belajar dan menumbuhkembangkan karakter siswa
Mahmud; Ajat Sudrajat; Shofia Rija Napitupulu
- 11 Pengembangan penuntun praktikum tipe discovery dan tipe Project Base Learning pada pembelajaran elektrolit dan non elektrolit di SMA
Zakiah; Albinus Silalahi; Zainuddin Muchtar
- 12 Pengaruh model pembelajaran *Contextual teaching and Learning CTL* terhadap hasil belajar siswa pada Pokok Bahasan Sistem Koloid
Saronom Silaban; Neeta Sri Debora Simangunsong
- 13 Uji efek antiinflamasi ekstrak etanol daun Binahong (*Anrederacordifolia*) secara oral
Yuziani

IKATAN TRANSLOKATOR PROTEIN DARI *Rhodobacter sphaeroides* DENGAN HEMIN DALAM DDM DAN DPC

The Binding Behavior of *Rhodobacter sphaeroides* TSPO with Hemin in DDM and DPC Detergents

Nora Susanti

Dosen Jurusan Kimia FMPA Universitas Negeri Medan

Email : nora.susanti.s2@gmail.com

Abstrak

Ikatan antara translokator protein (TSPO) dari *Rhodobacter sphaeroides* (RsTSPO) dengan hemin dalam dua detergen yang berbeda yaitu n-dodecyl- β -D-maltoside (DDM) and n-dodecylphosphocholine (DPC) detergents telah diteliti. RsTSPO dipilih sebagai model untuk TSPO mamalia. RsTSPO di tanam dalam *E.coli* kemudian dieksraksi menggunakan buffer Tris pH 7,5 dan diikuti dengan pemurnian menggunakan Fast Protein Liquid Chromotography (FPLC). Hasil fluoresensi triptofan intrinsik menunjukkan bahwa RsTSPO memiliki afinitas terhadap hemin yang lebih tinggi dalam DDM ($K_d = 0.35 \mu\text{M}$) dibandingkan DPC ($K_d = 0.575 \pm 0.005 \mu\text{M}$).

Kata kunci : translokator protein, hemin, fluoresensi, ikatan

Abstract

We have studied the binding behavior of *Rhodobacter sphaeroides* translocator protein (RsTSPO) to one of its ligand Hemin in n-dodecyl- β -D-maltoside (DDM) and n-dodecylphosphocholine (DPC) detergents. RsTSPO was chosen as the model for mammalian TSPO. The RsTSPO was expressed in *E.coli* and followed by extraction in Tris buffer pH 7.5 and purification by utilizing FPLC. The result of intrinsic tryptophan fluorescence quenching showed that RsTSPO has higher affinity to bind to Hemin in DDM ($K_d = 0.35 \mu\text{M}$) compare to DPC ($K_d = 0.575 \pm 0.005 \mu\text{M}$).

Key word : translocator protein, hemin, fluorescence quenching, binding

Introduction

The 18 kDa Translocator protein (TSPO) is an integral membrane protein of five transmembrane helices. Membrane proteins are proteins associated with a membrane bilayer, and they typically comprise about 25% of all proteins encoded for by the genome (Voet, 2011).

TSPO in mammals was discovered by Braestrup *et al.* (Braestrup, 1977) as a secondary receptor for diazepam. TSPO was then known as a peripheral benzodiazepine receptor (PBR). In bacteria, TSPO was known as a tryptophan rich sensory protein (TspO) (Yeliseev, 1995). Due to the multiple functions of this protein, Papadopoulos *et al.*

(Papadopoulos, 2006) in 2006 suggested new nomenclature: Translocator Protein (TSPO).

Eukaryotic TSPO is primarily located in the outer mitochondrial membrane bilayer. TSPO is particularly enriched at the sites of outer/inner mitochondrial contact (Taliani, 2011). The carboxyl end of TSPO is located on the outside of the mitochondrion and the amino terminal is inside the mitochondrion (Bernassau, 1993). TSPO comprises up to 2% of the outer mitochondrial membrane protein (Miller, 2013). At low levels, TSPO is also expressed in plasma and nuclear membranes (Olson, 1988). Endogenous ligands for TSPO include protoporphyrin IX (PPIX), hemin and cholesterol. Protoporphyrin IX is the biosynthetic precursor of heme in mammals and chlorophyll or bacteriochlorophyll in plants or bacteria.

TSPO has been associated with several health conditions, namely post-ischemic heart reperfusion injury, cancers and neurodegenerative diseases. TSPO has become a potential drug and imaging target for the above conditions. We started our investigation of TSPO ligand binding and structural characterization with the *Rhodobacter sphaeroides* TSPO (RsTSPO) homolog for a number of reasons. The TSPO protein is highly evolutionary conserved (Papadopoulos,

2006 dan Fan, 2012) from bacteria to eukaryotes. Moreover, *Rhodobacter* is one of the closest ancestors of mitochondria (Bui, 1996).

A suitable environment is critical for the isolation and purification of every membrane protein. An integral membrane protein such as TSPO is not soluble in water, thus detergents and lipids are commonly used to solubilize membrane proteins in aqueous media. In this research, we investigated the behaviour of RsTSPO in two different detergents, the n-dodecyl- β -D-maltoside (DDM) and n-dodecylphosphocholine (DPC) using hemin as the ligand.

Materials and Methods

2.1 Materials

Chemicals were purchased from Fisher and Sigma Aldrich. Dodecyl maltoside (DDM) and n-dodecylphosphocoline (DPC) (Anagrade) were purchased from Anatrace. Hemin was purchased from Frontier Scientific. Lysozyme was purchased from Sigma Aldrich. Ampicillin and isopropyl β -D-1-thiogalactopyranoside (IPTG) were purchased from Gold Biotechnology. Chloramphenicol was purchased from Acron Organics. Complete EDTA free Protease inhibitor was purchased from Roche.

2.2 Methods

2.2.1 Protein expression and purification
E.coli BL21 (DE3) pLysS containing pET23(a) plasmid was grown overnight in 25 mL Luria broth. The overnight culture (5 mL) was then transferred in to 1 L of LB and grown to optical density at 600 nm (OD_{600}) of 0.7. The protein expression in the culture was then induced by the addition of isopropyl β -D-1-thiogalactopyranoside (IPTG) (0.4mM final concentration). Cells were then grown for another 6 hours while shaking. Cells were harvested by centrifugation (20 min $4,400 \times g$, 4 °C). Cell pellets were resuspended in Tris buffer (20 mM Tris and 200 mM NaCl, pH 7.50) and immediately frozen in liquid nitrogen. Cell pellets were stored at -80°C until cell lysis. Before cell lysis, complete EDTA free protease inhibitor cocktail and lysozyme were added to the resuspended pellet. Cells were lysed by sonication and by passing through French Press three times. The cell lysate was centrifuged (20 min, $8,000 \times g$, 4 °C) to separate cell debris. Membranes containing TSPO were collected by ultracentrifugation (2 h, $170,000 \times g$, and 4 °C).

2.1.2 Extraction and purification of TSPO

Membranes were resuspended in Tris buffer(20 mM Tris and 200 mM NaCl, pH 7.50) supplemented with 2 % of either DDM (equal to $196 \times CMC$) or DPC

(equal to $57 \times CMC$). Protein was extracted by gently shaking the solution for 6 h at 4 °C. Solubilized proteins and the membranes were separated by ultracentrifugation (2 h, $170,000 \times g$, 4°C). The protein was then concentrated with Amicon Ultra centrifugal filter (30 kDa molecular weight cutoff) and filtered with Sterile Milex Filter unit (0.22 μm). Extracted protein was purified with AKTA FPLC system using GE health His Trap 1 mL FF Ni-affinity column. Unbound proteins were washed off the column using Tris buffer(20 mM Tris and 200 mM NaCl, pH 7.50)containing either 0.1 % DPC (equal to $2.6 \times CMC$) or 0.025 % DDM (equal to $2.5 \times CMC$) and supplemented with 50 mM imidazole. TSPO was eluted with Tris buffer pH 7.50 containing either 0.1 % DPC or 0.025 % DDM and supplemented with 500 mM imidazole. After purification with the Ni-affinity column, the sample fractions containing the protein was passed through Superdex75 size exclusion column using Tris size exclusion buffer (20 mM Tris and 30 mM NaCl, pH 7.50) containing either 0.1 % DPC or 0.025 % DDM. Purified protein then was run on 12% SDS-PAGE (sodium dodecyl sulfate poly acrylamide gel electrophoresis) and stained using Coomassie Blue to visualize the protein.

2.1.3 TSPO ligand binding studies

TSPO binding studies were carried out with purified, detergent solubilized protein (0.1 % DPC or 0.025 % DDM). All experiments in the binding studies were done with excitation wavelength at 285 nm and emission at 290-800 nm at room temperature (20 °C). Excitation and emission slits were set to 2.5 nm and 5 nm, respectively, with both excitation and emission filter were set to auto. After each addition of the ligand, the solution was incubated for 5 minutes either in dark or in ambient light before measuring fluorescence. We used a Varian Cary Eclipse Fluorometer and a Varian Cary UV-vis spectrophotometer. Fluorescence intensity (290-800 nm) and absorbance spectra (200-800 nm) were measured for every titration point. Fluorescence intensity was plotted against ligand concentration. Percent quenching was calculated using integration for every titration point.

2.1.4 Data Analysis

Data analysis was performed in collaboration with Dr. Jan Kubelka of University of Wyoming using Matlab (Mathworks Inc., Natick, MA). The following binding model was considered:



where E is the enzyme, S is substrate, f is an effective number of enzyme monomers binding the substrate and fES stands for the complex. The binding constant is expressed as follows:

$$K_d = [fE][S]/[fES] \quad (2)$$

Using the total concentration of the enzyme $[E]_0$ and substrate $[S]_0$:

$$[E]_0 = f([fE] + [fES]) \quad (3)$$

$$[S]_0 = [S] + [fES]$$

in (2), the concentration of the complex is:

$$[fES] = ([E]_0 + f[S]_0 + K_d - \sqrt{([E]_0 + f[S]_0 + K_d)^2 - 4f[E]_0[S]_0})/2f$$

The observed fluorescence intensity Φ is then:

$$\Phi = [fE] \Phi_{fE} + [fES] \Phi_{fES}$$

Where Φ is the fluorescence of the enzyme Φ_{fE} (f-mer), and Φ_{fES} is the fluorescence of the complex. It is assumed that the fluorescence of the f-mer in the absence of bound ligand, is simply f times that of the monomer.

Result and Discussion

3.1 Expression and purification of TSPO

We have successfully expressed and purified RsTSPO in *E.coli* in both DDM and DPC detergents.

3.2 Binding study

RsTSPO solubilized in the nonionic detergent (DDM) exhibited greater TSPO

fluorescence quenching than RsTSPO solubilized in the zwitterionic DPC, which is line with the previously reported results (Li, 2013). TSPO in DDM also exhibit higher affinity to bind hemin ($K_d = 0.35 \mu\text{M}$) compare to DPC ($K_d = 0.575 \mu\text{M}$)




Figure 1. Fluorescence quenching of RsTSPO with hemin in DDM (A) and DPC (B)

A number of previous studies highlighted the critical role of lipids in bacterial and eukaryotic TSPO folding and function. For instance, it was shown in a study of mouse TSPO (Lacapere, 2001) that sodium dodecylsulfate (SDS) solubilized mouse TSPO was only able to bind steroid ligands, but not others. TSPO activity was recovered after reconstitution into lipid membranes. In binding studies of the bacterial RsTSPO, Li *et al.* (Li, 2013)

compared the decylmaltoside (DM), a nonionic detergent, with DPC, a zwitterionic detergent. Their result showed greater TSPO fluorescence quenching in DM than in DPC. In our hands, efficient fluorescent quenching was observed for RsTSPO solubilized in either DDM or DCP, which indicates that RsTSPO was purified in a functional state in both detergents. However, RsTSPO solubilized in the nonionic detergent (DDM) exhibited greater TSPO fluorescence quenching than RsTSPO solubilized in DPC, which is line with the previously reported results (Li, 2013). Clearly, the nature of the detergent influenced the binding of ligands to TSPO, and the dissociation constants (K_d) observed for hemin was consistently lower in DDM, which indicated higher ligand affinity. One possible explanation is that nonionic detergents such as DM and DDM can be less deactivating than Zwitterionic detergents such as DPC, as reported in the literature before (Seddon, 2004). However, the effect of the nature (i.e. nonionic vs. Zwitterionic) and concentration (i.e. 0.1 % DPC or 0.025 % DDM) of the detergent on the hydrophobic porphyrin ligands (e.g. ligand aggregation) cannot be excluded as a possible explanation for this observation, and more experiments are required to provide a workable theory.

Conclusion

RsTSPO in DDM has higher affinity to bind hemin compare to RsTSPO in DPC.

References

- Bernassau, J. M.; Reversat, J. L.; Ferrara, P.; Caput, D.; Lefur, G., A 3D Model of The Peripheral Benzodiazepine Receptor and Its Implication in Intra Mitochondrial Cholesterol Transport. *Journal of Molecular Graphics* **1993**,*11* (4), 236-244.
- Braestrup, C.; Squires, R. F., Specific benzodiazepine receptors in rat brain characterized by high-affinity (^3H)diazepam binding. *Proceedings of the National Academy of Sciences of the United States of America* **1977**,*74* (9), 3805-3809.
- Bui, E. T. N.; Bradley, P. J.; Johnson, P. J., A common evolutionary origin for mitochondria and hydrogenosomes. *Proceedings of the National Academy of Sciences of the United States of America* **1996**,*93* (18), 9651-9656.
- Fan, J.; Lindemann, P.; Feuilloley, M. G. J.; Papadopoulos, V., Structural and Functional Evolution of the Translocator Protein (18 kDa). *Current Molecular Medicine* **2012**,*12* (4), 369-386.
- Lacapere, J. J.; Delavoie, F.; Li, H.; Peranzi, G.; Maccario, J.; Papadopoulos, V.; Vidic, B., Structural and functional study of reconstituted peripheral benzodiazepine receptor. *Biochemical and Biophysical Research Communications* **2001**,*284* (2), 536-541.
- Li, F.; Xia, Y.; Meiler, J.; Ferguson-Miller, S., Characterization and Modeling of the Oligomeric State and Ligand Binding Behavior of Purified Translocator Protein 18 kDa from Rhodobacter sphaeroides. *Biochemistry* **2013**,*52* (34), 5884-5899.
- Miller, W. L., Steroid hormone synthesis in mitochondria. *Molecular and Cellular Endocrinology* **2013**,*379* (1-2), 62-73.
- Olson, J. M. M.; Ciliax, B. J.; Mancini, W. R.; Young, A. B., Presence of peripheral-type benzodiazepine binding sites on human erythrocyte membranes. *European Journal of Pharmacology* **1988**,*152* (1-2), 47-53.
- Papadopoulos, V.; Baraldi, M.; Guilarte, T. R.; Knudsen, T. B.; Lacapere, J. J.; Lindemann, P.; Norenberg, M. D.; Nutt, D.; Weizman, A.; Zhang, M. R.; Gavish, M., Translocator protein (18 kDa): new nomenclature for the peripheral-type benzodiazepine receptor based on its structure and molecular function. *Trends in Pharmacological Sciences* **2006**,*27* (8), 402-409.
- Seddon, A. M.; Curnow, P.; Booth, P. J., Membrane proteins, lipids and detergents: not just a soap opera. *Biochimica Et Biophysica Acta-Biomembranes* **2004**,*1666* (1-2), 105-117.
- Taliani, S.; Pugliesi, I.; Da Settim, F., Structural Requirements to Obtain Highly Potent and Selective 18 kDa Translocator Protein (TSPO) Ligands. *Current Topics in Medicinal Chemistry* **2011**,*11* (7), 860-886.
- Voet, D.; Voet, J. G., Biochemistry. 4th ed.; John Wiley & Sons, Inc: New Jersey, 2011.
- Yeliseev, A. A.; Kaplan, S., A Sensory Transducer Homologous to the Mammalian Peripheral-type Benzodiazepine Receptor Regulates Photosynthetic Membrane Complex-formation in *Rhodobacter sphaeroides*. *Journal of Biological Chemistry* **1995**,*270* (36), 21167-21175.

