



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Melanoxylon Cotylelobium
Assignment title: Jurnal Januari
Submission title: Cotylelobium melanoxylon
File name: teria_Isolated_from_Cotylelobium_m..
File size: 631.05K
Page count: 7
Word count: 3,447
Character count: 18,998
Submission date: 13-Jan-2021 10:06AM (UTC+0700)
Submission ID: 1486662640

Int J Pharm Bio Sci 2016 April; 7(2): (B) 666 - 672

Original Research Article Microbiology

 International Journal of Pharma and Bio Sciences ISSN 0975-6299

ANTIMICROBIAL ACTIVITIES OF ENDOPHYTIC BACTERIA ISOLATED FROM COTYLELOBIUM MELANOXYLON (HOOK.F.) PIERRE.

IDRAMSA¹, ENDANG SUTARININGSIH SOETARTO^{2*}, LAURENTIUS. H. NUGROHO³ AND RARASOETI PRATIWI²

¹Department of Biology, Faculty of Mathematics and Natural Sciences, the State University of Medan, Indonesia
²Faculty of Biology, Gadjah Mada University, Yogyakarta, Indonesia

ABSTRACT

The current study used the culturable endophytic bacterium *Nocardiopsis* sp. BER-17 isolated from the bark of *Cotylelobium melanoxylon* to synthesize a novel antimicrobial compounds. *Cotylelobium melanoxylon* is one of therapeutic plant species used in traditional medicine. It is containing compounds with antibacterial activity and bacterial endophytes that provide bioactive metabolites for medicinal purposes, and source of novel compounds. An attempt has been made to employ endophytic bacterium isolated from medicinal plants in the synthesis of antibacterial compounds. The objectives of this study was to investigate the antibacterial activities produced by *Nocardiopsis* sp. BER-17. Dichloromethane endophytic bacterial extract and culture broth were tested for antibacterial activity using well diffusion method against pathogenic bacteria of *Escherichia coli* ATCC 35218, *Staphylococcus aureus* ATCC 25923, *Bacillus subtilis* FNCC 001 and *Pseudomonas aeruginosa* FNCC 0054. The antibacterial activities were assessed based by the presence of clear zones as inhibition zones. Composition of secondary metabolites in the dichloromethane bacterial extract was spectrometrically analyzed using GC-MS. The endophytic bacterial extract readily produced with clear zones against both Gram-negative and Gram-positive bacterial pathogens. Based on GC-MS result, dichloromethane extract endophytic contain components antibacterial activity.

KEYWORDS: *Cotylelobium melanoxylon*, culturable endophytic bacterium, *Nocardiopsis* sp., antibacterial activities.

  **ENDANG SUTARININGSIH SOETARTO**
Faculty of Biology, Gadjah Mada University, Yogyakarta, Indonesia

*Corresponding author

This article can be downloaded from www.ijphs.net
B - 666



Cotylelobium melanoxylon

by Melanoxylon Cotylelobium

THE
Character Building
UNIVERSITY

Submission date: 13-Jan-2021 10:06AM (UTC+0700)

Submission ID: 1486662640

File name: teria_Isolated_from_Cotylelobium_melanoxylon_Hook._F._Pierre.pdf (631.05K)

Word count: 3447

Character count: 18998

ANTIMICROBIAL ACTIVITIES OF ENDOPHYTIC BACTERIA ISOLATED FROM
COTYLELOBIUM MELANOXYLON (HOOK.F.) PIERRE.IDRAMSA¹, ENDANG SUTARININGSIH SOETARTO*², LAURENTIUS. H. NUGROHO²
AND RARASTOETI PRATIWI²¹Department of Biology, Faculty of Mathematics and Natural Sciences,
the State University of Medan, Indonesia²Faculty of Biology, Gadjah Mada University, Yogyakarta, Indonesia

ABSTRACT

The current study used the culturable endophytic bacterium *Nocardiopsis* sp. BER-17 isolated from the barks of *Cotylelobium melanoxyton* to synthesize a novel antimicrobial compounds. *Cotylelobium melanoxyton* is one of therapeutic plant species used in traditional medicine. It is containing compounds with antibacterial activity and bacterial endophytes that provide bioactive metabolites for medicinal purposes, and sources of novel compounds. An attempt has been made to employ endophytic bacterium isolate from medicinal plants in the synthesis of antibacterial compounds. The objectives of this study was to investigate the antibacterial activities produced by *Nocardiopsis* sp. BER-17. Dichloromethane endophytic bacterial extract and culture broth were tested for antimicrobial activity using well diffusion method against pathogenic bacteria of *Escherichia coli* ATCC 35218, *Staphylococcus aureus* ATCC 25923, *Bacillus subtilis* FNCC 0061 and *Pseudomonas aeruginosa* FNCC 0063. The antibacterial activities were assessed based by the presence of clear zones as inhibition zones. Composition of secondary metabolites in the dichloromethane bacterial extract was spectrometrically analyzed using GC-MS. The endophytic bacterial extract activity produced with clear zones against both Gram negative and Gram positive bacterial pathogens. Based on GC-MS result, dichloromethane extract endophytic contain components antibacterial activity.

KEYWORDS: *Cotylelobium melanoxyton*, culturable endophytic bacterium, *Nocardiopsis* sp., antibacterial activities.



ENDANG SUTARININGSIH SOETARTO
Faculty of Biology, Gadjah Mada University, Yogyakarta, Indonesia

*Corresponding author

THE Character Building UNIVERSITY

INTRODUCTION

Tropical forests occupy approximately 7% of the world's total land areas and 50-70% of plant species are located in the area¹. More or less 300,000 plant species become the hosts of endophytic microbes². The threats of forest denudation in the tropical forest ecosystem can reduce the diversity of vegetation as habitat for endophytic microbes³. The exploration of endophytic microbes in the tropical forest ecosystem is necessary to do for multiple purposes such as in agricultural, health, and industrial sectors⁷. Many opportunities to recover endophytic microbes, which can be used to improve plant growth, as natural raw materials of drugs, and compound producers of secondary metabolites, are still open with the higher level of diversity in plant species in tropical forest ecosystem⁴. The diversity of plant species in tropical forests is largely not investigated its endophytic bacteria⁵. The potential diversity of endophytic microbes as well as their compound contents of secondary metabolites produced are still not identified well⁶. *Raru* (*Cotylelobium melanoxylon*) plant is a plant living in tropical regions and largely growing in Sumatera and Borneo islands as well as in Malaysia Peninsular and Thailand⁸. *C. melanoxylon* plant is a perennial plant with a tree-like form. Emergent tree up to 52 m tall and 72 cm dbh. Stem with resin. A medium-sized to tall tree reaching up to 50 m in height, frequently with a twisted bole, branchless up to 30 m and up to 160 cm in diameter. Stipules ca 3 mm long. Twigs more or less smooth³. Leaves alternate, ovate-lanceolate, 5-10 x 2-6 cm, simple, penni-veined, venation inconspicuous and fine, lower surface mostly very light to almost whitish. Flowers ca. 13 mm in diameter, white, placed in small panicles. Fruits ca. 10 mm long, yellow-green to reddish, nut with 2 wings up to 60 mm long, wind dispersed⁹. It is included in a red list of endangered plant according to the International Union for Conservation of Nature (IUCN) in 1998⁸. The organ parts of *C. melanoxylon* are largely utilized for multiple purposes. For example, its stem can be used as raw material for construction and its stem bark has been used by a group of communities as the mixing materials for *tuak* (traditional alcoholic beverage) in North Sumatera. Some local communities in North Tapanuli have used the stem barks and leaves of *C. melanoxylon* plant to treat some diseases such as diarrhea, malaria, and diabetes⁹. The stem barks of *C. melanoxylon* contains compounds consisting of ampelopsin F, isoampelopsin F, ϵ -viniferin, vaticanol A, E, G, and lyoniresinol that are useful as anti-diabetic drugs¹⁰. Moreover, the stem barks of *C. melanoxylon* contain a group of flavonoid, tannin, and saponin compounds¹¹. In order to isolate endophytic microbes as well as the potential compounds of secondary metabolite in plants, the following aspects must be considered: 1) the host plants are originated from the specific environment and have the high level of diversity; 2) they have ethno-botanical values closely related to the needs of local communities; and 3) they are endemic plants in a region². Based on them, *C. melanoxylon* was included into the criteria of host plants that can produce endophytic microbes with the potential compounds of secondary metabolite. *Nocardio* sp

BER-17 was a culturable endophytic bacterium, isolated using specific method from the barks of *C. melanoxylon*¹³. It demonstrated their activity against Gram positive and Gram negative bacterial pathogens. Therefor, *Nocardio* sp BER-17 was promising novel antimicrobial compound producer.

MATERIALS AND METHODS

a. Bacterial Cultures

The pure culture of endophytic bacteria *Nocardio* sp of strain BER-17 was obtained from the isolation from the barks of *C. melanoxylon*. Endophytic bacteria *Nocardio* sp. BER-17 is identified based on 16S rRNA gene sequences¹². Identification of the plants *C. melanoxylon* conducted in Herbarium LIPI Cibinong, based on morphological characteristic compared with herbarium specimen data. The bacteria test used were *Escherichia coli* ATCC 35218, *Staphylococcus aureus* ATCC 25923, *Bacillus subtilis* FNCC 0061 and *P. aeruginosa* FNCC 0063 that were obtained from the Laboratory of Microbiology, Faculty of Medicine and the Laboratory of Food and Nutrient, Biotechnology, the Inter-University Center, Gadjah Mada University, Yogyakarta. All bacterial cultures were previously purified by the method of single cell culture before being used in the test¹³. Colony morphology and endophytic bacterial cells were characterized by Gram staining method.

a. Bacterial Cultivation for Antibacterial Production

The endophytic bacteria *Nocardio* sp. of strain BER-17 were grown in 250 ml Erlenmeyer flask containing TSB 100 ml medium (three-fold repetitions) on pH 7-8. They were incubated at a temperature of 37°C in shaker with speed of 180 rpm for 7 and 14 days. After the bacteria grown, the culture media was centrifugated with speed of 13,000 x g for 5 minutes and supernatants were filtered using whatman filter paper No. 5, so that filtrate media solution was gained. The supernatant were extracted with the same volume of the extracting solution for 3 times consecutively using ethyl acetate, dichloromethane and n-butanol. It was concentrated using vacuum rotary evaporator. The resultant extracts were stored in desiccators for use in further test¹².

b. Detection of Antibacterial Activity

The compounds of endophytic bacteria *Nocardio* sp. of strain BER-17 that were extracted by ethyl acetate, dichloromethane, and n-butanol were tested by Kirby-Baur method¹⁴. The bacteria *B. subtilis*, *E. coli*, *S. aureus* and *P. aeruginosa* were grown in nutrient agar medium for 24 hours. Furthermore, using ose, pathogenic bacteria were inoculated in nutrient broth media and incubated for 5-8 hours or a night at a temperature of 37°C. Furthermore, 1 ose was taken by using sterile cotton swabs that was previously inoculated by a scratch method on surface of petri cup containing nutrient agar medium and being left for 5 minutes²¹. Paper disc drops of as much as 10 μ L extraction endophytic bacteria *Nocardio* sp. BER-17 and dried. Paper disk was immersed into the extracts of endophytic bacteria *Nocardio* sp of strain BER-17 and placed in the sterile petri cup for wind drying.

Furthermore, the paper disk was placed by pinset on the surface of nutrient agar medium previously inoculated in the bacteria and incubated for 2-3 days at a temperature of 37°C. Inhibition zone diameter was measured (mm) using a millimeter ruler¹⁴.

b. GC-MS Analysis

Compounds resulted from extraction of endophytic bacteria *Nocardiopsis* sp of strain BER-17 were analyzed by using gaseous chromatography and mass spectroscopy (GC-MS). The analysis was carried out by using GC-MS QP2010S SHIMADZU, Japan. The operating procedures and conditions applied the type of capillary column AGILENT HP 5MS (30 M× 0.25 mm, I.D.). Helium was used as carrying gas with a constant rate of 0.40 mL/min, being injected for 1 µL (split ratio of 33:1) at an injector temperature of 310°C. The column was programmed at a temperature of 120°C (5 minutes) to 300°C (69 minutes) with the increase of temperature regulated by 5°C/minute. The conditions of GC-MS showed the ion source temperature of 250°C, the interface temperature of 305°C, the pressure of 13.7 kPa. The interpretation of mass spectrum GC-MS was identified by using the database from the National Institute Standard and Technology (NIST62.LIB and

WILEY229.LIB). The components of compounds assayed were recognized, i.e. name, weight, and structure of molecules²⁰.

RESULTS

The microorganisms of *B. subtilis* (FNCC 0061), *E. Coli* (ATCC 3521), *P. aeruginosa* (FNCC 0063) and *S. aureus* (ATCC 25923) were tested by using the dichloromethane, ethyl acetate and N-butanol extracts from endophytic bacteria of strain BER-17 (Table 1). The dichloromethane extracts from endophytic bacteria of strain BER-17 were able to inhibit bacterial growth of *B. subtilis* (25 mm), *E. coli* (17 mm) and *S. aureus* (18 mm). The ethyl acetate extracts and N-butanol from endophytic bacteria of strain BER-17 were able to inhibit bacterial growth of *P. aeruginosa* (18 mm) and (17 mm). Based on the results of the analysis, the dichloromethane extracts from endophytic bacteria of strain BER-17 were selected because the extracts showed wide inhibitory zone against many pathogenic bacteria.



Figure 1

Antibacteri activities test of endophytic *Nocardiopsis* sp. BER-17 cultivation for 7 and 14 days with extraction by diklorometan (DC), etil asetat (EA) and n-butanol (n-But). (A) *Escherichia coli* ATCC 35218 (DC 14 = 17 mm); (B) *Staphylococcus aureus* ATCC 25923 (DC 14 = 18 mm); (C) *Bacillus subtilis* FNCC 0061 (DC 14 = 25 mm and (D) *Pseudomonas aeruginosa* FNCC 0063 (EA 14 = 18 mm and n-But = 17 mm).

Table 1
Antibacterial activity of dichloromethane extracts from endophytic bacteria *Nocardiopsis* sp. BER-17

Bacteria test	Dichloromethane	Ethyl acetate	N-butanol
<i>B. subtilis</i> FNCC 0061	25	-	-
<i>E. coli</i> ATCC 35218	17	-	-
<i>P. aeruginosa</i> FNCC 0063	-	18	17
<i>S. aureus</i> ATCC 25923	18	-	-

The results of analysis using GC-MS showed eight peaks (Figure 2) with the retention times of 32.259, 31.533, 24.543, 24.025, 23.664, 23.268, 21.850, and 21.199. Based on the results of GCMS, eight components of compounds in the dichloromethane extracts from endophytic bacteria of strain BER-17 were found. The mass spectra of each of the compounds are shown from Figure 3 to 10.




Figure 2
GC-MS chromatogram of the dichloromethane endophytic bacterial crude extract




Figure 3
2-decene, 3-methyl




Figure 4
1,2-cyclopentanedione, 3,3,5,5-tetramethyl



Figure 5
1,4-diaza-2,5-dioxo-3-isobutyl bicyclo[4.3.0]nonane




Figure 6
Aziridine, 2-heptyl-3-methyl





Figure 7
1,3,2-Dioxaborinane, 2,4,6-triethyl-5-methyl-



Figure 8
Hexadecanoic acid ethyl ester



1
Figure 9
3-benzyl-1,4-diaza-2,5-dioxobicyclo[4.3.0]nonane



Figure 10
Dihydroergotamine

The components of compounds as the results of analysis using GC-MS: 2-decene, 3-methyl (13.80%), 1,2-cyclopentanedione, 3,3,5,5,-tetramethyl (9.76%), 1,4-diaza-2,5-dioxo-3-isobutyl bicyclo[4.3.0]nonane (9.09%), aziridine, 2-heptyl-3-(11.08%), 1,3,2-dioxaborinane, 2,4,6-triethyl-5-methyl-(20.37%), hexadecanoic acid ethyl ester (28.31%), 3-benzyl-1,4-diaza-2,5-dioxobicyclo[4.3.0]nonane (4.34%) and dihydroergotamine (3.25%) (Table 2).

Tabel 2
Compound identified in dichloromethane endophytic bacterial
***Nocardioopsis* sp BER-17 crude extract by GC-MS**

Chromatogram peak	Compound name	Molecular formula	Molecular weight	Peak area (%)	Retention time (min)	Sim. Index (%)
1	2-Decene, 3-methyl-(Z)-	C ₁₁ H ₂₂	154	13.80	21.200	79
2	1,2-Cyclopentanedione, 3,3,5,5,-tetramethyl	C ₉ H ₁₄ O ₂	154	9.76	21.850	77
3	1,4-diaza-2,5-dioxo-3-isobutyl bicyclo[4.3.0]nonane	C ₁₁ H ₁₈ N ₂ O ₂	210	9.09	23.267	83
4	Aziridine, 2-heptyl-3-methyl	C ₁₀ H ₂₁ N	155	11.08	23.667	77
5	1,3,2-Dioxaborinane, 2,4,6-triethyl-5-methyl-	C ₁₀ H ₂₁ BO ₂	184	20.37	24.025	80
6	Hexadecanoic acid, ethyl ester	C ₁₈ H ₃₆ O ₂	284	28.31	24.542	92
7	3-benzyl-1,4-diaza-2,5-dioxobicyclo[4.3.0]nonane	C ₁₄ H ₁₆ N ₂ O ₂	244	4.34	31.533	79
8	Dihydroergotamine	C ₃₃ H ₃₇ N ₅ O ₅	244	3.25	32.259	72

DISCUSSIONS

Nocardioopsis sp. of strain BER-17 is an endophytic microbe isolated from the bark of *raru* (*Cotylelobium melanoxylon*) and identified based on the sequence of 16S rRNA gene¹³. From the results of the study conducted, *Nocardioopsis* sp as endophytic microbe has been found in other vegetations such as yam²², mangrove²³, *Hygrophila spinosa*²⁴, *Ocimum sanctum*²⁵, and *Hibicus rosasinensis*²⁶. Endophytic bacteria *Nocardioopsis* contained in some plant tissues as endophytes or surface microflora serves to produce α -amylase and antifungal enzymes against lethal plant pathogens¹⁵. The dichloromethane extracts of endophytic bacteria *Nocardioopsis* sp. of strain BER-17 contain compounds with the activity against Gram-negative (*E. coli* ATCC 35218 and *P. aeruginosa* FNCC 0063) bacteria as well as Gram-positive (*B. subtilis* FNCC 0061 and *S. aureus* ATCC 25923) bacteria. The classes of *Actinobacteria* with genus *Streptomyces*, *Micromonospora* and *Nocardioopsis* had antimicrobial activity against pathogenic bacteria¹⁶. Member of the genus *Nocardioopsis* is clustered into the class *Actinobacteria* consisting of several species and able to grow in extreme and contaminated environment, so that during an adaptation process they produce chemical compounds that are useful for drugs and industry²⁷. The components of compounds resulted from the dichloromethane extracts from endophytic bacteria of *C. melanoxylon* plant had various biological activities (Table 2). The secondary metabolite compounds resulted from several species of *Nocardioopsis* sp. include 3-trehalosamine, phenazine, 1,6-dihydroxyphenazine, 1,6-dihydroxyphenazine-5,10-dioxide, indole alkaloid, pendolmycin, apoptolidin, kahakamides A dan B, neosidomycin, endo-1,3- β -glucanase, endo- β -1,4-D-glucanase, 3-

hydroxybutyrate, 3-hydroxyvalerate²⁸. Compounds of 1,4-diaza-2,5-dioxo-3-isobutyl bicyclo[4.3.0]nonane and 3-benzyl-1,4-diaza-2,5-dioxobicyclo[4.3.0]nonane had antioxidant activity extracted by chloroform from endophytic bacteria of BSI isolate¹⁷. In addition, reported compounds of 3-benzyl-1,4-diaza-2,5-dioxobicyclo[4.3.0]nonane extracted from isolates as bacteria sample from sea. *Burkholderia cepacia* has antibacterial and anticancer compounds¹⁸. The compound of hexadecanoic acid ethyl ester has antibacterial activities that inhibit bacterial growth of *E. coli*, *S. typhi* and *S. aureus*¹⁹. The hexadecanoic acid ethyl ester also has antioxidant, *hypcholesterolemic*, nematocide, pesticide, lubricant, antiandrogenic activities²⁰.

CONCLUSION

The study is an effort to seek the sources of raw materials for new natural medicines from endophytic bacteria. It investigated antimicrobial activities of the extract of *Nocardioopsis* sp of strain BER-17 against pathogenic bacteria *B. subtilis* (FNCC 061), *E. Coli* (ATCC 3521), *P. aruginosa* (FNCC 063), and *S. aureus* (ATCC 15923). The results of the analysis using GC-MS showed the components of active compounds with antibacterial activities, being then assayed in vivo as a model of antibacterial drugs.)

ACKNOWLEDGEMENT

The study was a part of my Ph.D research. The author thanks to the Directorate General of Higher Education providing scholarship for attending the doctoral program in Gadjah Mada University.

REFERENCES

1. Strobel G. Harnessing endophytes for industrial microbiology. *Curr Opin Microbiol.* 2006 Jun 27;9(3):240-244.
2. Strobel G, Daisy B. Bioprospecting for microbial endophytes and their natural products. *Microbiol Mol Biol Rev.* 2003 Dec;67(4):491-502.
3. Sekar S, Kandavel D. Interaction of plant growth promoting rhizobacteria (PGPR) and endophytes with medicinal plants—new avenues for phytochemicals. *J Phytol.* 2010 Nov;2(7):91-100.
4. Strobel GA. Endophytes as sources of bioactive products. *Microbes Infect.* 2003 May;5(6):535-544.

5. Izumi H. Diversity of endophytic bacteria in forest trees. In: Pirttila AM, Frank AC, editors. Endophytes of forest trees. London: Springer Dordrecht Heidelberg; 2011. p. 95-105.
6. Khan R, Shahzad S, Choudhary MI, Khan SA, Ahmad A. Communities of endophytic fungi in medicinal plant *Withania somnifera*. Pak J Bot. 2010 Jan;42(2):1281-1287.
7. Verma VC, Gond SK, Kumar A, Mishra A, Kharwar RN, Gange AC. Endophytic actinomycetes from *Azadirachta indica* A. Juss.: isolation, diversity, and anti-microbial activity. Microb Ecol. 2009 May;57(4):749-59.
8. The IUCN Red List of Threatened Species. *Cotylelobium melanoxylon* [Internet]. United Kingdom: IUCN Global Species Programme Red List Unit; 1998 [updated 1998 January 1; cited 2012 Dec 22]. Available from: <http://www.iucnredlist.org/details/33070/0>.
9. Soerianegara I, Lemmens RHJ. Plant resources of South-East Asia No.5 (1). Timber trees: Minor commercial timber. Bogor: Prosea Foundation; 1994.
10. Matsuda H, Aso Y, Nakamura S, Hamao M, Sugimoto S, Hongo M, et al. Antidiabetogenic constituents from the Thai traditional medicine *Cotylelobium melanoxylon*. Chem Pharm Bull. 2009 May;57(5):487-94.
11. Pasaribu G. Inhibition activity of alpha glucosidase from several stem bark of raru. Jurnal Penelitian Hasil Hutan. 2011 Mar;29(1):10-19.
12. Melo PFM de, Fiore MF, Moraes LAB de, Silva-Stenico ME, Scramin S, Teixeira M de A, et al. Antifungal compound produced by the cassava endophyte *Bacillus pumilus* MAIIM4A. Sci Agric. 2009 Sep;66(5):583-592.
13. Idramsa, Soetarto ES, Nugroho LH, Pratiwi R, Prasetya E. Endophytic bacteria inducing antibacterial synthesis of the bark of raru (*Cotylelobium melanoxylon*). Euro J Exp Bio. 2015 Oct;5(9):20-26.
14. Cappuccino JG, Sherman N. Microbiology, A Laboratory Manual. 9th ed. San Francisco: Pearson Benjamin Cummings; 2011.
15. Bennur T, Kumar AR, Zinjarde S, Javdekar V. *Nocardiosis* species: Incidence, ecological roles and adaptations. Microbiol Res. 2015 May;174:33-47.
16. Machavariani NG, Ivankova TD, Sineva ON, Terekhova LP. Isolation of endophytic actinomycetes from medicinal plants of the Moscow Region Russia. World Appl Sci J. 2014 Jan;30(11):1599-1604.
17. Bintang M, Purwanto UMS, Kusumawati DE, Yang JJ. Study of endophytic bacteria as novel source of antioxidant agent based on GC-MS analysis. Int Journal Chem Envi Biol Sci. 2015 Aug;3(5):368-369.
18. Gohar YM, Manal MA, El-Naggar MMA, Soliman MK, Barakat KM. Characterization of marine Burkholderia cepacia antibacterial agents. J Nat Prod. 2010 Oct;3:86-94.
19. Igwe OU, Okwu DE. GC-MS evaluation of bioactive compounds and antibacterial activity of the oil fraction from the seeds of *Brachystegia curycoma* (HARMS). Asian J Plant Sci Res. 2013;3(2):47-54.
20. Zayed MZ, Ahmad FB, Ho WS, Pang SH. GC-MS analysis of phytochemical constituents in leaf extracts of *Neolamarckia cadamba* (Rubiaceae) from Malaysia. Int J Pharm Pharma Sci. 2014 Aug;6(9):123-127.
21. Devi NN, Wahab F. Antimicrobial properties of endophytic fungi isolated from medicinal plant *Camellia sinensis*. Int J Pharm Bio Sci. 2012 Jul;3(3):420-427.
22. Stamford TLM, Stamford NP, Coelho LCBB, Araujo JM. Production and characterization of a thermostable α -amylase from *Nocardopsis* sp. endophyte of yam bean. Bioresour Technol. 2001 Jan;76(2):137-41.
23. Lin C, Lu C, Shen Y. Three new 2-pyrone derivatives from mangrove endophytic actinomycete strain *Nocardopsis* sp. A00203. Rec Nat Prod. 2010 May;4(4):176-179.
24. Pal A, Paul AK. Bacterial endophytes of the medicinal herb *Hygrophila spinosa* T. Aders and their antimicrobial activity. Br J Pharm Res. 2013 Mar;3(4):795-806.
25. Singh MJ, Padmavathy S. *Nocardopsis* sp. 5 endophytic to tulsj leaves-isolation and antimicrobial activity. Br Microbiol Res J. 2014 Aug;5(3):194-202.
26. Singh MJ, Sedhuraman P. Biosurfactant, polythene, plastic, and diesel biodegradation activity of endophytic *Nocardopsis* sp. minalini9 isolated from *Hibiscus rosasinensis* leaves. Bioresource and Bioprocess. 2015 Jan;2(2):1-7.
27. Ballav S, Dastager SG, Kerkar S. Biotechnological significance of Actinobacterial research in India. Recent Res Sci Technol. 2012 Feb;4(4):31-39.
28. Hozzein WN, Trujillo ME. Genus I. *Nocardopsis*. In: Goodfellow et al. Bergey's manual of systematic bacteriology. 2nd ed. London: Springer Dordrecht Heidelberg; 2012. p 1891-1906.

Cotylelobium melanoxyton

ORIGINALITY REPORT

9%

SIMILARITY INDEX

9%

INTERNET SOURCES

6%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

repositorio.ufsm.br

Internet Source

4%

2

mafiadoc.com

Internet Source


3%

3

www.asianplant.net

Internet Source

2%



Exclude quotes On

Exclude matches < 2%

Exclude bibliography On

THE *Character Building* UNIVERSITY