

# The effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise induced rats

*by* Novita Sari Harahap

---

**Submission date:** 08-Mar-2021 12:00AM (UTC-0800)

**Submission ID:** 1527243712

**File name:** F\_1000\_RESEARCH\_UPDATE\_MARET\_2020.pdf (838.31K)

**Word count:** 6566

**Character count:** 34982



## RESEARCH ARTICLE

# The effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise induced rats [version 1; peer review: 1 approved, 1 approved with reservations]

Novita Sari Harahap<sup>1</sup>, Aznan Lelo<sup>2</sup>, Ambrosius Purba<sup>3</sup>, Awaluddin Sibuea<sup>4</sup>, Rina Amelia<sup>5</sup>, Zulaini Zulaini<sup>1</sup>

<sup>1</sup>Department of Sports Sciences, Faculty of Sports Sciences, Universitas Negeri Medan, Medan, North Sumatra, 20221, Indonesia

<sup>2</sup>Department of Pharmacology, Faculty of Medicine, University of Sumatra Utara, Medan, North Sumatra, 20155, Indonesia

<sup>3</sup>Department of Physiology, Faculty of Medicine, Padjajaran University, Bandung, West Java, 45363, Indonesia

<sup>4</sup>Surgery Division, Dr. T. Mansyur Tanjung Balai Hospital, Tanjungbalai, North Sumatra, 21312, Indonesia

<sup>5</sup>Department of Public Health, Faculty of Medicine, University of Sumatra Utara, Medan, North Sumatra, 20155, Indonesia

v1 First published: 30 Jan 2019, 8:130 (<https://doi.org/10.12688/f1000research.17533.1>)

Latest published: 30 Jan 2019, 8:130 (<https://doi.org/10.12688/f1000research.17533.1>)

## Open Peer Review

### Reviewer Status

?



### Abstract

**Background:** Oxidative stress from exercise can contribute to damaging cells, increasing heat shock protein 70 (HSP70) and suppressing the immune system in the body. This research aimed to determine the antioxidant potential of red-fleshed pitaya extract on HSP70 and cortisol expression in rats which were subjected to strenuous exercise.

**Methods:** The subjects of this research were 32 Sprague Dawley male rats, aged 3 months, with an average weight of 200 g. Red-fleshed pitaya extract was obtained from methanol extraction process; a maceration technique was performed and the extract was concentrated using an air-drying method. Rats were randomly divided into four groups. Group 1 were subjected to strenuous exercise and treated with distilled water only; while Groups 2, 3 and 4 were subjected to strenuous exercise and treated with 100 mg/kg body weight, 200 mg/kg body weight and 300 mg/kg body weight of red-fleshed pitaya extract, respectively. Strenuous exercises in rats was performed by intense swimming of 20 min/day, 3 days a week for 3 weeks. HSP70 expression and cortisol were measured with Enzyme-Linked Immune Sorbent Assay (ELISA) method.

**Results:** There was a significant reduction of HSP70 ( $p=0.000$ ) and cortisol expression ( $p=0.000$ ) between the groups. Also, there was a significant difference in the average decreasing of HSP70 expression between group 4 and either groups 1 or 2 ( $p=0.000$ ). However, a significant difference between groups 4 and 3 was not observed ( $p=0.813$ ). Lastly, a significant difference was found in the average decrease of cortisol expression between groups 4 and 1 ( $p=0.000$ ), 2 ( $p=0.000$ ), and 3 ( $p=0.000$ ) respectively.

**Conclusion:** Red-fleshed pitaya is potential to be utilized as antioxidant to decrease the HSP70 and cortisol expression.

### Invited Reviewers

1 2

version 1

30 Jan 2019



1 Yusni Yusni Syiah Kuala University, Banda Aceh, Indonesia

2 Mohd Adzim Khalili Rohin Sultan Zainal Abidin University (UniSZA), Kuala Nerus, Malaysia

Any reports and responses or comments on the article can be found at the end of the article.

**Keywords**

Red-fleshed pitaya, HSP70, cortisol, strenuous exercise

**Corresponding author:** Novita Sari Harahap ([novitahrp74@gmail.com](mailto:novitahrp74@gmail.com))

**Author roles:** Harahap NS: Conceptualization, Investigation, Methodology, Resources, Writing – Original Draft Preparation; Lelo A: Investigation, Methodology; Purba A: Resources, Supervision; Sibuea A: Resources; Amelia R: Resources; Zulaini Z: Formal Analysis

**Competing interests:** No competing interests were disclosed.

**Grant information:** The authors would like to acknowledge Ministry of Research, Technology and Higher Education of Republic of Indonesia for funding this research via Research Grant of Universitas Negeri Medan with contract number of 027/UN33.8/LL/2018.

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2019 Harahap NS *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Harahap NS, Lelo A, Purba A *et al.* *The effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise induced rats [version 1; peer review: 1 approved, 1 approved with reservations]* F1000Research 2019, 8:130 (<https://doi.org/10.12688/f1000research.17533.1>)

**First published:** 30 Jan 2019, 8:130 (<https://doi.org/10.12688/f1000research.17533.1>)

## Introduction

Physical activity is an activity which has various influences and significant effects on the body. The effect of regular physical activity is a positive influence on biological functions, and will improve health and the antioxidant defense system in order to protect body from the negative effects of oxidative damage<sup>1</sup>. Strenuous exercise tends to trigger free radical compound production. Moreover, this impairs the balance of free radicals and antioxidants as a result of oxidative stress<sup>2-4</sup>. Research has discovered that oxidative stress from strenuous exercise reduces performance as it damages cells<sup>5</sup>, causing pain and <sup>37</sup> muscles fatigues<sup>6</sup>, lowering antioxidant levels<sup>7,8</sup>, increasing the expression of heat shock protein 70 (HSP70)<sup>9,10</sup> and suppressing the immune system<sup>11</sup>.

During an intense workout, self-defense and self-adaptation depend on the body condition which can be observed from HSP70 protein expression. Increases of HSP70 in muscles indicates a response to protect muscles cells from oxidative stress. HSP70 expression is an adaptation mechanism and a sign of damaging cells caused by oxidative stress<sup>12</sup>. Previous research reported that workouts increase HSP70 expression<sup>10</sup>. Strenuous exercise is a physical stressor in the body and as a result, adrenocorticotrophic hormone (ACTH) is secreted by hypothalamus hypo-physis anterior and triggers the adrenal cortex to produce cortisol<sup>13</sup>. The escalation of cortisol is influenced by the intensity and duration of training that leads to a suppression of the immune system, resulting in a decline of antibody. Cortisol can be a sign that the body encounters a decline in the immune system due to heavy training<sup>14</sup>. Antioxidants can detoxify the lipid peroxide produced during exercise, which can eliminate radicals and reduce the inflammatory response to exercise. Therefore, it can prevent a muscle damage from exercise<sup>15</sup>.

**13**

The body needs exogenous antioxidants to neutralize and prevent chain reactions from free radicals formed from heavy physical trainings<sup>16</sup>. Sources of exogenous antioxidants are Vitamin E, C and also beta carotene. External antioxidants from food or supplements can help the body to fight an excess of free radicals. In a previous study, proanthocyanidin from grape seed was given to rats for 2 weeks. As a result, it lowered malondialdehyde level and increased the superoxide dismutase and glutathione peroxidase was activated significantly. Furthermore, it reduced fatigue after physical activity<sup>17</sup>.

Red-fleshed pitaya (*Hylocereus polyrhizus*) is a unique fruit with a lot of benefits. The fruit is recently popular among Indonesians and appears to be a natural antioxidant. Several *in vitro* studies have revealed that red-fleshed pitaya extract has the potential to be an antioxidant<sup>18</sup>. This research aimed to investigate the antioxidant potential in red-fleshed pitaya extract on HSP70 and cortisol expression in rats that were subjected to strenuous exercise.

## Methods

### Experimental animals

**11**

The subjects of this research were 32 Sprague Dawley male rats, aged 3 months, with an average weight of 200 g, were obtained from the Animal Holding Unit of the Pharmacy laboratory, University of Sumatera Utara, Indonesia.

All rats were sustained and maintained in groups (four mice per cage) in experimental animal cages of the Pharmacy laboratory, University of Sumatera Utara, Indonesia. The cage is made of plastic (30 x 20 x 10 cm) and covered with fine wire mesh. The base of the cage is covered with rice husk as thick as 0.5 - 1 cm and replaced every day during the research. The room light was controlled to be exactly at 12 hours light and 12 hours dark cycle, while the temperature 25–27°C and humidity of the room were adjusted to a normal range and fed with standard rat pellets 551, and drink (tap water) was given *ad libitum*.

### Study design

The research applied a laboratory experiment method with random group posttest-only design. The male rats were obtained from the Pharmacy Laboratory, University of North Sumatra. The experimental animals were simple random sampling divided into four groups: Group 1 was subjected to strenuous exercise and treated with distilled water only; Group 2 was subjected to strenuous exercise and treated with dosage 100 mg/kg body weight of red-fleshed pitaya extract; Group 3 <sup>34</sup> subjected to strenuous exercise and treated with dosage 200 mg/kg body weight of red-fleshed pitaya extract; Group 4 was subjected to strenuous exercise and treated with dosage 300 mg/kg body weight of red-fleshed pitaya extract.

### Red-fleshed pitaya extract

Red-fleshed pitaya fruit, obtained from farmers, in Indonesia, was peeled, washed, cut into small pieces and then dried in a drying cabinet. After that, the fruit was blended using a blender. The fruit extract was isolated through maceration method by using ethanol 96% which has been distilled as much as 10 times the weight of red-fleshed pitaya powder. Red-fleshed pitaya fruit powder in a container had 96% ethanol added to it (ratio 1:7, fruit powder: ethanol), and then was soaked for 3 days then filtered and sealed. The macerates were collected in a container and then processed with rotary evaporator at a temperature of 45°C until the extract was thickened. After that, the same process were repeated the remaining ethanol 96% for 3 days. The less thickened extract was then evaporated in a water bath until a thick extract was obtained.

100 mg red-fleshed pitaya extract was weighed. Then, it was gently ground using a pestle and mortar. After that, carboxy methyl cellulose (CMC) Na 0.5% solution was slowly added and ground until a homogeneous phase was achieved. Finally, the suspension was added to a 10 mL measuring flask until it reached the mark <sup>19</sup>. The allocation of red-fleshed pitaya extract, dosage of 100 mg/kg body weight, for instance: weight <sup>19</sup> of 200 g, volume taken: 2 ml extract suspension. Dosage of 200 mg/kg body weight, for instance: weight <sup>19</sup> 200 g, volume taken: 4 ml extract suspension. Dosage of 300 mg/kg body weight, for instance: weight of 200 g, volume taken: 6 ml extract suspension.

### Experimental procedures

Strenuous exercise given to all rats was a morning swim between 8–9 am for 20 minutes/day, 3 days a week over 21 days. The equipment used in this research was a 10-cm length and 25-cm diameter bath as a pool. Group 1, the rats received distilled water only; Group 2, the rats received 2 ml red-fleshed pitaya

extract suspension; Group 3, the rats received 4 ml red-fleshed pitaya extract suspension; Group 4, the rats received 6 ml red-fleshed pitaya extract suspension. Administration of red-fleshed pitaya extract suspension and water was performed orally once daily for 21 days.

Testing for HSP70 and cortisol was conducted two days after the rats had completed a strenuous exercise, 3 days a week over 21 days. During the test, the rats were given a maximum training session by swimming as hard as they could until the rats drowned or showed fatigue symptoms such as the entire body almost dipped into water and limb movements slowed down. After that the rats were sacrificed by placing them in a jar containing cotton which was moistened with 10 ml of chloroform. After that, 2–3 ml blood was taken from the heart.

22

Blood samples were collected in micro tubes and centrifuged at 3000 rpm for 15 minutes. The serum was separated and stored at a temperature of 20°C until the analysis process would be carried out. Cortisol was measured with enzyme-linked immune sorbent assay (ELISA); Mouse Cortisol Elisa kit (catalog: E1483Mo, Brand Bioassay TL) and UV spectrophotometry at a wavelength of 450 nm. The HSP70 expression was recorded with ELISA; Mouse HSP70 Elisa Kit (catalog: E0302Mo, Brand Bioassay TL); the absorbance was indicated at 405 nm.

#### Statistical methods

Data was analyzed using SPSS 22 for Windows and displayed in tables and diagrams. Normality test was conducted through Shapiro-Wilk test ( $P > 0.05$ ) in order to determine the average of normal distribution of sample data which is presented as mean  $\pm$  SD. The result of the normality test was used for next analysis; parametric analysis was used for normal distribution, otherwise non-parametric analysis was used. The ANOVA statistical analysis was performed to indicate the effects of treatments for each group. If the significant result is obtained, then the procedure is followed by Least Significance Difference or Bonferroni tests.

#### Ethical approval

The research was performed on the animal subjects were in accordance with the ethical standards by the Animal Research Ethics Committees/AREC, Faculty of Mathematics and Natural Sciences University of Sumatera Utara, Indonesia (approval number 0011/KEPH-FMIPA/2018).

All efforts were made to reduce any suffering of the rats was during the experiments by following careful procedures and also by anaesthetizing the animal prior to sacrifice to prevent experiencing any pain.

#### Results

A normality test indicated that the data are normally distributed (Table 1). HSP70 expression was decreased across all groups (69.57 vs 46.04 vs 31.47 vs 27.65 pg/mL). Group 4 had the lowest expression compared with the other groups. This research reveals a significant decrease in HSP70 expression ( $p=0.000$ ) between the groups (Table 2).

Cortisol expression was also decreased across all groups (119.02 vs 86.11 vs 62.94 vs 40.86 pg/mL). Group 4 had the lowest expression compared with the other groups. An ANOVA test revealed a significant decrease in cortisol expression ( $p=0.000$ ) between the groups (Table 2).

Figure 1 indicates a significant difference in the average decrease in HSP70 expression between group 4 and either group 1 or 2 ( $p=0.000$ ). However, a significant difference between group 4 and group 3 ( $p=0.813$ ) was not found. This means that group 4, with strenuous exercise and given 300 mg/kg body weight red-fleshed pitaya extract was indicated to be more effective for

**Table 1. Normality test for heat shock protein 70 (HSP70) and cortisol expression.**

Parameter	Group	Normality test	
		statistic	p-value
HSP70	Group 1	12	0.969 0.888
	Group 2	12	0.907 0.332
	Group 3	12	0.922 0.449
	Group 4	12	0.875 0.168
Cortisol	Group 1	12	0.923 0.451
	Group 2	12	0.862 0.833
	Group 3	12	0.939 0.603
	Group 4	12	0.900 0.290

Shapiro-wilk test,  $P>0.05$

**Table 2. Average heat-shock protein (HSP70) and cortisol expression in rats who were subjected to strenuous exercise and treated with distilled water (group 1), and 100 mg/ kg body weight (group 2), 200 mg/ kg body weight (group 3) and 300 mg/ kg body weight (group 4) red-fleshed pitaya extract, respectively.**

Parameter	24				p-value
	Group 1 (n=8)	Group 2 (n=8)	Group 3 (n=8)	Group 4 (n=8)	
HSP70 (pg/ml)	69.57 $\pm$ 7.27	46.04 $\pm$ 6.45	31.47 $\pm$ 1.24	27.65 $\pm$ 1.72	0.000*
Cortisol (pg/ml)	119.02 $\pm$ 5.56	86.11 $\pm$ 11.40	62.94 $\pm$ 4.61	40.86 $\pm$ 10.94	0.000*

The mean  $\pm$  SD HSP70 and cortisol expression is shown.

reducing HSP70 expression compared to group 1 and 2, with no big variance from group 3.

**Figure 2** confirms that a significant difference is revealed in the average decrease of cortisol expression between groups 4 and 1 ( $p=0.000$ ), 2 ( $p=0.000$ ), and 3 ( $p=0.000$ ). It can be concluded that group 4 with strenuous exercise and 300 mg/kg body weight red-fleshed pitaya extract was more effective in reducing cortisol expression compared to the other groups.

### Discussion


Based on the results of this research, it is found that strenuous exercise combined with daily red-fleshed pitaya extract consumption contributes to a declining expression of HSP70 and cortisol. The dosage of 300 mg/kg body weight red-fleshed pitaya extract was found to <sup>29</sup> the optimum amount in decreasing cortisol compared to <sup>100</sup> and <sup>200</sup> mg/kg body weight dosage. However, for both 300 mg/kg body weight and 200 mg/kg body weight dosage of red-fleshed pitaya extract could not provide any difference to HSP70 expression. Therefore, the red-fleshed pitaya extract can be categorized as a potential exogenous antioxidant to eliminate free radicals formed during strenuous exercise.

<sup>31</sup> Free radicals are an element that possesses one or more unpaired electrons in its outermost orbital. Consequently, it is very reactive to cells or cell components in its surroundings. Commonly, a reactive element finds its pair by attacking and binding with adjacent electrons. Then, if this element reacts with another radical element, a new radical element will be formed. This will consistently continue to occur leading to an unavoidable chain reaction<sup>10</sup>. During strenuous exercise, oxygen consumption rises by 20 times. The excess oxygen triggers the formation of free radicals with electrons released from the respiratory chain. Free radicals production from activity, especially superoxides, increases in the mitochondria<sup>20</sup>. When an imbalance happens due to the excess of free radicals, oxidative stress occurs and damages the DNA. Moreover, proteins will lose their structure and function like enzymes and membrane receptor. Also, there will be damage in the lipid bilayer structure<sup>21</sup>.


Strenuous exercise triggers oxidative stress to occur, therefore, HSP70 expression will elevate, resulting in a decrease of endogenous antioxidant activity. HSP70 is an important protein molecule for cell healing and preventing homeostasis, and also increased expression as a cyto-protective effect<sup>22</sup>. HSP70 is induced strongly due to oxidative stress as cyto-protective to prevent oxidative damage and heal broken proteins<sup>23</sup>. The increase of HSP70 expression is aimed to balance between ischemic condition, high temperature and increased production of free radical<sup>24</sup>. In this research, it is found that HSP70 expression tends to decrease in a group with strenuous exercise combined with red-fleshed pitaya extract.

The red-fleshed pitaya extract combined with strenuous exercise is proven to be able to inhibit the increasing HSP70 expression. This occurred due to the potential of red-fleshed pitaya as an antioxidant that is able to balance the increasing amount of free radicals formed and impairs HSP70 synthesis induced by strenuous exercise<sup>25</sup>. Therefore, HSP70 expression is lower compared to the group <sup>27</sup> that are not given by red-fleshed pitaya extract. The results of this research is in accordance to a research conducted by Petiz *et al.* (2017) which discovered that Vitamin A combined with high intensity activities could prevent tissue damage and reduce endogenous antioxidant defense regulation in rats, which also suppressed HSP70 expression<sup>26</sup>. Other researches have also revealed that antioxidant supplementation is proven to be effective to slow down the HSP70 synthesis caused by high intensity exercise<sup>27</sup>.

Physical training combined with low to medium dosage of red-fleshed pitaya extract were measured based on the ability and lead to immunomodulation effect that affect the body immune system and protect the body from cell damages, resulting in an effective condition to reduce the oxidative stress<sup>28</sup>. Stress resulted from physical activity like oxidative stress is responded by hypothalamus to secrete corticotrophin releasing hormone (CRH) which then delivers a message to pituitary anterior. The pituitary produced adrenocorticotrophic hormone (ACTH) which is useful to activate or affect adrenal cortex where



**Figure 1. Effect of red-fleshed pitaya extract on HSP70 expression in rats that were given strenuous exercise.**



**Figure 2.** Effect of red-fleshed pitaya extract on cortisol in rats that were given strenuous exercise.

cortisol hormone is secreted. Cortisol contributes a massive influence to immune responses<sup>13</sup> as a sign that the body is suffering from oxidative stress<sup>17</sup>. In this research, cortisol hormone level decreased in the group with strenuous exercise combined with red-fleshed pitaya extract.

This occurs since the red-fleshed pitaya extract is an effective antioxidant to reduce the risk of oxidative stress and is able to decrease the secretion of cortisol by reducing ACTH secretion in the hypothalamus and CRH in the pituitary gland<sup>18</sup>.

### Conclusion

Based on this research, it can be concluded that red-fleshed pitaya extract has the potential to be antioxidant with its anthocyanin content, and is able to eliminate oxidative stress due to strenuous exercise. It can be observed by the decreased pattern of HSP70 and cortisol expression in the strenuous exercise combined with red-fleshed pitaya extract group. Furthermore, the optimum result was shown in the dosage of 300 mg/kg body weight red-fleshed pitaya extract.

### Data availability

#### Underlying data

Open Science Framework: The effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise induced rats, <https://doi.org/10.17605/OSF.IO/MGX4K><sup>20</sup>

<sup>15</sup>

Data are available under the terms of the Creative Commons Zero "No rights reserved" data waiver (CC0 1.0 Public domain dedication).

### Funding information

The authors would like to acknowledge Ministry of Research, Technology and Higher Education of Republic of Indonesia for funding this research via Research Grant of Universitas Negeri Medan with contract number of 027/UN33.8/LL/2018.

<sup>14</sup>

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

### References

- Yavari A, Javadi M, Mimiran P, et al.: Exercise-induced oxidative stress and dietary antioxidants. *Asian J Sports Med.* 2015; 6(1): e24898. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Escribano BM, Tinez I, Requena F, et al.: Effects of an aerobic training program on oxidative stress biomarkers in bulls. *Vet Med (Praha)*. 2010; 55(9): 422–428. [Publisher Full Text](#)
- El Abed K, Rebai H, Bloomer RJ, et al.: Antioxidant status and oxidative stress at rest and in response to acute exercise in judokas and sedentary men. *J Strength Cond Res.* 2011; 25(9): 2400–9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Steinbacher P, Eckl P.: Impact of oxidative stress on exercising skeletal muscle. *Biomolecules*. 2015; 5(2): 356–77. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vaiko M, Leibfritz D, Moncol J, et al.: Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007; 39(1): 44–84. [PubMed Abstract](#) | [Publisher Full Text](#)
- Macedo DV, Lazarim FL, Catanho da Silva FO, et al.: Is lactate production related to muscular fatigue? A pedagogical proposition using empirical facts. *Adv Physiol Educ.* 2009; 33(4): 302–7. [PubMed Abstract](#) | [Publisher Full Text](#)
- Kürkü R, Tekin A, Özda S, et al.: The effects of regular exercise on oxidative and antioxidative parameters in young wrestlers. *African J Pharm Pharmacol.* 2010; 4(5): 244–251. [Reference Source](#)
- Azizbeigi K, Azarbayjani MA, Peerfi M, et al.: The effect of progressive resistance training on oxidative stress and antioxidant enzyme activity in erythrocytes in untrained men. *Int J Sport Nutr Exerc Metab.* 2013; 23(3): 230–8. [PubMed Abstract](#) | [Publisher Full Text](#)
- Folkesson M, Mackey AL, Langberg H, et al.: The expression of heat shock protein in human skeletal muscle: effects of muscle fibre phenotype and training background. *Acta Physiol (Oxf)*. 2013; 209(1): 26–33. [PubMed Abstract](#) | [Publisher Full Text](#)
- Liu Y, Gampert L, Nethsing K, et al.: Response and function of skeletal muscle

- heat shock protein 70.** *Front Biosci.* 2006; 11: 2802–27.  
[PubMed Abstract](#) | [Publisher FullText](#)
11. Biller-Takahashi JD, Takahashi LS, Mingotto FE, et al.: **The immune system is limited by oxidative stress: Dietary selenium promotes optimal antioxidative status and greatest immune defense in pacu *Piaractus mesopotamicus*.** *Fish Shellfish Immunol.* 2015; 47(1): 360–7.  
[PubMed Abstract](#) | [Publisher FullText](#)
  12. Staib JL, Turner N, Powers SK: **Increased temperature and protein oxidation lead to HSP72 mRNA and protein accumulation in the *in vivo* exercised rat heart.** *Exp Physiol.* 2009; 94(1): 71–80.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  13. Usui T, Yoshikawa T, Ueda SY, et al.: **Effects of acute prolonged strenuous exercise on the salivary stress markers and inflammatory cytokines.** *Jpn J Phys Fitness Sports Med.* 2011; 60(3): 295–304.  
[Publisher FullText](#)
  14. Hill EE, Zacki E, Battaglini C, et al.: **Exercise and circulating cortisol levels: the intensity threshold effect.** *J Endocrinol Invest.* 2008; 31(7): 587–91.  
[PubMed Abstract](#) | [Publisher FullText](#)
  15. Peternelj TT, Coombes JS: **Antioxidant supplementation during exercise training: beneficial or detrimental?** *Sports Med.* 2011; 41(12): 1043–69.  
[PubMed Abstract](#) | [Publisher FullText](#)
  16. Gomez-Cabrer MC, Vifia J, Ji LL: **Interplay of oxidants and antioxidants during exercise: implications for muscle health.** *Phys Sportsmed.* 2009; 37(4): 116–23.  
[PubMed Abstract](#) | [Publisher FullText](#)
  17. Belviranli M, Gokbel H: **Acute exercise induced oxidative stress and antioxidant changes.** *Eur J Gen Med.* 2006; 3(3): 126–131.  
[Publisher FullText](#)
  18. Nurul SR, Asmah R: **Variability in nutritional composition and phytochemical properties of red pitaya (*Hylocereus polyrhizus*) from Malaysia and Australia.** *Int Food Res J.* 2014; 21(4): 1689–1697.  
[Reference Source](#)
  19. Sahlén K, Shabalina IG, Mattsson CM, et al.: **Ultraendurance exercise increases the production of reactive oxygen species in isolated mitochondria from human skeletal muscle.** *J Appl Physiol (1985).* 2010; 108(4): 780–7.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  20. Gomes EC, Silva AN, de Oliveira MR: **Oxidants, antioxidants, and the beneficial roles of exercise-induced production of reactive species.** *Oxid Med Cell Longev.* 2012; 2012: 756132.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
  21. Abruzzo PM, Esposito F, Marchionni C, et al.: **Moderate exercise training induces ROS-related adaptations to skeletal muscles.** *Int J Sports Med.* 2013; 34(8): 676–87.  
[PubMed Abstract](#) | [Publisher FullText](#)
  22. Krause M, Heck TG, Bittencourt A, et al.: **The chaperone balance hypothesis: the importance of the extracellular to intracellular HSP70 ratio to inflammation-driven type 2 diabetes, the effect of exercise, and the implications for clinical management.** *Mediators Inflamm.* 2015; 2015: 249205.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  23. Lanneau D, Brunet M, Frisan E, et al.: **Heat shock proteins: essential proteins for apoptosis regulation.** *J Cell Mol Med.* 2008; 12(3): 743–61.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  24. Khassaf M, McArdle A, Esanu C, et al.: **Effect of vitamin C supplements on antioxidant defence and stress proteins in human lymphocytes and skeletal muscle.** *J Physiol.* 2003; 549(Pl.2): 645–52.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  25. Petz LL, Girardi CS, Bortolin RC, et al.: **Vitamin A Oral Supplementation Induces Oxidative Stress and Suppresses IL-10 and HSP70 In Skeletal Muscle of Trained Rats.** *Nutrients.* 2017; 9(4): pii: E353.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  26. Baghaiee B, Karimi P, Siahkouhian M, et al.: **Moderate aerobic exercise training decreases middle-aged induced pathologic cardiac hypertrophy by improving Klotho expression, MAPK signaling pathway, and oxidative stress status in Wistar rats.** *Iran J Basic Med Sci.* 2018; 21(9): 911–919.  
[PubMed Abstract](#) | [Publisher FullText](#) | [Free Full Text](#)
  27. Turner-Cobb JM, Palmer J, Aronson D, et al.: **Diurnal cortisol and coping responses in close relatives of persons with acquired brain injury: a longitudinal mixed methods study.** *Brain Inj.* 2010; 24(6): 893–903.  
[PubMed Abstract](#) | [Publisher FullText](#)
  28. Kraemer WJ, Ratamess NA: **Hormonal responses and adaptations to resistance exercise and training.** *Sports Med.* 2005; 35(4): 339–61.  
[PubMed Abstract](#) | [Publisher FullText](#)
  29. Harahap, Novita S: "The Effect of Red-Fleshed Pitaya (*Hylocereus Polyrhizus*) on Heat Shock Protein 70 and Cortisol Expression in Strenuous Exercise Induced Rats." OSF. 2019.  
<http://www.doi.org/10.17605/OSF.IO/MGX4K>

## Open Peer Review

### Current Peer Review Status:



Version 1

Reviewer Report 04 March 2020

<https://doi.org/10.5256/f1000research.19175.r59870>

8  
© 2020 Rohin M. This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

✓ Mohd Adzim Khalili Rohin 15

Faculty of Health Sciences, Sultan Zainal Abidin University (UniSZA), Kuala Nerus, Malaysia

1. It is an interesting manuscript. However, there are several grammatical issues and some typos that you should check very carefully.
2. My main concern about your manuscript is that you have not mentioned any strengths or limitations in your study.
3. The small number of rats per group is indeed a limitation and should be mentioned and taken into consideration before making conclusions.
- 2  
4. This is a prospective study on the effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise-induced rats.
5. The conclusion is weak in my opinion.
6. The study could be considered more as a pilot study than a proper study; however the results are interesting and should be indexed.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

5

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Functional Food & Nutraceuticals

1

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 08 November 2019

1

<https://doi.org/10.5256/f1000research.19175.v55B88>

1

© 2019 Yusni Y. This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Yusni Yusni**



Department of Physiology, Syiah Kuala University, Banda Aceh, Indonesia

**Introduction:**

1. Please explain why red-fleshed pitaya is thought to act as antioxidants that will affect the heat shock protein 70 and the expression of cortisol: include theoretical and other studies that support.  
20
2. The novelty of the article must be clearly explained in the introduction.

**Methods:**

1. Why use 3 doses, whereas the purpose of the study did not assess the effective dose? In the conclusion, also mention the effective dose.
2. Explain the results of the identification of the chemical content of red-fleshed pitaya and also the results of herbarium analysis because there is a possibility that Indonesia's geographical differences will affect its chemical content.
3. Explain the time of blood-sampling because cortisol production is influenced by circadian.

**Discussion:**

2. It is important to explain the role or mechanism of action of red-fleshed pitaya on heat shock protein 70 and cortisol expression.

**Reference:**

1. Use the latest references (from the last 5-10 years).

**Note:** There are some grammatical errors, you should use the proof-read service.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

5

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Medical Physiology: exercise physiology; endocrine and metabolic physiology

5  
I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response ID No. 18

Novita Sari Harahap, Faculty of Sports Sciences, Universitas Negeri Medan, Medan, Indonesia

**Introduction:**

1. Please explain why red-fleshed pitaya is thought to act as antioxidants that will affect the heat shock protein 70 and the expression of cortisol: include theoretical and other studies that support.

Red fleshed pitaya has antioxidant activity. Red Dragon Fruit contains powerful antioxidants because it consists of various antioxidants, namely flavonoids, vitamin E, vitamin C (Lianiwiati, 2011). The results of several studies stated the red dragon fruit (*Hylocereus polyrhizus*) has an anthocyanin content (Kanner et al., 2001) around 26,457 ppm. Dragon fruit contains flavonoids (Damayanti, 2016). Anthocyanin is a pigment that is classified as a type of water-soluble flavonoids. Anthocyanins are anthocyanidin and glucose compounds in organic acids (Shipp and Abdel Aal, 2010). Anthocyanins are believed to have a very good antioxidant effect. A study conducted at the University of Michigan in America showed that anthocyanin can destroy free radicals, more effectively than vitamin E which has been known as a powerful antioxidant. Other studies in the United States prove that anthocyanin is the most powerful antioxidant among other flavonoid classes. The content of anthocyanin is believed to inhibit various free radicals such as superoxide radicals and hydrogen peroxide. Anthocyanins and various forms of their derivatives can inhibit various oxidation reactions by various mechanisms (Astawan and Kasih, 2008). The results of previous studies ripe dragon fruit contains many organic acids (Stinzing, et al., 2004), protein (Le Bellec, et al., 2006), minerals such as potassium, magnesium, calcium and iron and

vitamin C (Cal et al., 2003).

Increased RONS (Reactive Oxygen and Nitrogen Species) production during exercise has potential negative effects, transient increases in RONS seem to be a trigger for many exercise-induced adaptations in skeletal muscle (Powers et al. 2010). Exercise can stimulate an increased concentration of HSP70 in muscles and plasma.

Oxidative stress due to exercise can cause physiological functional changes because physical exercise is considered as a stimulus received by the hypothalamus, then the hypothalamus signals to the HPA axis, the HPA axis responds and gives positive and negative responses to the body (Guilliam & Edward, 2010). Then stimulates the hypothalamus and causes secretion of the hormone corticotrophin-releasing hormone (CRH), which stimulates the hypothalamus for ACTH secretion. Increased ACTH secretion, causes increased cortisol secretion (Usui et al, 2012).

#### Methods:

1. Why use 3 doses, whereas the purpose of the study did not assess the effective dose? In conclusion, also mention the effective dose.

Giving 3 doses of 100 mg/kg BW, 200 mg/kg BW and 300 mg/kg BW to find out what is the optimal dose to neutralize oxidative stress due to strenuous physical exercise. A dose of 100 mg/kg BW can also reduce oxidative stress but not as well as a dose of 300 mg/kg BW. So it is necessary to experiment with 3 doses, the purpose of the study to find out the potential of Red-fleshed pitaya as an antioxidant and once to know the optimal dose to reduce oxidative stress.

2. Explain the results of the identification of the chemical content of red-fleshed pitaya and also the results of herbarium analysis because there is a possibility that Indonesia's geographical differences will affect its chemical content.

Red-fleshed pitaya is the cactus fruit of the genera *Hylocereus* and *Selenicereus*. Dragon fruit is very popular and widely planted in Indonesia because it is known by the public as herb medicine (Evi, et al., 2007). From the results of previous studies, ripe dragon fruit contains many organic acids (Stinzing, et al., 2004), protein (Le Bellec, et al., 2006), minerals such as potassium, magnesium, calcium and iron and vitamin C (Cal et al., 2003).

3. Explain the time of blood-sampling because cortisol production is influenced by circadian.

Cortisol levels are affected by circadian rhythm. Blood drawn for examination of cortisol is done in the morning because it is a basal state of the body after resting at night. Also besides, our bodies have biological variations according to time, meaning that the analytical levels examined in the morning can give different results if examined in the afternoon. On examination of Cortisol, the levels will increase in the morning and reach the lowest levels in the afternoon. Then the blood draw is done in the morning according to the normal human biological clock, the most efficient and does not result in excessive fatigue.

#### Discussion:

1. It is important to explain the role or mechanism of action of red-fleshed pitaya on heat shock

protein 70 and cortisol expression.

*Research conducted by Rebecca et al (2010) shows that red dragon fruit contains the most polyphenols compared to other species, namely  $86.13 \pm 17.02$  mg in 0.50 g of dried extract of red dragon fruit. Red dragon fruit (*Hylocereus polyrhizus*) is one of the plants that can be used as a source of antioxidants. Severe physical exercise is associated with increased HSP70 expression due to oxidative stress (Paulsen et al. 2007).*

*Oxidative stress causes damage to cells that will cause stress responses in the form of heat shock response (HSR). HSR is regulated at the level of transcription by a mechanism involving heat shock transcription factor (HSF), especially HSF-1. Increased HSF-1 will help in increasing HSP 70 expression in certain amounts/levels in serum and blood plasma (Baird et al., 2006).*

*Physical activity appears to occur concomitantly with the up-regulation of endogenous antioxidant systems (Powers and Jackson 2008) and heat shock proteins (HSP) in skeletal muscle (Morton et al. 2009b). While the main role of the antioxidant enzymes is to reduce oxidation and prevent oxidative damage, HSPs can prevent and reverse damage to proteins. Intriguingly, the HSPs work with the antioxidant systems, and collectively they have essential roles in cell homeostasis.*

*Up-regulation of these proteins is, therefore, important adaptations for increased protection and recovery capacity in the face of cellular stress and damage induced by high-intensity exercise (Powers and Jackson 2008; Morton et al. 2009b).*

*Antioxidants have been shown to abolish the acute exercise-induced increases in HSP70 protein levels (Khassaf et al. 2003; Jackson et al. 2004).*

*Increased cortisol levels after physical activity depends on the level of fitness, exercise intensity and exercise program [Leite et al., 2011]. Exercise with heavy intensity tends to produce free radicals which can cause the production of the hormone cortisol to increase (Stachowicz, 2016).*

*The provision of red dragon fruit that contains antioxidant polyphenols can reduce the production of free radicals due to exercise thereby reducing cortisol levels.*

**Competing Interests:** I declare no competing interests

#### The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact [research@f1000.com](mailto:research@f1000.com)

F1000Research

# The effect of red-fleshed pitaya (*Hylocereus polyrhizus*) on heat shock protein 70 and cortisol expression in strenuous exercise induced rats

ORIGINALITY REPORT



PRIMARY SOURCES

- 1 P Ravi Shankar, Kundan K Singh, Ajaya Dhakal, Arati Shakya, Rano M Piryani. "Student feedback about the use of literature excerpts in Sparshanam, a Medical Humanities module", F1000Research, 2012  
Publication 1 %
- 2 [sinta3.ristekdikti.go.id](http://sinta3.ristekdikti.go.id) 1 %  
Internet Source
- 3 [www.academicjournals.org](http://www.academicjournals.org) 1 %  
Internet Source
- 4 Submitted to University of Birmingham 1 %  
Student Paper
- 5 Poisot, Timothée, Thomas Bell, Esteban Martinez, Claire Gougam-Barbera, and Michael E Hochberg. "Terminal investment induced by a bacteriophage in a rhizosphere bacterium", F1000 Research, 2012.  
Publication 1 %

6

global.oup.com

Internet Source

1 %

7

www.omicsonline.org

Internet Source

1 %

8

Bettina Bewernick, Thomas E Schlaepfer.  
"Update on Neuromodulation for Treatment-  
Resistant Depression", F1000Research, 2015

Publication

9

Submitted to Universitas Muhammadiyah  
Surakarta

Student Paper

1 %

10

physreports.physiology.org

Internet Source

1 %

11

doaj.org

Internet Source

<1 %

12

mhrbeo.org

Internet Source

<1 %

13

N S Harahap, A Sunarno, N Simatupang,  
Suprayitno. " The Effect of Red Dragon Fruit  
Juice Towards Cholesterol Level and Maximum  
Aerobic Capacity (VO max) on Sport Science  
Students Treated with Heavy Physical Exercise  
", Journal of Physics: Conference Series, 2020

Publication

<1 %

14

www.frontiersin.org

Internet Source

<1 %

- 
- 15 Luck, Gary W, Kai MA Chan, and Carissa J Klien. "Identifying spatial priorities for protecting ecosystem services", F1000 Research, 2012. Publication <1 %
- 
- 16 15projects.com Internet Source <1 %
- 
- 17 Kenneth A. Moselle, Ernie Chang. "CovidSIMVL – Agent-Based Modeling of Localized Transmission within a Heterogeneous Array of Locations – Motivation, Configuration and Calibration", Cold Spring Harbor Laboratory, 2020 Publication <1 %
- 
- 18 Novita Sari Harahap, Nurhayati Simatupang, Suprayitno Suprayitno. "The Effect of Physical Activity and Red Dragon Fruit (*Hylocereus polyrhizus*) in Red Blood Cell and Hemoglobin in Trained People", Open Access Macedonian Journal of Medical Sciences, 2021 Publication <1 %
- 
- 19 Seyed Sajjad Eghbali, Samad Akbarzadeh, Afshar Bargahi, Kamran Mirzaei, Nasrollah Maleki, Mehrzad Bahtouee, Ahmad Yazdanpanah. "Aqueous extract of *Parkinsonia aculeata* has an anti-urolithiatic effect in rats", <1 %

# Research Square, 2021

Publication

- 
- 20 medsci.org **<1 %**  
Internet Source
- 
- 21 Giuseppe Musumeci, Ali Mobasher, Marta Anna Szychlinska, Rosa Imbesi, Paola Castrogiovanni. "Post-operative rehabilitation in osteoarthritis ", F1000Research, 2014 **<1 %**  
Publication
- 
- 22 Submitted to University of Newcastle **<1 %**  
Student Paper
- 
- 23 bestjournal.untad.ac.id **<1 %**  
Internet Source
- 
- 24 www.pubmedcentral.nih.gov **<1 %**  
Internet Source
- 
- 25 E.M. Bomhard, S.Y. Brendler-Schwaab, A. Freyberger, B.A. Herbold, K.H. Leser, M. Richter. " -Phenylphenol and its Sodium and Potassium Salts: A Toxicological Assessment ", Critical Reviews in Toxicology, 2008 **<1 %**  
Publication
- 
- 26 International Journal of Housing Markets and Analysis, Volume 9, Issue 2 (2016) **<1 %**  
Publication
- 
- 27 Novita Sari Harahap, Aznan Lelo, Ambrosius Purba, Diky Setya Diningrat. "Influence of **<1 %**

**Combination of Weight Training with Sport  
Massage on 200 Meter Runners", Journal of  
Medical Sciences, 2018**

Publication

- 
- 28 [dentistry.tbzmed.ac.ir](http://dentistry.tbzmed.ac.ir) <1 %  
Internet Source
- 29 L. Pari, M. Latha. " Antihyperglycaemic Effect of  
. Effect on Key Metabolic Enzymes of  
Carbohydrate Metabolism in Streptozotocin-  
Induced Diabetes ", Pharmaceutical Biology,  
2008 <1 %  
Publication
- 30 Sihombing, Eidi, Manihar Situmorang,  
Timbangen Sembiring, and Nasruddin  
Nasruddin. "The Development Of Mercury Ion  
Selective Electrode With Ionophore 7,16-Di-(2-  
Methylquinolyl)-1,4,10,13-Tetraoxa-7,16-  
Diazacyclooctadecane (DQDC)", Modern  
Applied Science, 2015. <1 %  
Publication
- 31 [brage.bibsys.no](http://brage.bibsys.no) <1 %  
Internet Source
- 32 [isindexing.com](http://isindexing.com) <1 %  
Internet Source
- 33 [library.oapen.org](http://library.oapen.org) <1 %  
Internet Source

34

[www.farmaciadimuria.it](http://www.farmaciadimuria.it)

Internet Source

<1 %

35

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

36

[www.science.gov](http://www.science.gov)

Internet Source

<1 %

37

"Heat Shock Proteins and Whole Body  
Physiology", Springer Science and Business  
Media LLC, 2010

Publication

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

Off