

The mitochondrial cytochrome c oxidase subunit I (COI) for identification of batoids collected from landed sites in Medan, Indonesia

MUFTI SUDIBYO¹, KHAIRIZA LUBIS^{1*}, ACHMAD FARAJALLAH², NISFA HANIM¹

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Medan, Jl. Willem Iskandar Pasar V, Medan 20221, North Sumatra, Indonesia. Tel.: +62-66-1-4002, Fax.: +62-66-13319, *email: khairizalubis@unimed.ac.id, khairizalubis15@gmail.com.

²Department of Biology, Faculty of Mathematics and Natural Sciences, Institut Pertanian Bogor, Jl. Meranti, Kampus IPB Damsara, Bogor 16680, West Java, Indonesia

Manuscript received: 5 June 2020. Revision accepted: 26 October 2020.

Abstract. Lubis K, Sudibyo M, Farajallah A, Hanim N. 2020. Short Communication: The mitochondrial cytochrome c oxidase subunit I (COI) for identification of batoids collected from landing sites in Medan, Indonesia. *Biodiversitas* 21: 5414-5421. Batoids are member of Elasmobranch subclass which consist of many species. Most of batoids species are overexploited, especially in Medan Indonesia. Up to presents, the information about diversity of rays on the east coast of North Sumatra, Indonesia was very limited. Therefore, this research aimed to investigate the diversity of rays on the east coast of North Sumatra. We examined the morphological trait of 82 individuals of batoid from three landing sites on the east coast of North Sumatra, namely: Tanjung Balai, Belawan, and Percut, then identify its species based on determination key. After that, we collected pectoral muscle tissue from an individual in each species which successfully identified to extract its genomic DNA. Molecular based identification was carried out by using DNA fragment from COI gene. The successfully amplified COI gene DNA fragment then was sequenced and analyzed. Based on morphological trait, we successfully identifying nine species of batoid, which is *Maculabatis gerrardi*, *Gymnura poecilura*, *Dasyatis zugei*, *Brevitrygon heterura*, *Neotrygon kuhlii*, *Hemitrygon bennettii*, *Rhinobatos jemberanensis*, *Rhinoptera javanica*, and *Taeniura lymma*. The result of identification based on COI gene DNA fragment was in congruent with morphological-based identification based on data BLAST-N and genetic distance value within same species. The nucleotide diversity within same species ranged from 0-15 nucleotide variants.

Keywords: COI gene, rays, Medan, Indonesia

INTRODUCTION

Indonesian marine waters is known as having the richest diversity of Elasmobranch fishes and also as the center of diversity in the world. Several studies on Elasmobranch diversity had been conducted. A total of 207 species from 44 families of Chondrichthyans were found in Indonesian marine waters, which consist of 109 sharks, 96 batoids, and two ghost sharks (chimera) (Fahmi 2010). Another study reported that at least 221 species of sharks and batoids found in Indonesian marine waters (Sadili et al. 2015). Recently, a study of elasmobranch in the North Sumatra reported that eight species of sharks and batoids were caught and landed in Belawan and Batubara (Fadhilah et al. 2019).

Batoid is one of member of Elasmobranch subclass and closely related to sharks. This group consists of fishes that had moderately to greatly flattened body (dorsoventrally), ventral gill slit; enlarged pectoral fin, lack of anal fin, and its enlarged pectoral fin were fused to the side of the head and snout (Compagno 1999; McEachran and de Calvarho 2002). The body and conjoined of head enlarged pectoral fin and snout forming a disc, that delimited from its tail (Compagno 1999; McEachran and de Calvarho 2002). Batoid disk's shape was varied there are wedge-shape, circular, oval, and rhomboidal (Compagno 1999). Besides

its shape, the size (length) of this group were varied from 100 mm to more than 7000 mm (Compagno 1999). As of 7 November 2015, the number of batoid species in the world which has been reported was 630 species (Wiegmann 2016). Several species of batoid had been going through taxonomic review, due to new feature (morphology or molecular) were found, i.e. *Himantura gerrardi*, *Dasyatis bennetti*, *Himantura walga* had been under taxonomic review and now were called *Maculabatis gerrardi*, *Brevitrygon walga* and *Hemitrygon bennettii*, respectively (Last et al 2016; Froese and Pauly 2020).

Morphology-based identification methods were used to identifying species of certain organisms based on morphological features. This identification process was done by comparing the morphological features of sample with morphological features description in the determination key. While this method were of the fastest and cheapest approach, there is certain circumstances that we need to collect or exploit more biological data to successfully identify species of sample, such as specimen species that has similar morphological trait with other species from the same genus, cryptic species, immature specimen, incomplete specimen and hybrid species (Dudgeon et al. 2012). In this circumstances DNA-based identification were solution to collect or exploit more biological data from specimen. DNA-based identification

method or DNA barcoding is a method that used standardized DNA fragment sequence to identify species of organism interest (Hebert et al. 2003a). The mitochondrial genome is accumulating high percentage of neutral mutations that can be helpful in species identification (Siddappa et al. 2013). Up to present, there is several DNA marker from mitochondrial DNA that was known can be used in DNA barcoding of fish, one of them is DNA fragmen from COI gene (Dudgeon et al. 2012). Due to commonly used as DNA marker in DNA barcoding (Dudgeon et al. 2012), the amount of data about COI gene in GenBank was huge (11680 data for shark and batoids) so we used DNA fragment from COI gene for this study.

Batoid fishes was known have high economic value such as it skin used for basic materials of shoes, bags and belts and also it oil, teeth and bones can also be used for medicinal ingredients and glue (Subrata et al. 2017). Based on fisherman explanation, batoid were popular good for expor (personal communication). Due to that, nowadays, batoid were overexploited (Sadili et al. 2015). While the situation is like that, up to present the information about batoid in Indonesia were limited especially molecular information. In North Sumatra the information about batoid that available is about diversity of batoid that inhabit marine water in there (Puteri et al. 2017; Fadhillah et al. 2019). Herein, the present study is devoted to species identification and diversity of batoids in the east coast of North Sumatra, Indonesia based on mitochondrial cytochrome c oxidase subunit I (COI) gene.

MATERIALS AND METHODS

Flesh sample collection and morphological identification

The present study was conducted from September 2019 to October 2019. A total of 82 individuals of batoid fishes, which caught by local fishermen in Malaka Strait area, at 3 landing site in the east coast of North Sumatra, namely Belawan, Tanjung Balai and Percut (Figure 1) were examined for its morphological features and were identified for its species based on White et al (2006). Small part of pectoral fin muscle tissue under the skin from one individual from each species, which has successfully identified, were collected. The muscle tissue sample, then, were placed in 15 mL tube, after that alcohol 96% were added until muscle tissue submerged.

Table 1. Reference DNA sequence of COI gene of batoids

Species	Accession number
<i>Brevitrygon heterura</i>	MG792099
<i>Dasyatis zugei</i>	EU398759
<i>Dasyatis bennetti</i>	KF604910
<i>Himantura gerrardi</i>	MH230945
<i>Neotrygon kuhlii</i>	KU497951
<i>Taeniura lymma</i>	KM881715
<i>Gymnura poecilura</i>	EU398804
<i>Rhinoptera javanica</i>	JF494383
<i>Rhinobatos jimbaramensis</i>	EU398994




Figure 1. Map of three landed sites of batoids in east Sumatra coast of North Sumatra Province, Indonesia; purple circle is Belawan landing site, orange circle is Percut landing site and green circle is Tanjung Balai landing site

Genomic DNA extraction

Small parts from preserved samples were cut with a sterile surgical scissor, put in the 1.5 mL tube, then soaked in sterile water to remove the preserving liquid for 20-30 minutes. After that, the tubes containing small parts of sample were centrifuged 5000 rpm for 1 minute then the sterile water was removed. The tissue samples were cut to fine pieces in the tubes with a sterile surgical scissor. Then, genomic DNA extraction was carried out using DNA GENE AID Genomic DNA Mini Kit (Tissue) (Geneaid, Canada) and were done according to the protocol of the manufacturer.

Polymerase Chain Reaction (PCR) and Sequencing

The mitochondrial *COI* gene was amplified by polymerase chain reaction (PCR) method using the forward primer AF585 5'-ACCAACCACAAAGACATTGGCAC and the reverse primer AF586 5'-ACTTCTGGGTGCCAAAG AATCA, which was modified from primer C_FishF11 5'-TGTA AAAACGACGGCCAGTCGACTAATCATAAAGA TATCGGCAC and C_FishR11 5'-CAGGAAACAGCTAT GACACCTCAGGGTGTCCGAARAAYCARAA Ward et al. 2005). Amplification of *COI* gene fragment was conducted by using GoTaq Green Mastermix® I SA Polymerase (Promega) with the following steps: pre-denaturation for 3 minutes at 95°C, followed by denaturation, 30 cycles at 95 °C for 1 minute, annealing for 15 seconds at 52°C, and an extension for 10 minutes at 72°C. All products of PCR were separated using polyacrylamide gel electrophoresis and stained using silver staining method (Byun et al. 2009). These products of PCR were sent to 1st Base for DNA sequencing.

Molecular identification and species diversity analysis

The DNA sequencing products were displayed and edited using BioEdit 7.0.9.0 software (Hall 1999) and followed by aligning sequence from primer forward and reverse from the same sample using ClustalW which embedded in MEGA7 software (Kumar et al. 2016). After that, we combined/contig the two sequences to produce the *COI* gene DNA sequence that was ready to be analyzed further. To identify its species, firstly similarity analysis was conducted, using the GenBank BLAST-N program (Altschul et al. 1990) to compare *COI* gene DNA fragments from each sample with other mitochondrial DNA fragments in database. After we get the BLAST-N result, the top one sequence was downloaded and used as reference sequence (Table 1), then we aligned all *COI* gene DNA sequence from all sample with it using ClustalW which embedded in MEGA7 software using then we conduct genetic distance analysis with p-distance method to further identifying species of each sequence. Besides that, we also conducting nucleotide variation and construct phylogenetic tree. The phylogenetic tree was constructed using Neighbour-Joining for statistical methods (Saitou and

Nei 1987) with Kimura 2-Parameter for substitution model (Kimura 1980) and bootstrap method for test of phylogeny (Felsenstein 1985) with number of bootstraps was 1000. Neighbor-Joining method with nucleotide substitution model Kimura 2-parameter was used due to this option was the most used option to analyze the diversity pattern of *COI* gene (Hajibabei et al. 2006; Puillandre 2009)

RESULTS AND DISCUSSION

Results

Eighty-two batoids, which were examined its morphological features directly in the three landing sites, were successfully identified belonged to 2 ordo, 4 family, and 9 species (Table 2). Morphological features of the 9 species of batoid were presented in Table 3. Species diversity that we found in this study were different from the previous study of Fadhillah et al. (2019) which also took place in North Sumatra, around Malaka Strait. Fadhillah et al. (2019) only found 6 species of batoid while we found nine (3 species more). Only two species of batoid that were found both in this study and Fadhillah et al. (2019) that is *T. lynna* and *N. kuhlii*, while the other seven species were only found in this study. *Brevitrygon heterura* was batoid species with the most numbered individual which we found in these three landing sites (Belawan, Percut, and Tanjung Balai) and this species had different size of disc width (DW) and tail length (CL) with *B. walga* (*B. heterura* was species of *B. walga* that inhabit in Indonesian marine water) that was described by Simpfendorfer et al. (2017). The size of disc width (DW) and tail length (CL) of this species in our study are 17.14 ± 1.92 cm and 21.36 ± 4.87 respectively, while in Simpfendorfer et al. (2017) is 23-32 cm and 23-50.6 cm (1-2.2 times DW) respectively. We also found 1 individual of *Rhinobatos jimbaranensis*, which were previously reported by Last et al. (2006) as endemic species in Jimbaran Bay (Southern Bali).

Table 2. List of successfully identified batoid fishes and individual frequency of each species which found in three landed site

Order	Species	Number of individuals
Family		
Myliobatiformes		
Dasyatidae	<i>Brevitrygon heterura</i>	28
	<i>Dasyatis sayi</i>	6
	<i>Hemitrygon bennetti</i>	8
	<i>Maculobatis gerrardi</i>	4
	<i>Neotrygon kuhlii</i>	16
	<i>Taeniura lynna</i>	6
Gymnuridae	<i>Gymnura poeciliaria</i>	2
Myliobatidae	<i>Rhinoptera javanica</i>	11
Rajiformes		
Rhinobatidae	<i>Rhinobatos jimbaranensis</i>	1

Table 3. Morphological features of nine batoid species

Species	Qualitative morphological features	Quantitative morphological features				
		W (g)	TL (cm)	CL (cm)	DL (cm)	DW (cm)
<i>Brevitrygon heterura</i>	<p>1) skin fold on ventral surface of tail</p> <p>Profile of disc somewhat oval</p> <p>Tail short, not whip-like (end bulbous in adult females)</p> <p>Tail thorns very elongate, bases nearly half eye diameter in length</p> <p>Mid-disc thorns absent or rudimentary</p> <p>Low skin fold present on ventral surface of tail</p> <p>Tail not banded posterior to sting</p> <p>Snout extremely elongate</p> <p>Anterior margin of disc distinctly concave</p> <p>No oral papillae</p> <p>Skin fold present on ventral surface of tail</p> <p>The hind of tail to sting is not striped</p> <p>Very long tail</p> <p>Very small eyes</p> <p>Enlarged thorns were found on the middle of disk and along the tail</p> <p>Whip-like tail</p> <p>There are small denticles skin in the middle of dorsal</p> <p>Tail is covered by small thorn</p> <p>No skin fold on ventral surface of tail</p> <p>Profile of disc quadrangular</p> <p>Tail whip-like, with alternating light and dark bands</p> <p>Upper surface greyish brown with numerous white spots, variably distributed over disc (from almost plain to fully spotted)</p> <p>Low skin fold present on ventral surface of tail</p> <p>Low skin fold on dorsal tail beyond sting</p> <p>Tail with broad black and white bands, not whip-like</p> <p>Snout short with black bar through eyes</p> <p>Large, bright, blue spots on upper disc</p> <p>Usually no thorns on tail before sting</p> <p>Central skin fold on tail moderately deep, extending to tail tip</p> <p>Profile of disc oval</p> <p>Upper surface with bright blue spots</p> <p>Stings present (usually 2), located near end of tail</p> <p>Tail with a long blue stripe extending alongside before sting</p> <p>Dorsal fin absent</p> <p>Tail lacking a sting, and with ~9 dark bands</p> <p>Dorsal surface usually plain, sometimes with faint pale spots</p> <p>The snout strongly notched medially to form two lobes</p> <p>The tail is short</p> <p>The head is rather wide</p> <p>Have dark spots on dorsal area, not white spot</p> <p>Lower lobe of the tail is short</p> <p>Denticles along the midline of dorsal area are not clear</p> <p>Slightly large nostril</p>	187.86 ± 83.64	36.61 ± 5.72	21.36 ± 4.87	18.55 ± 2.29	17.14 ± 1.92
<i>Dasyatis zugei</i>		255.00 ± 220.43	46.33 ± 12.74	27.17 ± 7.39	21.83 ± 5.67	19.25 ± 4.12
<i>Hemirhynchon bonasus</i>		460.00 ± 278.23	50.48 ± 8.49	31.16 ± 5.46	20.31 ± 4.63	25.06 ± 4.20
<i>Maculabatis gerraulti</i>		375 ± 150	79 ± 6.78	62 ± 7.79	20.3 ± 1.71	21.4 ± 2.18
<i>Neotrygon kuhlii</i>		460.00 ± 278.23	50.48 ± 8.49	31.16 ± 5.46	20.31 ± 4.63	25.06 ± 4.20
<i>Taeniana lynnae</i>		655.0 ± 68.63	61.2 ± 3.25	37.7 ± 2.34	27.0 ± 2.28	43.09 ± 3.14
<i>Gymnura poecilura</i>		720	39	1845	24	45
<i>Rhinoptera javanica</i>		1,315.45 ± 314.69	66.18 ± 6.52	40.45 ± 5.37	30.27 ± 2.83	43.09 ± 3.14
<i>Rhinobatos jimbaramensis</i>		1300	77	43	40	24

Note: W: weight; TL: Total Length; CL: Tail Length; DL: Disc Length; DW: Disc Width; g: gram; cm: centimeter.

Table 5. Genetic distance of COI gene fragment between nine batoid species

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
1																			
2	0.201																		
3	0.182	0.134																	
4	0.146	0.191	0.174																
5	0.168	0.166	0.157	0.177															
6	0.190	0.163	0.149	0.180	0.128														
7	0.199	0.199	0.207	0.187	0.188	0.184													
8	0.179	0.191	0.184	0.161	0.165	0.172	0.212												
9	0.199	0.203	0.207	0.176	0.203	0.204	0.223	0.204											
10	0.000*	0.201	0.182	0.146	0.168	0.190	0.199	0.179	0.199										
11	0.204	0.003*	0.134	0.195	0.169	0.166	0.203	0.191	0.206	0.204									
12	0.184	0.136	0.002*	0.176	0.158	0.149	0.209	0.185	0.209	0.184	0.136								
13	0.142	0.193	0.176	0.003*	0.176	0.180	0.188	0.161	0.177	0.142	0.196	0.177							
14	0.166	0.168	0.158	0.176	0.002*	0.127	0.190	0.163	0.201	0.166	0.171	0.160	0.174						
15	0.185	0.163	0.152	0.176	0.128	0.006*	0.179	0.171	0.206	0.185	0.166	0.152	0.176	0.127					
16	0.199	0.199	0.207	0.187	0.188	0.184	0.000*	0.212	0.223	0.199	0.203	0.209	0.188	0.190	0.179				
17	0.185	0.191	0.180	0.168	0.168	0.166	0.210	0.006*	0.207	0.185	0.191	0.182	0.168	0.166	0.165	0.210			
18	0.207	0.204	0.198	0.182	0.206	0.201	0.222	0.195	0.024*	0.207	0.207	0.199	0.184	0.207	0.203	0.222	0.198		

Note: * - genetic distance between same species; 1. *B. heterura*; 2. *D. zugei*; 3. *Hemirhynchon bennerii*; 4. *M. gerrardi*; 5. *N. kuhlii*; 6. *T. lymanza*; 7. *G. poecilura*; 8. *Rhinoptera javanica*; 9. *Rhinobatos jimbaramensis*; 10. MG792099 *B. heterura*; 11. EU398759 *D. zugei*; 12. KT504910 *D. bennerii*; 13. M1230945 *Himantura gerrardi*; 14. KU497951 *N. kuhlii*; 15. KM881715 *T. lymanza*; 16. EU398804 *G. poecilura*; 17. MG792069 *Rhinoptera javanica*; 18. EU 398994 *Rhinobatos jimbaramensis*




Figure 2. Phylogenetic tree of nine batoid species which found in three landed site based on COI gene fragment constructed using Neighbour-joining method with Kimura 2-Parameter model and 1000 bootstrap

Table 4. BLAST-N result of nine batoid species which found in three landed site

Organism species of sample sequence	Organism species of reference sequence	Accession Number	Query Cover	E-Value	Identity percentage
<i>B.heterura</i>	<i>Brevitrygon heterura</i>	MG792099	99%	0.0	100.00%
<i>D. zugei</i>	<i>Dasyatis zugei</i>	EU398759	96%	0.0	99.68%
<i>Hemirhynchus bennetti</i>	<i>Dasyatis bennetti</i>	KF604910	99%	0.0	99.85%
<i>M. gerrardi</i>	<i>Himantura gerrardi</i>	MH230945	100%	0.0	99.69%
<i>N. kuhlii</i>	<i>Neotrygon kuhlii</i>	KU497951	100%	0.0	99.85%
<i>T. lymma</i>	<i>Taeniura lymma</i>	KM881715	100%	0.0	99.39%
<i>G. poecilura</i>	<i>Gymnura poecilura</i>	EU398804	99%	0.0	100.00%
<i>Rhinoptera javanica</i>	<i>Rhinoptera javanica</i>	JF494383	99%	0.0	99.39%
<i>Rhinobatos jimbaranensis</i>	<i>Rhinobatos jimbaranensis</i>	EU398994	99%	0.0	97.55%

Table 6. Nucleotide variant of COI gene fragment between same species

Sample >< Reference Sample	Total number of nucleotide variant
<i>B. heterura</i> >< MG792099 <i>B. heterura</i>	0
<i>D. zugei</i> >< EU398759 <i>D. zugei</i>	2
<i>Hemirhynchus bennetti</i> >< KF604910 <i>D. bennetti</i>	1
<i>M. gerrardi</i> >< MH230945 <i>Himantura gerrardi</i>	2
<i>N. kuhlii</i> >< KU497951 <i>N. kuhlii</i>	1
<i>T. lymma</i> >< KM881715 <i>T. lymma</i>	4
<i>G. poecilura</i> >< EU398804 <i>G. poecilura</i>	0
<i>Rhinoptera javanica</i> >< MG792069 <i>Rhinoptera javanica</i>	4
<i>Rhinobatos jimbaranensis</i> >< EU398994 <i>Rhinobatos jimbaranensis</i>	15

Table 7. Batoids conservation status by International Union for Conservation of Nature (IUCN) and trade status by Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) in 2019.

Species	IUCN	CITES
<i>Rhinobatos jimbaranensis</i>	Vulnerable	Not evaluated
<i>G. poecilura</i>	Near threatened	Not evaluated
<i>Rhinoptera javanica</i>	Vulnerable	Not evaluated
<i>M. gerrardi</i>	Vulnerable	Not evaluated
<i>T. lymma</i>	Near threatened	Not evaluated
<i>B. heterura</i>	Not Evaluated	Not evaluated
<i>D. zugei</i>	Near threatened	Not evaluated
<i>N. kuhlii</i>	Data deficient (DD)	Not evaluated
<i>Hemirhynchus bennetti</i>	Not Evaluated	Not evaluated

We successfully get *COI* gene fragment sequence from nine species of batoid which had been identified based on morphological features. The length of *COI* gene fragment after we conduct editing and combining primer forward and reverse sequence was 655 bp. By using this 655 bp *COI* gene fragment we conduct similar analysis using BLAST-N, the result of this analysis can be seen in Table 4. The result of BLAST-N in all sample sequences above 97 % in "Identity Percentage" value, the lowest was *Rhinobatos jimbaranensis* (97.55%) and the highest was *B. heterura* and *G. poecilura* (100.00%). After we aligning sequence sample and reference sequence, the length of *COI* fragment sequence which can be used in further analysis was 632 bp. Within same species, the range of genetic distance was 0.000-0.024, while between different species was more than 0.1. The lowest value of genetic distance (0.000) was found in *G. heterura* and *G. poecilura*, while the highest was in *Rhinobatos jimbaranensis*.

Our result in nucleotide diversity analysis showed that between same species, nine batoids species which we found, differ in 0-15 nucleotide (Table 6). The lowest number of variant nucleotides found in *B. heterura* and *G. poecilura* (0 nucleotide variant), while the highest (15 nucleotide variant) were found in *Rhinobatos jimbaranensis*. The phylogenetic tree showed that Dasyatidae was divided into two clades, one clade was consist of *B. heterura*, *M. gerrardi*, *Rhinoptera javanica*, and *Rhinobatos jimbaranensis*, and the other clade consist of *D. zugei*, *Hemirhynchus bennettii*, *N. kuhli*, *T. lymma* and *G. poecilura* (Figure 2). Even though Dasyatidae divided into 2 clades, in each clade, every species of Dasyatidae were forming smaller clade, separated from the other family. *Maculabatis gerrardi* were forming a small clade with *B. heterura* and *D. zugei* were forming a small clade with *Hemirhynchus bennettii*, *N. kuhli*, and *T. lymma*.

Based on these results, we conclude that from 82 individuals we found in three landed sites (Tanjung Balai, Belawan, and Percut) were successfully identified belonged to nine species of batoid, both using morphological features identification and DNA identification using *COI* gene fragment. The mitochondrial cytochrome c oxidase subunit I (*COI*) gene identification could be used to evaluate batoid diversity, monitoring its conservation and fisheries management. This study about the use of DNA fragment to identifying batoid species needs further investigation by using another mitochondrial gene fragment to strengthening this finding. Furthermore, almost all batoid species which we found in these three landing sites its population is already.

ACKNOWLEDGEMENTS

This research was funded by Universitas Negeri Medan, Research Expertise Lecturer Group in 2019 budget year, No. 292D/UN33.8/PL/2019.

REFERENCES

- Almerón-Souza F, Sperb C, Castilho CL, Figueiredo PICC, Gonçalves LT, Machado R, Oliveira LR, Valati VH, Fagundes NJR. 2018. Molecular identification of shark meat from local markets on Southern Brazil based on DNA barcoding: evidence for mislabeling and trade of endangered species. *Front Genet* 9 (138): 1-12.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol*. 215 (3): 403-410, 10.1016/S0022-2836(05)80360-2.
- Blanco M, Perez-Martin RI, Sotelo CG. 2008. Identification of shark species in seafood products by forensically informative nucleotide sequencing (FINS). *J Agric Food Chem* 56: 9858-9874.
- Byun SO, Fang Q, Zhou H, Hickford JG. 2009. An effective method for silver-staining DNA in large numbers of polyacrylamide gels. *Anal Biochem*. 385 (1): 174-5.
- Ceratti-Pereira F, Meekun MG, Wei NWV, O'Shea O, Bradshaw CJA, Austin CM. 2012. Identification of Rays through DNA Barcoding: An Application for Ecologists. *PLOS ONE* 7 (6): 1-10.
- Compagnon LJV. 1999. Batoid fishes. In: Carpenter KE (eds) *The Living Marine Resources of the Western Central Pacific*, Vol 3: Batoid Fishes, Chimaeras, and Bony Fishes Part 1 (Elopidae to Linophrynidae). FAO, Rome.
- Dudgen CL, Blower DC, Broderick D, Giles JJ, Holmes BJ, Kashiwagi T, Krick NC, Morgan JAT, Tillet BJ, Overden JR. 2012. A review of the application of molecular genetics for fisheries management and conservation of sharks and rays. *J Fish Biol* 80: 1789-1843.
- Fadhilah A, Susetya, IE, \ Simoon BM. 2019. Elasmobranch catch composition of North Sumatra fishers. *IOP Conference Series: Earth and Environmental Science*. 260 (2109): 1-7.
- Fahmi. 2010. Sharks and rays in Indonesia. *Marine Research in Indonesia*. 35(1): 43-54.
- Felsenstein J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39:783-791.
- Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PDN. 2006. DNA barcodes distinguish species of tropical Lepidoptera. *PNAS* 103: 968-971.
- Hall TA. 1998. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser* 41: 95-98.
- Hebert PDN, Cywinska A, Ball SL, DeWaard JR. 2003a. Biological identifications through DNA barcodes. *Proceeding of the Royal Society B*. 270: 313-321.
- Hebert PDN, Ratnasingham S, deWaard JR. 2003b. Barcoding animal life: cytochrome c oxidase subunit I divergence among closely related species. *Proceeding of the Royal Society of London B*. 270: S96-S99.
- Hoelzel A. 2001. Shark fishing in fin soup. *Conservation Genetics*. 2: 69-72.
- Kimura M. 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16:111-120.
- Kinakesi SM, Wahyudewantoro G. 2017. Kajian jenis ikan Pari (Dasyatidae) di Indonesia. *Fauna Indonesia*. 16 (2): 17-25. [Indonesia]
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33 (7): 1870-1874.
- Last PR, Naylor GJ, Manjaji-Matsumoto BM. 2016. A revised classification of the family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insight. *Zootaxa*. 4193 (3): 345-368.
- Last PR, Naylor GJP, Manjaji-Matsumoto BM. 2016. A revised classification of the Family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insights. *Zootaxa*. 4139 (3): 345-368.
- Last PR, White WT, Fahmi. 2006. *Rhinobatos jimbaranensis* and *R. perigalli*, two new shovelnose rays (Batoidea: Rhinobatidae) from Eastern Indonesia. *Cybiu*. 30 (3): 261-271.
- Lim KC, Lim P-E, Chong VC, Loh K-H. 2015. Molecular and morphological analyses reveal phylogenetic relationships of stingrays focusing on the Family Dasyatidae (Myliobatiformes). *PLoS One*. 10 (4): e0120518. DOI: journal.pone.0120518.
- Maddappa H, Ayuningtyas RU, Subhan B, Anfar D, Prahadi. 2018. Exploited but unevaluated: DNA barcoding reveals skates and

- stingrays (Chordata, Chondrichthyes) species landed in the Indonesian fish market. *Ilmu Kelautan*, 21 (1): 29-36.
- McEachran JD, de Carvalho MR. 2002. Batoid fishes. In: Carpenter KE (eds) *The Living Marine Resources of the Western Central Pacific, Vol 1: Introduction, Molluscs, Crustaceans, Hagfishes, Sharks, Batoid Fishes and Chimaeras*. FAO, Rome.
- Puillandre N, Strong EE, Bouchet P, Boisselier MC, Couloux A, et al. 2009. Identifying gastropod spawn from DNA barcodes: possible but not yet practicable. *Mol Ecol Resour* 9: 1211-1221.
- Puteri D, Sitrus H, Muhtadi A. 2017. Fish diversity at mangrove ecosystem of Jaring Halus Village in Langkat Regency, North Sumatra. *Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 6 (2): 145-152.
- Rathipriya A, Marx KK, Jayashakila R. 2019. Molecular identification and phylogenetic relationship of flying fishes of Tamil Nadu coast for fishery management purposes. *Mitochondrial DNA Part A*, 3 0(3): 500-510.
- Riyanto B, Nurhayati T, Pujastuti AD. 2013. Karakterisasi Glikosaminoglikan dari Tulang Rawan Ikan Pari Air Laut (*Neotrygon kuhlii*) dan Pari Air Tawar (*Himantura signifer*). *Jurnal Pengolahan Hasil Perikanan Indonesia*, 16 (3): 224-232. [Indonesian]
- Rodrigues Filho LFdaS, Feitosa LM, Silva Nunes JL, Onodera Palmeira AR, Martins APB, Giarrizzo T, Carvalho-Costa LF, Monteiro ILP, Gemaque R, Gomes F, Souza RFC, Sampaio I, Sales JBdeL. 2020. Molecular identification of ray species traded along the Brazilian Amazon coast. *Fisheries Research*, 223 (105407): 1-10.
- Rodrigues Filho LFdaS, da Rocha TC, do Rego PS, Schneider H, Sampaio I, Vallinoto M. 2009. Identification and phylogenetic inferences on stocks of sharks affected by the fishing industry off the Northern coast of Brazil. *Genetics and Molecular Biology*, 32: 405-413.
- Sadili D, Dharmadi, Fahmi, Saminohadi, Ramli I, Sudarsono. 2015. Rencana Aksi Nasional (RAN) Konservasi dan Pengelolaan Hiu dan Pari. Direktorat Konservasi dan Keanekaragaman Hayati Laut, Ditjen Pengelolaan Ruang Laut, Kementerian Kelautan dan Perikanan, Indonesia. www.kkp.go.id
- Saitou N, Nei M. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406-425.
- Siddappa CM, Saini M, Das A, Doseswamy R, Sharma, AK, Gupta, PK. 2013. Sequence characterization of mitochondrial 12S rRNA Gene in Mouse Deer (*Moschiola indica*) for PCR-RFLP Based Species Identification. *Molecular Biology International*, 2013: 783925. DOI: 10.1155/2013/783925.
- Simpfendorfer C, Moore A., Elhassan I., Owfi F, Akhilesh KV. 2017. *Brevitrygon walga*. The IUCN Red List of Threatened Species 2017: e.T104176764A111015783. DOI: 10.2305/IUCN.UK.2017-2.REITS.T104176764A111015783.en.
- Subrata A, Wulandari D, Rizalinda. 2017. Inventarisasi Jenis Ikan Subkelas Elasmobranchii di Teluk Nuri Kecamatan Pulau Maya Kabupaten Kayong Utara. *Protobiot*, 6 (2): 45-49.
- Tinti F, Ungaro N, Pasolini P, De Paulis M, Garza F, Guarniero I, Sabelli B, Marano G, Piccinetti C. 2003. Development of molecular and morphological markers to improve species-specific monitoring and systematics of Northeast Atlantic and Mediterranean skates (Rajiformes). *J Exp Mar Biol Ecol* 288: 149-165.
- Ward RD, Zentak TS, Innes BH, Last PR, Hebert PD. 2005. DNA barcoding Australia's fish species. *Phil Trans R Soc B Biol Sci* 360 (1462): 1847-1857.
- Weigmann S. 2016. Annotated checklist of the living sharks, batoids and chimeras (Chondrichthyes) of the world, with a focus on biogeographical diversity. *J Fish Biol* 88: 837-1037.
- White WT, Last PR, Stevens JD, Yearsley GK, Fahmi, Dharmadi. 2006. Economically important sharks and rays of Indonesia. Lamb Print, Perth, WA.
- Wijayanti F, Abrari MP, Fitriana N. 2018. Keanekaragaman spesies dan status konservasi ikan pari di Tempat Pelelangan Ikan Muam Angke Jakarta Utara. *Jurnal Biodjati* 3 (1): 23-35. [Indonesian]
- Zhang J, Hamer R. 2012. Molecular Approach to the Identification of fish in the South China Sea. *PLoS One* 7 (2): e30621. DOI: 10.1371/journal.pone.0030621.

Nanopartikel-4

ORIGINALITY REPORT

17%

SIMILARITY INDEX

14%

INTERNET SOURCES

5%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	pt.scribd.com Internet Source	6%
2	Submitted to Universitas Jenderal Soedirman Student Paper	4%
3	biodiversitas.mipa.uns.ac.id Internet Source	1%
4	www.cvwd.org Internet Source	1%
5	Kevin McDonough. "Mammals of Mexico", Reference Reviews, 2016 Publication	<1%
6	protocolexchange.researchsquare.com Internet Source	<1%
7	N H Darlan, B D A Nugroho, S S Arif, P Sudira. "Spatial and temporal change of rainfall pattern in east coast of North Sumatera", IOP Conference Series: Earth and Environmental Science, 2019 Publication	<1%

8	Z A Harahap, H K Nasution, R Leidonald. " Relationship of spatial and temporal characteristics of water conditions and production in the Malacca Strait using satellite imagery ", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %
9	tubbatahareef.org Internet Source	<1 %
10	worldwidescience.org Internet Source	<1 %
11	www.hindawi.com Internet Source	<1 %
12	mafiadoc.com Internet Source	<1 %
13	Bárbara Serra-Pereira. "Molecular barcoding of skates (Chondrichthyes: Rajidae) from the southern Northeast Atlantic : Barcoding of skates from the southern Northeast Atlantic", Zoologica Scripta, 01/2011 Publication	<1 %
14	www.tandfonline.com Internet Source	<1 %
15	www.jove.com Internet Source	<1 %

16 Fangfang Zhao, Huitong Zhou, Shaobin Li, Qian Fang, Yuzhu Luo, Jon G. H. Hickford. "Growth and carcass trait association with variation in the somatostatin receptor 1 (SSTR1) gene in New Zealand Romney sheep", New Zealand Journal of Agricultural Research, 2018
Publication <1%

17 www.researchsquare.com
Internet Source <1%

18 S. Weigmann. "Reply to Borsa (2017): Comment on 'Annotated checklist of the living sharks, batoids and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity by Weigmann (2016)'", Journal of Fish Biology, 2017
Publication <1%

19 Dharmadi Dharmadi, Fahmi Fahmi, William White. "BIODIVERSITY OF SHARKS AND RAYS IN SOUTH EASTERN INDONESIA", Indonesian Fisheries Research Journal, 2017
Publication <1%

20 M Sirait, K Sinulingga, N Siregar, R S D Siregar. "Synthesis of hydroxyapatite from limestone by using precipitation method", Journal of Physics: Conference Series, 2020
Publication <1%

21

Internet Source

<1%

22

A Fadhilah, I E Susetya, B M Simeon. " catch composition of North Sumatera Fishers ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1%

23

ejournal-balitbang.kkp.go.id

Internet Source

<1%

24

"Abstracts of Papers Presented at the Fourteenth Annual Meeting", The Journal of Asian Studies, 2011

Publication

<1%

25

Nur Fadli, Siti Azizah Mohd Nor, Ahmad Sofiman Othman, Hizir Sofyan, Zainal A. Muchlisin. "DNA barcoding of commercially important reef fishes in Weh Island, Aceh, Indonesia", PeerJ, 2020

Publication

<1%



Exclude quotes On

Exclude matches Off

Exclude bibliography On

Nanopartikel-4

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8



THE
Character Building
UNIVERSITY