

# Nutritional Practices and Body Mass Index Among Secondary School Students

*by Syed Kamaruzaman Bin Syed Ali*

---

**Submission date:** 03-Jun-2022 02:16PM (UTC+0700)

**Submission ID:** 1849620960

**File name:** 2442-5209-1-PB.pdf (436.02K)

**Word count:** 7096

**Character count:** 39674



[www.ijemst.net](http://www.ijemst.net)

## Nutritional Practices and Body Mass Index among Secondary School Students

45

**Syed Kamaruzaman Bin Syed Ali**

University of Malaya, Malaysia

**Misbon Onn Bin Hassan**

University of Malaya, Malaysia

**Nguang Ung Siong**

University of Malaya, Malaysia

**Imran Akhmad**

Universitas Negeri Medan, Indonesia

50

**Zulakbal Abd Karim**

Sultan Idris Education University, Malaysia

### To cite this article:

Kamaruzaman Bin Syed Ali, S., Onn Bin Hassan, M., Siong, N. U., Akhmad, I., & Abd Karim, Z. (2022). Nutritional practices and body mass index among secondary school students. *International Journal of Education in Mathematics, Science, and Technology (IJEMST)*, 10(3), 618-631. <https://doi.org/10.46328/ijemst.2442>

The International Journal of Education in Mathematics, Science, and Technology (IJEMST) is a peer-reviewed scholarly online journal. This article may be used for research, teaching, and private study purposes. Authors alone are responsible for the contents of their articles. The journal owns the copyright of the articles. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of the research material. All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations regarding the submitted work.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

## Nutritional Practices and Body Mass Index among Secondary School Students

Syed Kamaruzaman Bin Syed Ali, Misbon Onn Bin Hassan, Nguang Ung Siong, Imran Akhmad, Zulakbal Abd Karim

39

### Article Info

#### Article History

Received:

18 September 2021

Accepted:

02 March 2022

#### Keywords

Nutritional practices

Body Mass Index

Physical and health

education

Secondary school students

### Abstract

This study investigated the nutritional practices and body mass index (BMI) among secondary school students in Kerian District, Perak. The respondents of this study consisted of 363 first, second, and fourth-grade students who were selected through a simple random sampling method in secondary schools in Kerian District, Perak. Questionnaires were distributed to selected schools in this study. The questionnaire in the study was obtained from the Meal Pattern Questionnaire (MPQ), which requires the socio-economic status of parents' monthly income, students' body mass index, and students' eating habits. The study's findings showed that the nutritional practices among secondary school students are in the presence of moderate, whereas the body mass index of students is also at the level of underweight. Meanwhile, students' socio-economic status is of the low-income group. In addition, there were significant differences in terms of nutritional practices among secondary school students based on the family's socio-economic status. The study's findings also showed no significant relationship between dietary practices and body mass index (BMI) of secondary school students. The findings of this study can be used as a reference and provide conclusive assistance in improving the nutritional practices through learning sessions and learning of Physical and Health Education in schools.

### Introduction

Physical and Health Education in the Secondary School Standard-Based Curriculum (KSSM) emphasizes healthy eating practices as these components are essential to maintain a healthy lifestyle. Nutritional practices are a component that needs to be studied in the Physical and Health Education syllabus in secondary schools (KPM, 2015; 2016; 2017). In the nutrition practice component, aspects that need to be exposed to students include healthy and safe eating practices, unhealthy eating practices, and the impact of eating habits. The importance of nutritional practices is always emphasized during the teaching and learning sessions of physical and health education because the practice of healthy eating style is an essential topic in Physical and Health Education which is enacted based on the National Education Philosophy in order to produce generations

balanced in terms of physical, emotional, spiritual, intellectual and social. Thus, good nutrition practices among secondary school students in Malaysia need to be given serious attention as the total population of secondary school students in Malaysia is around 2.007 million (MOE, 2019).

Although nutrition practices are focused on teaching and learning sessions in secondary schools, socio-economic developments in the country over the past decade have driven lifestyle changes among Malaysians, especially in terms of nutrition practices that affect nutritional status and body mass index (BMI) across all ages in the Malaysian population for high, medium and low socio-economic groups which include secondary school students. Unhealthy eating habits are among the major contributing factors in the issue of overweight and obesity, and this phenomenon does not only occur among students consisting of children and adolescents but also involves adults (Ismail et al., 2009; IPH, 2008; 2014; 2015).

Obesity in Malaysia has caused various chronic health problems, alarming, either urban or rural. As Malaysia continues to develop rapidly in the economy, the population's health is expected to decline (Ismail et al., 2002). The latest Asian Development Bank Institute report places Malaysia first among the countries in Southeast Asia and ninth in Asia regarding obesity prevalence. It is also revealed that the obesity rate has been increasing dramatically in Malaysia (Helble & Francisco, 2017). In addition, the issue of malnutrition and weight gain also occurs in Malaysia. The issue of obesity and underweight disease is closely intertwined with dietary practices (NCCFN; 2006). Changes in eating habits are associated with changes in health status, and it contributes to an increased risk of chronic diseases like diabetes, cardiovascular disease, hypertension, and cancer. It also negatively impacts the quality of life standards and increases psychosocial problems such as depression, lack of self-confidence, employment discrimination, and other forms of social stigmatization (Harnois & Gabriel, 2002).

### **Statement of Problem**

Healthy eating styles focus on appropriate nutritional practices based on macronutrients and micronutrients as needed to address the issue of obesity and being underweight among secondary school students (KPM, 2015; 2016; 2017). However, nutritional practices among children and adolescents, especially school students in Malaysia, have undergone negative changes. These changes occurred due to the increase in the number of meals consumed at various food premises and an increase in the intake of beverages from milk-based to sweet and sugary drinks apart from the habit of skipping breakfast intake and an increase in snack intake (Ismail et al., 2003). In Malaysia, buying outside food or eating fast food has become a phenomenon among the community (Abdullah, Mokhtar & Bakar, 2015; Tan, 2016). This situation increases obesity among adolescents, especially school students. In addition, unhealthy eating practices have become a national issue due to a lack of awareness about the importance of healthy eating practices in daily life. As a result, it causes an increment in symptoms of obesity among school students in Malaysia because it was found that 90% of school students do not meet the nutritional intake recommended by the Ministry of Health Malaysia (Suhaili, 2007). Meanwhile, the school students also did not practice proper eating habits in their daily diet (Moy, Gan & Siti, 2006), whereas previous studies found out 3 out of 10 students did not practice healthy eating habits (Ismail et al., 2009). Therefore, this

situation has caused Malaysia to face a double burden in coping with nutritional practices such as over-nutrition and malnutrition among the low-income urban population (IPH, 2017).

In addition, Malaysians who do not practice a healthy lifestyle and are not aware of food and nutrition intake are vulnerable to diseases such as hypertension or high blood pressure (IPH, 2014; 2015). Only a slight 11.2 percent of students practice healthy eating style even 73.8 percent of school students know nutritional practices. Thus, unhealthy eating habits coupled with a worsening obesity crisis have been observed in developing countries (Goh, Ali, McCullough & Mitra, 2021). An Asian Development Bank Institute study also concluded that Malaysia has the most obese population in Southeast Asia (Helble & Francisco, 2017). This increase in the percentage of obesity is related to unhealthy eating habits that will lead to chronic diseases, including heart attack and stroke being the significant diseases causing death, commonly occur among adolescents which includes secondary school students and also adults (Abdullah et al., 2015; Tan, 2016). Thus, dietary practices are factors contributing to the obesity problem among Malaysians, especially school students.

Similarly, The National Health and Morbidity Survey (IPH, 2017) also found that school students did not practice healthy eating habits. Body Mass Index (BMI) data from the Assessment of Physical Activity, Sports and Co-curriculum (PAJSK) has shown that Year Six students in the category at risk of being overweight also increased to 75,171 people (17.3 percent), compared to 56,584 people (13.4 percent) in 2017 (Jafar, & Nasbah, 2018). The increase in the BMI data of the students is very upsetting and needs to be given extreme attention by all parties because obesity can be particularly detrimental to be the cause of various health problems now and in the future. BMI data also showed that a total of 32,142 (7.4 percent) school students were found to have lost weight in the BMI test (Jafar & Nasbah, 2018).

Furthermore, socio-economic factors also contribute to the issue of dietary practices in Malaysia (Shahar et al., 2019). Body mass index (BMI) changes according to age and dietary practices for children and adolescents, especially secondary school students. Being underweight is also related to nutritional practices, especially among school students who live in poverty and come from low-income socio-economic status families (Shahar et al., 2019; Alam et al., 2016). In 2016, an estimated 340 million children and adolescents, including school students, were overweight and obese (Abarca-Gómez et al., 2017). This trend has increased drastically from 4% in 1975 to 18% in 2016. In addition, an estimated 192 million school children and adolescents were also underweight in 2016 (Abarca-Gómez et al., 2017). The World Health Organization (WHO) also expressed concern over this situation, and systematic steps need to be taken to address the issue (Helble & Francisco, 2017). Therefore, teenagers in Malaysia, especially secondary school students, are not excluded from this issue. Meanwhile, weight and dietary practices have been shown to have a very significant correlation (Fogelholm & Kukkonene-Harjula, 2000). Therefore, the Malaysian government has launched the National Nutrition Action Plan III from 2016 to 2025 to promote healthier eating habits (MHM, 2016).

### **Research Questions**

This study aims to provide answers to the following research questions:

1. What are the nutritional practices among secondary school students?
2. What is the body mass index (BMI) among secondary school students?
3. Are there significant differences in nutritional practices among secondary school students based on family socio-economic status?
4. Is there a significant relationship between dietary practices and the body mass index (BMI) of secondary school students?

## Literature Review

### Nutritional Practices and Socio-economic Status

Mukhari and Yasin (2010) have studied nutritional practices among secondary school students living in Johor in a survey. The study's findings reported that a total of 88 respondents, male and female students practiced a regular intake of breakfast, lunch, and dinner every day. Most of the respondents consume foods that contain various nutrients needed by the body according to the correct meal time every day. The study also found that those who are knowledgeable and have higher socio-economic status have a more positive attitude and adopt better eating habits. Similarly, Abdullah & Ali (2011) has also conducted a study to review nutritional practices among university students and their perceptions of proper nutritional practices. This study involved 102 respondents consisting of undergraduate students from Year 1 to Year 3 of various faculties in the main campus of Universiti Kebangsaan Malaysia, Bangi. The survey results concluded that most respondents have sufficient knowledge and awareness of the aspects of proper and healthy nutrition. However, this study did not consider the socio-economic factors of the students' families.

Furthermore, Karim et al. (2014) also conducted a study to determine the association of socio-demographic factors with eating habits among preschool children in Peninsular Malaysia. A total of 1,933 preschool children aged 4–6 years participated in the study. Parents or guardians were interviewed based on their children's socio-demographic characteristics and eating habits. Height and weight of preschool children were measured; BMI for age, weight for age, and height for age were also determined. The study found that the average monthly household income is RM3,610, with 59.6% of the parents having secondary school-level education. Most preschoolers also consumed breakfast, lunch, and dinner daily, with the number of children skipping main meals at only 15.0%. The level of parental education and household income are fundamental about the intake of fruits, vegetables, milk and dairy products, and fast food. However, there was no significant relationship between children's weight status and frequency of eating staple foods, fruits, vegetables, milk, dairy products, and fast foods.

The study of Sharif et al. (2016) differs from previous studies that investigated energy intake, nutrients, and category of food intake involving 749 urban children (1-10 years old) based on household income status. Children's nutritional intake was obtained using "food recall" and recorded for two days. Adequate diet taken in this study is assessed based on recommended energy and nutrient intake and meals by food category. The study found that children from low-income groups have a low level of energy intake, and most of them do not meet the recommended energy and nutrient intake compared to children from middle and high-income groups. Thus,

these findings suggested that low socio-economic status, such as low household income, may limit access to adequate diets, especially for teenagers. However, this study does not consider the factors of dietary practice as a variable.

### Nutritional Practices and Body Mass Index (BMI)

A survey study by Chong et al. (2016) was conducted to investigate children's nutrition practices through a questionnaire using South East Asian Nutrition Surveys (SEANUTS) in Malaysia. It has identified that out of 2797 children aged 2 to 12 years, only 56.1% of children consume three main meals every day. About 20% of children consume snacks three times a day, while 9.7% eat fast food weekly. This irregular eating pattern is associated with low micronutrient intake, and it happens more frequently among Malay teenagers and those living in rural areas. However, the study did not consider the factors of weight and body mass index (BMI). Meanwhile, a study by Umairah et al. (2012) involving 204 students from Selangor, aged 7 to 10 years, was conducted to determine the relationship between dietary patterns and body mass index (BMI) among primary school children. The results of this cross-sectional study showed a significant relationship between the type of diet and body mass index.

Moreover, the cross-sectional study was also conducted by Howe, Black, Wong, Parnell, and Skidmore (2013) to examine the relationship between nutritional practices and adolescent body composition. Information on food intake and current nutritional status was collected using a web-based survey involving 681 adolescents attending schools in Otago, New Zealand. The study found a significant relationship between nutritional practices and overall adiposity but no significant relationship with body mass index (BMI). These findings are more significant for male adolescents than female adolescents. A study by Law, Nasir, and Hazizi (2013) was conducted on the factors related to breakfast neglect among adolescents in Sarawak. This cross-sectional study was conducted to determine differences in weight status, socio-demographic, behavioral, and psychological characteristics between daily breakfast takers and breakfast neglecters. Data were collected from 275 forms from four secondary schools students. Weight and height were measured using standard procedures to determine the questionnaire's body mass index (BMI) scores. The study's findings showed a significant relationship between breakfast neglect and the body mass index (BMI) status of secondary school students.

In addition, the Family Diet Study by Yang (2017) involved a total of 236 Malay students in five national primary schools in Malaysia. The study aimed to determine the relationship between food intake and body mass index (BMI) among children aged 8 to 12 years in Malaysia. It was found that there was a moderately positive correlation between food intake and weight. Thus, the body mass index of primary school children showed a significant relationship with the type of food intake but did not significantly correlate with breakfast intake and total food intake in a day. However, this study only focused on the population among primary school students.

### Method

This study uses a survey study design to obtain necessary information about nutritional practices and body mass

index among secondary school students through a questionnaire. This survey method is also used to collect information in the form of comparative data (quantifiable information), a set of similar questions given to a large sample of studies in this study. Therefore, this survey method is highly appropriate (Cresswell & Garret, 2008; Roberts, Spink, & Pemberton, 1999). Furthermore, the survey method is very suitable to measure opinions, achievements, and attitudes of respondents towards what is happening, and it is also suitable for collecting information on independent and dependent variables (Kerlinger, 1973; Konting, 2005).

The population in this study consists of secondary school students in rural areas in the district of Kerian, Perak. The sample in this study was secondary school students between 13, 14, and 16 years old. The population of this study involves 6628 students, which consists of 3300 male students and 3328 female students, respectively. The total sample selected is 363 students from 7 secondary schools from Perak State Education Department in the Kerian district. The simple random sampling technique was used in this study by dividing 15 secondary schools in Kerian district, Perak according to the zone and later selecting seven schools from the entire district independently. The total sample was selected using the method Krejcie and Morgan (1970) introduced. From the population of 6628 students, a total of 363 secondary school students were selected by simple random sampling in this study which consisted of 189 (52%) male students and 175 (48%) female students from 13, 14, and 16 years old from families with different socio-economic background such as high, medium and low status. Simple random sampling will give equal opportunity to all respondents to be selected. The study's findings will be able to be generalized to a population (Konting, 2005).

#### **Validity and Reliability**

The questionnaire is the measurement instrument chosen in carrying out this study. This instrument contains three sections, A, B, and C. Section A is a demographic instrument containing information about the demographics of the respondents, such as the socio-economic status of the parents' monthly income. Part B is an instrument that contains questions related to the status of body mass index (BMI) and information on the weight and height of students (WHO/IASO/IOTF, 2000). Section C contains questions about students' nutritional practices obtained from the Meal Pattern Questionnaire (MPQ) (Alfonsson et al., 2015). The reliability value of the MPQ instrument was between 0.63-0.89 for individuals (Alfonsson et al., 2015). The questionnaire was referred to five experts specializing in language, content, and field of physical education and health to review the validity of the questionnaire's content and the consistency of translation of the questionnaire. Suggested amendments and improvements from the feedback were made to the instrument from the experts consulted to ensure its validity.

In a pilot study, researchers have identified 30 respondents consisting of secondary school students from the state of Perak. A total of 10 respondents of form 1, 2, and 4 students were randomly selected in the pilot study. Respondents were given two days to answer the research instrument. The instrument's reliability was tested through Cronbach's Alpha coefficient, and if the value of the reliability coefficient is higher than 0.60m, it is indicated that the instrument is suitable for use in real studies (Konting, 2005). The overall Cronbach's value reported for the nutritional practice items in the pilot study is 0.668. This result suggests that each item of



nutritional practice among secondary school students can be maintained and is suitable for the actual study.

## Results

### Question 1: What are the nutritional practices among secondary school students?

As shown in Table 1, the highest mean score for the nutritional practice among secondary school students is lunch practice (M = 3.79, SD = 1.31). The mean score for the practice of taking snacks after breakfast (M = 1.88, SD = 0.99) was the lowest. So, the secondary school students are more focused on lunch intake practices than other dietary practices. The value of the overall mean score for nutritional practices among secondary school students is at a moderate level (M = 28.0, SD = 6.63).

Table 1. Nutritional Practices of Secondary School Students

No	In the last seven days, how many times have you taken	Mean Score	SD
...			
1	Breakfast	2.91	1.44
2	Snacks after breakfast	1.88	0.99
3	Lunch	3.79	1.31
4	Snacks after lunch	2.19	1.44
5	Dinner	3.49	1.39
6	Snacks after dinner	2.19	1.20
7	Meals during school breaks	3.24	1.32
8	Snacks	2.26	1.02
9	Fruits	3.14	1.18
10	Vegetables	2.95	1.39
	Nutritional Practices	28.0	6.63

### Question 2: What is secondary school students' body mass index (BMI)?

Based on Table 2, the study's findings showed that out of 363 respondents involved, a total of 44.1% were underweight while 33.6% were of normal weight. Furthermore, the findings show that 15.4% are in this category for overweight. In addition, only 6.9% of respondents are in the obesity category. These findings explain that 44.1% of respondents among secondary school students have underweight symptoms.

Table 2. Body Mass Index (BMI) Among Secondary School Students

Mass Index	Frequency	Percentage
Less weight	160	44.1%
Normal (ideal weight)	122	33.6%
Overweight	56	15.4%
Obesity	25	6.9%
Overall	363	100.0%

**Question 3: Are there significant differences in nutritional practices among secondary school students based on family socio-economic status?**

The Kruskal Wallis test showed that there were significant differences in nutritional practices among secondary school students based on the socio-economic status of the family with values of  $\chi^2 (2) = 11.748$  and sig = 0.003 ( $p < 0.05$ ) as shown in Table 3. In terms of mean ranking, students with socio-economic status having an income of RM5000 and above (MR = 224.12) recorded higher nutritional practices than students with socio-economic status having income between RM2501 to RM4999 (MR = 199.41) and students with socio-economic status, having an income of RM2500 and below (MR = 167.14). Since the Kruskal Wallis test showed significant differences, the Mann Whitney U test was conducted to look in detail at the differences for each group of students according to socio-economic status. The results of the Mann Whitney U test are in Table 3.

**Table 3.** Kruskal Wallis Test of Differences in Nutritional Practices among Secondary School Students based on Family Socio-economic Status

Income	N	Mean Ranking	$\chi^2$	Df	Sig.
RM2500 and below	215	167.14	11.748	2	0.003
RM2501-RM4999	123	199.41			
RM5000 and above	25	224.12			

Based on Table 4, Mann Whitney U test showed that there was a significant difference in the nutritional practices of secondary school students between students with family socio-economic status having a parental income of RM2500 and below and students with family socio-economic status having parental income between RM2501 to RM4999 with U value = 10872.000 and sig = 0.006 ( $p < 0.05$ ). In terms of mean ranking, it is shown that students with family socio-economic status having parental income between RM2501 to RM4999 (MR = 188.61) had higher nutritional practices compared to students with family socio-economic status having a parental income of RM2500 and below (MR = 158.57). Furthermore, there is a significant difference in the nutritional practices of secondary school students between students with family socio-economic status having a parental income of RM2500 and below and students with family socio-economic status having a parental income of RM5000 and above with values of U = 1844.00 and sig = 0.010 ( $p < 0.05$ ).

**Table 4.** Mann Whitney U Test of Differences in Nutritional Practices among Secondary School Students based on Family Socio-economic Status

Parental Income	N	MR	SR	U	Sig.
RM2500 and below	215	158.57	34092.00	10872.000	0.006
RM2501-RM4999	123	188.61	23199.00		
RM 2500 and below	215	116.58	25064.00	1844.000	0.010
RM5000 and above	25	154.24	3856.00		
RM2501-RM4999	123	72.80	8954.00	1328.000	0.283
RM5000 and above	25	82.88	2072.00		

In terms of mean ranking, students with family socio-economic status having a parental income of RM5000 and above (MR = 154.24) have higher nutritional practices than students with family socio-economic status with a parental income of RM2500 and below (MR = 116.58), as shown in Table 4. While the results of the Mann Whitney U test showed that there was no significant difference in the nutritional practices of secondary school students between students with family socio-economic status having parental income between RM2501 to RM4999 and students with family socio-economic status having a parental income of RM5000 and above with U = 1328.00 and sig = 0.283 (p > 0.05). In terms of mean ranking, it is shown that students with family socio-economic status having a parental income of RM5000 and above (MR = 82.88) had dietary practices similar to students with family socio-economic status having parental income between RM2501 to RM4999 (MR = 72.80).

**Question 4: Is there a significant relationship between nutritional practices and secondary school students' body mass index (BMI)?**

In this study, the Spearman correlation was used to examine the relationship between nutritional practices and secondary school students' body mass index (BMI). Spearman correlation test is used because there is one variable in the study that uses an ordinal scale: body mass index (BMI) (Pallant, 2007). Table 5 shows the correlation coefficient value between the mean score of Nutritional Practices and Body Mass Index (BMI) among secondary school students. The findings showed that there was no significant relationship between Nutritional Practices and Body Mass Index (BMI) with a value of r= 0.042, sig = 0.421 (p> 0.05). The relationship between the two variables was insignificant. This result indicates that their nutritional practices do not influence secondary school students' Body Mass Index (BMI).

Table 5. Relationship between Nutritional Practices and Body Mass Index (BMI) Among Secondary School Students

Relationship	Body Mass Index (BMI)		Interpretation
	r	Sig.	
Nutritional Practices	0.042	0.421	-

**Discussion**

Socio-economic status is one of the factors that is often emphasized in influencing a student's eating habits. Students' nutritional practices involving food such as breakfast have become a widespread nationwide issue discussed among various parties and levels of society, including the Ministry of Education Malaysia. This issue has resulted in a proposal to provide complimentary breakfast to all primary school students in the B40 group starting 2020. It is in line with the government's extension plan from the Supplementary Meal Plan program, which has long been implemented by the Ministry of Education Malaysia to help the low-income groups, B40, reduce the cost of living in urban and rural areas.

Students with high socio-economic status are recognized to have better nutritional practices than students with

low and medium socio-economic status. In the context of this study, it is proven that the practice of nutrition as a personal factor is significantly different due to the influence of the environment and the socio-economic status of students' families. This situation suggests that socio-economic status factors can affect a person's eating habits, whether good, moderate, or wrong. Students' eating practices through the rate of food intake, breakfast, fruits, and vegetables were influenced by family socio-economic status (Yannakoulia et al., 2015). Indeed, the findings of this study are in line with statements from Skårdal, Western, Ask, and Overby (2014), studies in Norway, where there are significant differences in nutritional practices among secondary school students based on family socio-economic status. They found that either low or high family socio-economic status can have positive or negative influences on a student's eating habits.

In addition, it was found that students from low socio-economic status families are experiencing food shortages while students from high socio-economic status families obtain more nutritious food with their eating habits and have better-eating habits (Mukhari & Yasin, 2010). Thus, students from the socio-economic status group of high-income families are more likely to suffer from obesity than other socio-economic status groups of families (Soo et al., 2011). This situation is likely to cause an increase in the number of students who are underweight and obese either in urban or rural areas. However, the findings of this study were contrary to research conducted by Karim et al. (2014), which concluded that the socio-economic status of the family does not influence the nutritional practices of the students when their findings indicated that there were no significant differences in dietary practices among students based on socio-economic status factors.

Based on the findings of this study, it was also found that there is no significant relationship between nutritional practices and body mass index (BMI) among secondary school students in the district of Kerian, Perak. It clearly shows that their dietary practices do not influence secondary school students' body mass index (BMI). The results of this study are different from previous studies that found that there is a significant relationship between dietary practices and body mass index (BMI). This study contrasts with Umairah et al. (2012), who found that body mass index (BMI) had a significant association with the types of diet among school students. This finding is further supported by Law et al.' (2013) study, which concluded that students who regularly practice breakfast intake are regular weight students compared to breakfast neglecters because students who were overweight and rarely ate breakfast but consumed other meals in large quantities such as lunch and dinner compared to average, overweight students (Ha et al., 2016). However, low intake of fruits and vegetables is also associated with weight gain and BMI, which is at risk of obesity (Brunst, Rhee, & Zhong, 2008).

However, the results of this study are similar to the findings of the study by Howe et al. (2013), who also found that there was no significant relationship between nutritional practices and body mass index (BMI) because differences in students' eating habits, in which students who did not eat breakfast and were not experiencing weight gain compared to students who regularly ate breakfast (Berkey et al., 2003). Thus, breakfast intake and the amount of food intake per day did not affect the status of body mass index (BMI). This finding is also in line with the study of Yang (2017), who also found that there was no significant relationship between total food intake in a day and breakfast intake with body mass index (BMI) among students. Therefore, neglecting breakfast, consuming fewer fruits, fewer drinks, and frequent snacks intake do not affect a student's body mass

index (BMI) (Al-Muammar, El-Shafie & Feroze, 2014). The insignificant relationship between nutritional practices and body mass index (BMI) among secondary school students in this study is likely to occur due to genetic or hereditary factors. Despite eating a lot, likely, a person will not be obese due to the genetic factor of being thin. In addition, there is also the possibility that the attitude of a person who likes to skip meals causes the risk of becoming obese or obesity to be low due to fewer calories burned. This situation shows that the level of body mass index (BMI) of secondary school students in Kerian district, Perak has no relationship with the dietary practices.

## Conclusion

Overall, it can be concluded that the family's socio-economic status - in the context of this study refers to family income- is an essential factor in determining students' nutritional practices in their daily lives. In the context of Physical and Health Education, teaching and learning sessions are needed to increase students' understanding of the importance of healthy eating practices every day, especially the intake of breakfast, eating during school breaks, lunch, dinner, vegetables, fruits, and implications of skipping meals and fast food intake towards students' health status. This study has provided meaningful ideas and contributions, and it can be used as a guide for other individuals, leading to optimal improvements in consumers' nutritional practices. The findings of this study can be used as a reference and provide a meaningful contribution, especially in improving the quality of teaching and learning of Physical and Health Education in rural secondary schools so that students' healthy lifestyle practices can be improved. Therefore, it is recommended that future researchers will explore more profound in this field of study in order to produce more dynamic and conclusive researches that will help to contribute to the teaching and learning process of physical and health education and to review nutritional practices and the body mass index (BMI) as well as its relationship with the socio-economic status of students. This suggestion is essential in strengthening the physical and health education syllabus, which can also be integrated with other subjects such as Science and Sports Science.

## Acknowledgments

I would like to thank all the leaders of the Faculty of Education, University of Malaya, and all colleagues who have supported this research.

## References

- Abarca-Gómez, L., Abdeen, Z. A., Hamid, Z. A., Abu-Rmeileh, N. M., Acosta-Cazares, B., Acuin, C., Aguilar-Salinas, C. A. (2017). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *The Lancet*, 390(10113), 2627-2642.
- Abdullah, M. A., & Ali, N. (2011) Amalan pemakanan dalam kalangan pelajar universiti dan implikasinya terhadap pembelajaran. *Jurnal Personalia Pelajar*, 14, 59-68.
- Abdullah, N. N., Mokhtar, M. M., Bakar, M. H. A. (2015). Trend on Fast Food Consumption in Relation to

- Obesity among Selangor Urban Community. *Procedia-Social and Behavioral Sciences*, 202, 505-513.
- Alam, M. M., Siwar, C., Wahid, A. N. M., Talib, B. A. (2016). Food security and low-income households in the Malaysian east coast economic region: an empirical analysis. *Review of Urban and Regional Development Studies*, 28, 1-15.
- Alfonsson, S. A., Sewall, H., Lidholm, T., & Hursti. (2015). The Meal Pattern Questionnaire: A psychometric evaluation using the Eating Disorder Examination. *Eating Behaviors*, 21, 7-10.
- Al-Muammar, M. N., El-Shafie, M., & Feroze, S. (2014). Association between dietary habits and body mass index of adolescent females in intermediate schools in Riyadh, Saudi Arabia. *Eastern Mediterranean Health*, 20(1), 39-45.
- Berkey, C. S., Rockett, H. R. H., Gillman, M. W., Field, A. E., & Colditz, G. A. (2003). Longitudinal study of skipping breakfast and weight change in adolescents. *International Journal of Obesity*, 27(10), 1258-1266.
- Brunt, A., Rhee, Y. & Zhong, L. M. S. (2008). Differences in Dietary Patterns Among College Students According to Body Mass Index. *Journal of American College Health*, 56(6), 629-634.
- Chong, K. H., Wu, S. K., Hafizah, Y. N., E., M. C., & Poh, B. K. (2016). Eating Habits of Malaysian Children: Findings of the South East Asian Nutrition Surveys (SEANUTS). *Asia Pacific Journal of Public Health*, 28(5), 59-73.
- Fogelholm, M., & Kukkonen-Harjula, K. (2000). Does physical activity prevent weight gain? A systematic review. *Obesity Review*, 1, 95-111.
- Goh, E. V., Ali, S. A., McCullough, F., & Mitra, S. R. (2021). The nutrition transition in Malaysia: Key drivers and recommendations for improved health outcomes. *BMC Nutrition*, 6(32), 1-14.
- Ha, S. A., Lee, S. Y., Kim, K. A., Seo, J. S., Sohn, C. M., Park, H. R., & Kim, K. W. (2016). Eating habits, physical activity, nutrition knowledge, and self-efficacy by obesity status in upper-grade elementary school students. *Nutrition Research and Practice*, 10(6), 597-605.
- Harnois, G., & Gabriel, P. (2002). *Mental health and work: Impact, issues and good practices*. Geneva: World Health Organization.
- Helble, M., & Francisco, K. (2017). *The Imminent Obesity Crisis In Asia And The Pacific: First Cost Estimates*. Tokyo: Asian Development Bank Institute.
- Howe, A. S., Black, K. E., Wong, J. E., Parnell, W. R., & Skidmore, P. M. (2013). Dieting status influences associations between dietary patterns and body composition in adolescents: A cross-sectional study. *Nutrition Journal*, 12(1), 1-10.
- Institute for Public Health (IPH). (2008). *National Health and Morbidity Survey 2006 (NHMS III): Nutritional Status*. Putrajaya: Ministry of Health, Malaysia.
- Institute for Public Health (IPH). (2014). *National Health and Morbidity Survey 2014: Malaysian Adult Nutrition Survey (Volume II)*. Putrajaya: Ministry of Health, Malaysia.
- Institute for Public Health (IPH). (2015). *National Health and Morbidity Survey 2015: Non-Communicable Diseases, Risk Factors & Other Health Problems (Volume II)*. Putrajaya: Ministry of Health, Malaysia.
- Institute for Public Health (IPH). (2017). *National Health and Morbidity Survey (NHMS) 2017: Adolescent Nutrition Survey*. Putrajaya: Ministry of Health, Malaysia., Malaysia.

- Ismail M. N., Chee, S.S., Nawawi, H., Yusoff, K., Lim, T.O., & James, W.P.T. (2002). Obesity in Malaysia. *The International Association for the Study of Obesity. Obesity Reviews*, 3, 203-208
- Ismail, M. N., Norimah, A. K., Poh, B. K., Ruzita, A. T., Mazlan, N. M., Shanita, N. S., Roslee, R. & Zakiah, N, M. S. (2009). *Nutritional status and dietary habits of primary school children in Peninsular Malaysia (2001-2002)*. Department of Nutrition and Dietetics, Faculty of Allied Health Sciences, Universiti Kebangsaan Malaysia.
- Jafar, F. & Nasbah, N, N. (2018). Tahap kanak-kanak 12 tahun meningkat. *Berita Harian Online*. <https://www.bharian.com.my/berita/pendidikan/2018/11/503397/tahap-obesiti-kanak-kanak-12-tahun-meningkat>
- Karim A., N., Mohd Nasir, M. T., Hazizi, A. S., Suraya, P., Loh, S. H., & Nurliyana, A. R. (2014). Association of body weight status and socio-demographic factors with food habits among preschool children in Peninsular Malaysia. *Malaysian Journal of Nutrition*, 20(3), 303-315.
- Kementerian Pendidikan Malaysia (KPM). (2015). *Dokumen standard kurikulum dan pentaksiran pendidikan jasmani dan kesihatan tingkatan satu*. Putrajaya: Bahagian Perkembangan Kurikulum.
- Kementerian Pendidikan Malaysia (KPM). (2016). *Dokumen standard kurikulum dan pentaksiran pendidikan jasmani dan Kesihatan tingkatan dua*. Putrajaya: Bahagian Perkembangan Kurikulum.
- Kementerian Pendidikan Malaysia (KPM). (2017). *Dokumen standard kurikulum dan pentaksiran pendidikan jasmani dan kesihatan tingkatan tiga*. Putrajaya: Bahagian Perkembangan Kurikulum.
- Kerlinger, F.N. (1973). *Foundation of Behavioural Research*. New York. Holt. Rinehand and Hinston.
- Krejcie, R.V., & Morgan, D.W. (1970). Determining Sample Size for Research Activities. *Educational and Psychological Measurement*, 30, 607-610
- Konting, M. M. (2005). *Kaedah penyelidikan pendidikan*. Kuala Lumpur: Dewan Bahasa dan Pustaka.
- Law, L. S., Nasir, M. T. & Hazizi, A. S. (2013) Factors Associated with Breakfast Skipping among School-going Adolescents in Sarawak, Malaysia. *Malaysian Journal of Nutrition*, 19(3), 401-407.
- Ministry of Education Malaysia (MOE). (2019). *Quik Fact 2019: Malaysia Educational Statistics*. Putrajaya: Educational Planning and Research Division.
- Ministry of Health Malaysia (MHM) (2016). *National Plan of Action for Nutrition of Malaysia III 2016-2025*. Kuala Lumpur: National Coordinating Committee on Food and Nutrition (NCCFN).
- Moy, F. M., Gan, C.Y., Siti, Z. M. K. (2006). Eating patterns of school children and adolescents in Kuala Lumpur. *Malaysian Journal of Nutrition*, 12, 1-10.
- Mukhari, A. W. B., & Yasin, I. A. B. M (2010) Amalan Pemakanan Di Kalangan Remaja Di Kawasan Felda Bukit Ramun, Kulai, Johor. *Universiti Teknologi Malaysia Institutional Repository*, 1-9.
- National Coordinating Committee on Food and Nutrition (NCCFN). (2006). *National Plan of Action for Nutrition of Malaysia*. Putrajaya: Ministry of Health, Malaysia.
- Roberts, G. C., Spink, K.S., & Pemberton, C. L. (1999). *Learning Experiences in Sport Psychology (2nd Ed.)* Champaign, IL: Human Kinetics.
- Shahar, S., Lau, H, Puteh, S. E.W., Amara, S., & Razak, N. A. (2019). Health, access and nutritional issues among low-income population in Malaysia: Introductory note. *BMC Public Health*, 19(4), 1-5
- Sharif, R., Chong, K. H, Zakaria, N. H., Ong, M. L., Reilly, J. J., Wong, J. E., Saad, H. A., & Poh, B. K. (2016). Results From Malaysia's 2016 Report Card on Physical Activity for Children and Adolescents. *Journal*


- of Physical Activity and Health*, 13(11), 201-205.
- Skårdal, M., Western, I. M., Ask, A. M. S., & Øverby, N. C. (2014). Socio-economic differences in selected dietary habits among Norwegian 13–14 year-olds: a cross-sectional study. *Food & Nutrition Research*, 58(1), 1-8.
- Soo, K. L., Wan, A. M., Abdul, M. H., & Lee, Y. Y. (2011). Dietary practices among overweight and obese Chinese children in Kota Bharu, Kelantan. *Malaysian Journal of Nutrition*, 17(1), 87-95.
- Suhaili S (2007). *Corak pengambilan buah-buahan dan sayur-sayuran di kalangan kanak-kanak di Sekolah Kebangsaan Serdang. (Tesis Sarjana)*. Kuala Lumpur: Universiti Kebangsaan Malaysia.
- Tan L. Y. (2016) *Fast Food Consumption Behaviour Among Generation Behaviour among generation Y in Malaysia. (Master Thesis)*. Selangor: Universiti Tunku Abdul Rahman
- Umairah, S. N., Yahya, B. T., Datin, M., & Yusof, S. M. (2012). Relationship between dietary pattern and body mass index among primary school children. *Asian Journal of Clinical Nutrition*, 4(4), 142-150.
- WHO/IOTF/IASO. (2000). *The Asia-Pacific perspective: Redefining Obesity and its Treatment*. Hong Kong: World Health Organization, International Obesity Task Force, International Association for the Study of Obesity.
- Yang, W. Y., Burrows, T., MacDonald-Wicks, L., Williams, L. T., Collins, C. E., Chee, W. S., & Colyvas, K. (2017). Body Weight Status and Dietary Intakes of Urban Malay Primary School Children: Evidence from the Family Diet Study. *Children*, 4(1), 1-16.
- Yannakoulia, M., Lykou, A., Kastorini, C. M., Saranti Papasaranti, E., Petralias, A., & Veloudaki, A. (2015). Socio-economic and lifestyle parameters associated with diet quality of children and adolescents using classification and regression tree analysis: the DIATROFI study. *Public Health Nutrition*, 19(2), 339–347.

---


#### Author Information

---


**Syed Kamaruzaman Bin Syed Ali**

 <https://orcid.org/0000-0002-4804-1237>  
Faculty of Education  
University of Malaya  
Malaysia


**Misbon Onn Bin Hassan**

 <https://orcid.org/0000-0003-2685-7523>  
Faculty of Education  
University of Malaya  
Malaysia  
Contact e-mail: [onn2017@siswa.um.edu.my](mailto:onn2017@siswa.um.edu.my)


**Nguang Ung Siong**

 <https://orcid.org/0000-0002-5891-4791>  
Faculty of Education,  
University of Malaya  
Malaysia

**Imran Akhmad**

 <https://orcid.org/0000-0001-5193-8005>  
Faculty of Sport Science  
Universitas Negeri Medan  
Indonesia

**Zulakbal Abd Karim**

 <https://orcid.org/0000-0001-9883-5767>  
Faculty of Sports Science and Coaching  
Sultan Idris Education University  
Malaysia

---



# Nutritional Practices and Body Mass Index Among Secondary School Students

## ORIGINALITY REPORT

**25%**  
SIMILARITY INDEX

**21%**  
INTERNET SOURCES

**13%**  
PUBLICATIONS

**6%**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [pesquisa.bvsalud.org](https://pesquisa.bvsalud.org) **3%**  
Internet Source

**2** Submitted to Universitas Negeri Semarang **2%**  
Student Paper

**3** [medcraveonline.com](https://medcraveonline.com) **2%**  
Internet Source

**4** [umexpert.um.edu.my](https://umexpert.um.edu.my) **2%**  
Internet Source

**5** [worldwidescience.org](https://worldwidescience.org) **1%**  
Internet Source

**6** Submitted to Universitas Negeri Medan **1%**  
Student Paper

**7** Syed Kamaruzaman Syed Ali, Misbon Onn Hassan, Zulakbal Abd Karim. "The importance of eating habits and physical activities among school students", International Physical Medicine & Rehabilitation Journal, 2019 **1%**  
Publication

8	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1 %
9	Shariff, Zalilah Mohd, Khor Geok Lin, Sarina Sariman, Huang Soo Lee, Chin Yit Siew, Barakatun Nisak Mohd Yusof, Chan Yoke Mun, and Maznorila Mohamad. "The relationship between household income and dietary intakes of 1-10 year old urban Malaysian", Nutrition Research and Practice, 2015. Publication	1 %
10	<a href="http://psasir.upm.edu.my">psasir.upm.edu.my</a> Internet Source	1 %
11	Submitted to Universiti Kebangsaan Malaysia Student Paper	1 %
12	<a href="http://nutrition.moh.gov.my">nutrition.moh.gov.my</a> Internet Source	1 %
13	<a href="http://scialert.net">scialert.net</a> Internet Source	<1 %
14	<a href="http://researchspace.ukzn.ac.za">researchspace.ukzn.ac.za</a> Internet Source	<1 %
15	<a href="http://jyx.jyu.fi">jyx.jyu.fi</a> Internet Source	<1 %
16	<a href="http://www.iosrjournals.org">www.iosrjournals.org</a> Internet Source	<1 %

17 Thomas, Liz, Quinn, Jocey. "First Generation Entry into Higher Education", First Generation Entry into Higher Education, 2006  
Publication <1 %

---

18 [iku.moh.gov.my](http://iku.moh.gov.my)  
Internet Source <1 %

---

19 [www.oapub.org](http://www.oapub.org)  
Internet Source <1 %

---

20 [studentsrepo.um.edu.my](http://studentsrepo.um.edu.my)  
Internet Source <1 %

---

21 Qaiser Suleman, Ishtiaq Hussain, Zaib-un-Nisa. "Effects of Parental Socioeconomic Status on the Academic Achievement of Secondary School Students in District Karak (Pakistan)", International Journal of Human Resource Studies, 2014  
Publication <1 %

---

22 Ali Saad, Hadeel Hamady, Tanya Kronos, Ludmila Sandler, Dorit Itah. "Trends in Psychiatric Admissions of Children and Adolescents During Early Corona Period Year (2020) Compared to Pre-Corona Year (2018): A Cross Sectional Study in Israel", Research Square Platform LLC, 2022  
Publication <1 %

---

23 [dspace.plymouth.ac.uk](http://dspace.plymouth.ac.uk)  
Internet Source <1 %

---

24	<a href="http://actacommercii.co.za">actacommercii.co.za</a> Internet Source	<1 %
25	<a href="http://core.ac.uk">core.ac.uk</a> Internet Source	<1 %
26	Kar Hau Chong, Suet Kei Wu, Yatiman Noor Hafizah, Marjolijn C. E. Bragt, Bee Koon Poh. "Eating Habits of Malaysian Children", Asia Pacific Journal of Public Health, 2016 Publication	<1 %
27	Richard J. Deckelbaum, Christine L. Williams. "Childhood Obesity: The Health Issue", Obesity Research, 2001 Publication	<1 %
28	<a href="http://digitalcommons.andrews.edu">digitalcommons.andrews.edu</a> Internet Source	<1 %
29	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1 %
30	<a href="http://www.tci-thaijo.org">www.tci-thaijo.org</a> Internet Source	<1 %
31	Submitted to Universiti Teknologi MARA Student Paper	<1 %
32	<a href="http://irep.iium.edu.my">irep.iium.edu.my</a> Internet Source	<1 %
33	<a href="http://nutriweb.org.my">nutriweb.org.my</a> Internet Source	<1 %

34

[sosyaldergi.usak.edu.tr](http://sosyaldergi.usak.edu.tr)

Internet Source

&lt;1 %

35

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

&lt;1 %

36

Silambarasi Kuralneethi, Sarina Sariman, Vaidehi Ulaganathan. "Gender and age differences in the relationship between calorie, macronutrients intake and growth status of school-aged Aboriginal children at Labu, Negeri Sembilan", British Food Journal, 2020

Publication

&lt;1 %

37

Zahra Gorji, Hamed Kord Varkaneh, Sam talaee, Ali Nazary-Vannani et al. "The effect of green-coffee extract supplementation on obesity: A systematic review and dose-response meta-analysis of randomized controlled trials", Phytomedicine, 2019

Publication

&lt;1 %

38

[discovery.ucl.ac.uk](http://discovery.ucl.ac.uk)

Internet Source

&lt;1 %

39

[ijte.net](http://ijte.net)

Internet Source

&lt;1 %

40

[vuir.vu.edu.au](http://vuir.vu.edu.au)

Internet Source

&lt;1 %

41

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

42

Submitted to Pennsylvania State System of  
Higher Education

Student Paper

<1 %

43

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

44

[pt.scribd.com](http://pt.scribd.com)

Internet Source

<1 %

45

[www.e-iji.net](http://www.e-iji.net)

Internet Source

<1 %

46

S. E. H. Tung, X. H. Ng, Y. S. Chin, M. N. Mohd  
Taib. "Associations between parents'  
perception of neighbourhood environments  
and safety with physical activity of primary  
school children in Klang, Selangor, Malaysia",  
Child: Care, Health and Development, 2016

Publication

<1 %

47

[epdf.tips](http://epdf.tips)

Internet Source

<1 %

48

[eprints.ums.edu.my](http://eprints.ums.edu.my)

Internet Source

<1 %

49

[etheses.whiterose.ac.uk](http://etheses.whiterose.ac.uk)

Internet Source

<1 %

50

[htsm2019.gaics.org](http://htsm2019.gaics.org)

Internet Source

<1 %

51

[industrialpsychiatry.org](http://industrialpsychiatry.org)

Internet Source

<1 %

52

[ir-library.ku.ac.ke](http://ir-library.ku.ac.ke)

Internet Source

<1 %

53

[jurcon.ums.edu.my](http://jurcon.ums.edu.my)

Internet Source

<1 %

54

[mdpi-res.com](http://mdpi-res.com)

Internet Source

<1 %

55

[mobt3ath.com](http://mobt3ath.com)

Internet Source

<1 %

56

[ugspace.ug.edu.gh](http://ugspace.ug.edu.gh)

Internet Source

<1 %

57

[www.academia.edu](http://www.academia.edu)

Internet Source

<1 %

58

[www.cambridge.org](http://www.cambridge.org)

Internet Source

<1 %

59

[www.vlgafc.ru](http://www.vlgafc.ru)

Internet Source

<1 %

60

Abdullah Khamaiseh, Mohammed  
ALBashtawy. "Oral health knowledge,  
attitudes, and practices among secondary

<1 %

school students", British Journal of School  
Nursing, 2013

Publication

---

61

Sharifah Nur Umairah, Binti Tuan Yahya,  
Madya Datin, Safiah Md. Yusof. "Relationship  
Between Dietary Pattern and Body Mass  
Index Among Primary School Children", Asian  
Journal of Clinical Nutrition, 2012

Publication

---

<1 %

62

Wai Yang, Tracy Burrows, Lesley MacDonald-  
Wicks, Lauren Williams, Clare Collins, Winnie  
Chee, Kim Colyvas. "Body Weight Status and  
Dietary Intakes of Urban Malay Primary  
School Children: Evidence from the Family  
Diet Study", Children, 2017

Publication

---

<1 %

63

Yi-Shin Lee, John Komar, Michael Yong Hwa  
Chia. "Physical Activity Measurement  
Methodologies: A Systematic Review in the  
Association of South East Asian Nations  
(ASEAN)", Sports, 2021

Publication

---

<1 %

64

Dorottya Kisfalusi. "Interethnic Relations  
among Roma and Non-Roma Students in  
Hungary", Corvinus University of Budapest,  
2016

Publication

---

<1 %



---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On