

Subsurface Structure Determination of Geothermal Area in Siogung-ogung Samosir District by Using Magnetic Method

by Juniar Hutahaeen

Submission date: 11-Nov-2022 02:23PM (UTC+0700)

Submission ID: 1950972167

File name: 3_a._Subsurface_Structure_Determination_ICIESC_IOP_2018.pdf (804.65K)

Word count: 2448

Character count: 13326

PAPER • OPEN ACCESS

Subsurface Structure Determination of Geothermal Area in Siogung-ogung Samosir District by Using Magnetic Method

To cite this article: Togi Tampubolon *et al* 2018 *J. Phys.: Conf. Ser.* **970** 012002

View the [article online](#) for updates and enhancements.

You may also like

- 14 - [causing factors of declining trend on](#)
- 17 - [samosir panorusan goat population as endangered species in Samosir Island](#) Nurzainah Ginting, A Manik, Hasnudi *et al.*
- 6 - [Magnetic anomaly inversion using magnetic dipole reconstruction based on the pipeline section segmentation method](#) Qi Pan, De-Jun Liu, Zhi-Yong Guo *et al.*
- 6 - [A brief review of magnetic anomaly detection](#) Yue Zhao, Junhai Zhang, Jiahui Li *et al.*



*Benefit from connecting
with your community*

ECS Membership = Connection

ECS membership connects you to the electrochemical community:

- Facilitate your research and discovery through ECS meetings which convene scientists from around the world;
- Access professional support through your lifetime career;
- Open up mentorship opportunities across the stages of your career;
- Build relationships that nurture partnership, teamwork—and success!

Join ECS!

Visit electrochem.org/join



Subsurface Structure Determination of Geothermal Area in Siogung-ogung Samosir District by Using Magnetic Method

Togi Tampubolon¹, Juniar Hutahaean¹, Suryani N.J. Siregar¹

Department of Physics, State University of Medan, Indonesia

Corresponding author: topartam@gmail.com

Abstract. Underwater research often uses geomagnets. It is one of the geophysical methods for measuring magnetic field variations. This research was done to identify how the subsurface rock structure is and determine kinds of rock based on its susceptibility value in Siogung-ogung geothermal area, Pangururan, Samosir District. The tool measurement of total magnetic field called Proton Precision Magnetometer, positioning using Global Position System, and north axis determination using geological compass. Data collection was done randomly with 51 measuring points obtained. Data analysis started with International geomagnetics Reference Field correction to obtain the total magnetic field anomaly. Then, the data analysis of total magnetic anomaly was done by using surfer program 12. To get a magnetic anomaly cross section used Magdc For Windows program. Magnetic measurement results indicated that the variation of magnetic field strength in each point with the lowest magnetic intensity value of 41785.67 nano tesla. The highest magnetic intensity value is 43140,33. From the results of qualitative interpretation, the magnetic anomaly value is at -200.92 to 1154.45 whereas the quantitative interpretive results of model show the existence of degradation and andesitic rocks, with the value of susceptibility

1. Introduction

The Indonesian archipelago locates near with the boundary of the Eurasian and Indo-Australian tectonic plates. The type of boundary between these two plates is convergent. Indo-Australian Plate is a plate that dips under the Eurasian plate (subduction). In addition, to the east, it met 3 tectonic plates at once, namely the Philippines, Pacific, and Indo-Australian plate [1]. Subduction between two plates causes the formation of volcanoes and oceanic troughs. Similarly, the subduction between the Indo-Australian Plate and the Eurasian Plate causes the formation of volcanoes in rows, that is Bukit Barisan on Sumatra Island and volcanoes along Java island, Bali and Lombok, and the ocean trench, that is Java Trough (Sunda). Tectonic plates continue to move on each other, their movements causing friction with each other to produce big energy that causes magma rise to the surface and volcanic activity increased or the emergence of geothermal sources on the surface. Therefore, Indonesia is in the fire rings area that have enormous geothermal potential.

Population development in Indonesia is accompanied by the increasing of energy demand. So far, energy supplies still utilize energy sources that dominated by fossils such as petroleum, gas and coal. Dependence on fossil energy causes these energy reserves to be depleted, it is necessary to find alternative renewable energy sources. Renewable alternative energy sources that are very potential and it is time to be developed in Indonesia is geothermal. One of the undeveloped geothermal energy



sources is geothermal Siogung-ogung, Pangururan District, Samosir District. So far, Geothermal Siogung-ogung is only used as a hot spring water for tourism object [2].

By looking at the potential of this geothermal, it is necessary to conduct comprehensively research so that geothermal energy resources can be utilized better. In this research, the method for identification of subsurface structure is magnetic method. The application of magnetic methods to geothermal exploration is based on the demagnetization of a material due to temperature. The magnetism of a material will decrease as the temperature rises and even disappears when it reaches the Curie temperature. The observation of the magnetic field is based on measuring the magnetic field strength values. Variations in magnetic field strength values are caused by differences of existing rock susceptibility value at the measurement site as an internal factor and the influence of celestial bodies, especially solar activity as an external factor.

2. Research Method

2.1. Measurement

The position of each measuring point is determined by Garmin Extrex GPS including latitude, longitude, and elevation. The strength measurement of the total magnetic field was carried out using two tools, they are Proton Precision Magnetometer (PPM) G-856 AX model at base point and PPM elsec 770 type at 51 point stations. Measurements at the base point are periodical at certain intervals, while the measurements at each station point are random and repeatable for three times.

2.2. Correction

The magnitude of the measured magnetic field at each station point is essentially the contribution of the earth's main magnetic field, the outer magnetic field, and the anomalous magnetic field. To obtain the anomalous magnetic field strength that produced by the rock beneath of each measurement point surface, it is necessary to make correction to the main magnetic field and external magnetic field which is also measured through PPM. The corrections which are done include:

1. Daily correction (diurnal correction) is a correction to remove the influence of external magnetic field mainly from solar activity. Daily correction is obtained from a variation in magnetic field strength over a specified time interval at the base point corresponding to the measuring time at each station point. If there is a decrease in magnetic field strength, it must be added for its decrease to the measurement result at the station point, and vice versa.
2. Topographic correction is a correction to remove magnetic field magnetic field influence due to topographic differences. There is no general rule in this correction, but if it is deemed as unmagnetized, the correction made only refers to the elevation difference, that is 0.015 nT/m for the equatorial area.
- 3) IGRF Correction is a correction to eliminate the influence of the main magnetic field or International Geomagnetic Reference Field (IGRF). The price of IGRF is certain for each point (Latitude, Arc, and elevation) on the earth's surface [3].

The final result of the magnetic method is the strength of the anomalous magnetic field after correction with the equation:

$$\Delta H = H_{OBS} \pm \Delta H_{vh} \pm H_T - H_{IGRF} \quad (1)$$

with:

ΔH	= Magnetic Anomaly
H_{OBS}	= Measured magnetic field price
ΔH_{vh}	= Daily variation price
H_T	= Topography correction price
H_{IGRF}	= IGRF correction price

Suceptibility is the degree of the magnetization of an object due to the influence of the magnetic field. To estimate the magnetic susceptibility value at the measurement point, it can be determined from the equation:

$$K = \frac{\Delta H}{H} \quad (2)$$

with: K = Susceptibility of the magnetic field at the point of measurement
 ΔH = Magnetic field Anomaly strength at measurement point (nT)
 H = Average magnetic field strength in the measurement area (nT)

The value of susceptibility is very important for anomalous objects searching, because of its very characteristic for each type of mineral or metal mineral.

2.3. Interpretation

Interpretation of geomagnetic data is done both qualitatively and quantitatively. Qualitative interpretation is done by presenting the magnetic field strength of anomaly in two-dimensional map using Surfer 12. The contour pattern of anomalous magnetic field shows the distribution of geomagnetic objects or geological structures at Earth subsurface. Furthermore, the magnetic field anomaly pattern is interpreted based on local geological information in the form of the distribution of magnetic objects or geological structures, as the basic for estimating the actual geological condition. Quantitative interpretation aims to describe the subsurface structure to determine the lithology of the research area [5]. Interpretation is done to create a geomagnetic sectional model using Mag2dc software. The geomagnetic sectional model is linked to a table of susceptibility values in rock type and geological data of the research area to determine the subsurface rock structure.

3. Results

3.1. Distribution Patterns of Magnetic Anomalies

The distribution pattern of anomalous magnetic field strength in the measurement area is shown in Figure 1 below:




Figure 1. Distribution Patterns of Siogung-ogung Magnetic Anomalies

Based on the picture above, it is found that the highest magnetic field strength is 1154.45 (nT) at A27 point and the lowest anomaly is -200,92 (nT) at A1 point. It is accordance with the survey area where the area is a magnetic measurement point that close to the emergence of hot water. The price of low and medium magnetic anomalies is spread over the west which is closely related to the formation of geothermal manifestations in this area. This strong magnetic field states the effect of rocks

demineralization due to high temperatures. This area is interpreted as a geothermal prospect area that needs to be examined more thoroughly to ensure its energy potential [6].

3.2. Susceptibility of Siogung-ogung Geothermal Rock

Based on the magnetic field strength of anomaly, then the susceptibility calculation is using equation 2). The pattern of anomalous magnetic field strength distribution in the measurement area is shown in Figure 2 below:




Figure 2. Pattern of Susceptibility Distribution at Siogung-ogung

From the calculation of susceptibility value, it is obtained that Siogung-ogung geothermal region has susceptibility value from the lowest to highest that is: $-0,00479 \times 10^3$ until $0,0275 \times 10^3$. The susceptibility value obtained will be used to determine the type of rock at subsurface of Siogung-ogung geothermal area through quantitative analysis with modeling using Mag2dc software.

3.3. Magnetic Anomaly Modeling

A quantitative interpretation is made to create a geomagnetic sectional model using Mag2dc software for susceptibility data. Modeling is done by making the line section (AA' path) from the lowest anomaly to the highest that suspected as a source of magnetic anomalies. In numerical modeling, some geometric magnetic field parameters are needed in the research area including IGRF value (41987.1 nT), declination angle (-0.1210 °), inclination angle (-0.0164 °) and other parameters. The modeling results are shown in Figure 3 below:




Figure 3. Sectional Model of Siogung-ogung Geomagnetic Using Mag2dc

Figure 3 shows the AA' cross-sectional model, where the x-axis denotes the length of the latitude, the positive y-axis represents the strength value of magnetic anomaly, field and the negative y-axis

represents depth. From the modeling results, it can be determined rocks lithology based on the susceptibility value. The top layer with a value of $k = -0.0254$ is interpreted as pyroclastic rock at a depth of 0 to 20 m which serves as cup rock on a geothermal system. Layers with values of $k = 0.0499$ are interpreted as andesite rocks at depths of 4 to 25 m which serve as a barrier to the loss of geothermal steam. The layer with the value of $k = 0.0877$ at a depth of 10 to 25 m, $k = 0.1734$ at a depth of 10 to 35 m, $k = 0.0356$ at a depth of 41 to 60 m, and $k = 0.0579$ at a depth of 60 to 100 m, interpreted as a strong alternated rock or have undergone many changes and decreased magnetism properties due to heat. The base of this layer is a reservoir zone and beneath this layer is the geothermal host rock and geothermal vapor.

The contrast of negative and positive susceptibility values to area A is interpreted as a Horst Graben structure, a fracture used as a geothermal vapor exit and in that area there is a fumarole crater which is a geothermal manifestation. If there it is related with the value of the susceptibility table by Telford [8], the rock is andesite that have a susceptibility of 0.0100 to 0.0500 in line with the geological data of the research area. This result is in line with the previously obtained that andesite rocks have a susceptibility between 0.0200 to 0.0400 [9].

4. Conclusion

From the research results, it can be concluded, such as (1) The magnetic anomaly value in the survey area ranges from - 200.92 nT at coordinates 237039 N 9840345 E to 1154.45 nT at coordinates 237063 N 9840356 E and (2) Based on the value of susceptibility obtained, the types of rocks found in this research area are desiccation and andesitic rocks with the value of susceptibility (0.0254, 0.0499, 0.0877, 0.1734, 0.0356, 0.00579)

References

- [1] Badan Geologi, (2010), *Hidup di atas tiga lempeng : Energy, Mineral dan Geodiversity*, Badan Geologi, Kementerian Energy dan Sumber Daya Mineral, Bandung.
- [2] Badan Pusat Statistik, (2015), *Statistika Daerah Kabupaten Simalungun 2015*, Badan Pusat Statistika Kabupaten Samosir, Medan : Samosirkab.bps.go.id (diakses pada 09 Oktober 2016).
- [3] Broto, S., T. T. Putranto, (2011), *Aplikasi Metode Geomagnet Dalam Eksplorasi Panas Bumi*, UNDIP: Semarang.
- [4] <http://swdcwww.kugi.kyoto-u.ac.jp/igrf/point/index.html>, model field at a point by IGRF accessed, August, 23, 2016.
- [5] Winarsih, F. P., (2014), *Identifikasi Litologi daerah Manifestasi Panas Bumi Parangwedang Kabupaten Bantul DIY dengan Metode Magnetik*, Skripsi Fakultas Sains dan Teknologi, UIN Sunan Kalijaga, Yogyakarta.
- [6] Singarimbun, A., Cyrke, A.D.N.B, and Riva, C.F., (2013). *Penentuan Struktur Bawah Permukaan Area Panas Bumi Patuha dengan Menggunakan Metode Magnetik*, *Jurnal Matematika dan Sains*, Agustus 2013, Vol. 18 No.2. Sulawesi Utara.
- [7] Setiawan, Sigit, (2012). *Energi Panas Bumi Dalam Kerangka MP3EI : Analisis Terhadap Prospek, Kendala, dan Dukungan Kebijakan*, *Jurnal Ekonomid dan Pembangunan*, Vol. XX (1).
- [8] Telford, W. M., L. P. Geldart dan R. E. Sheriff., (1991), *Applied Geophysics*, Second Edition. Cambridge University Press: USA.
- [9] Lita, F., (2012), *Identifikasi Anomali Magnetik di Daerah Prospek Panasbumi Arjuna- Welirang*, Skripsi, FMIPA, Universitas Indonesia (UI), Depok.

Subsurface Structure Determination of Geothermal Area in Siogung-ogung Samosir District by Using Magnetic Method

ORIGINALITY REPORT

22%

SIMILARITY INDEX

18%

INTERNET SOURCES

16%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | Submitted to UC, Boulder
Student Paper | 4% |
| 2 | Sehah, M A Kurniawan, I Andriyanto, B Arismawan, A Risyad, A M Wibowo, S A Raharjo, Sugito. "Estimation of Potential of Iron Sand in The Eastern Coastal Area of Cilacap Regency Based on The Local Magnetic Anomalies Data", Journal of Physics: Conference Series, 2020
Publication | 2% |
| 3 | www.repository.cam.ac.uk
Internet Source | 2% |
| 4 | factsofindonesia.com
Internet Source | 2% |
| 5 | Lestari Sinaga Domini, Togi Tampubolon. "Temperature Analysis Of Tinggi Raja Geothermal Reservoir, Simalungun District", Journal of Physics: Conference Series, 2020
Publication | 1% |

6	<p>N.E. Parcutela, C.B. Dimalanta, L.T. Armada, R.S. Austria, J.A. Gabo-Ratio, G.P. Yumul. "Band processing of Landsat 8-OLI multi-spectral images as a tool for delineating alteration zones associated with porphyry prospects: A case from Suyoc, Benguet, Philippines", IOP Conference Series: Earth and Environmental Science, 2022</p> <p>Publication</p>	1 %
7	<p>jurnal.untan.ac.id</p> <p>Internet Source</p>	1 %
8	<p>"International Conference on Innovation in Education, Science and Culture (ICIESC-2017)", Journal of Physics: Conference Series, 2018</p> <p>Publication</p>	1 %
9	<p>mafiadoc.com</p> <p>Internet Source</p>	1 %
10	<p>idoc.pub</p> <p>Internet Source</p>	1 %
11	<p>media.neliti.com</p> <p>Internet Source</p>	1 %
12	<p>U. R. Christensen. "Scaling properties of convection-driven dynamos in rotating spherical shells and application to planetary magnetic fields", Geophysical Journal International, 7/2006</p> <p>Publication</p>	1 %

13	Zimmerman, S.. "Rsp1p, a J Domain Protein Required for Disassembly and Assembly of Microtubule Organizing Centers during the Fission Yeast Cell Cycle", <i>Developmental Cell</i> , 200404 Publication	1 %
14	iopscience.iop.org Internet Source	1 %
15	lib.unnes.ac.id Internet Source	1 %
16	www.slideshare.net Internet Source	1 %
17	A Y Fitrianna, C Novtiar. "The role of education, values, culture and character in online learning for learning mathematical outcomes of prospective teachers", <i>Journal of Physics: Conference Series</i> , 2020 Publication	<1 %
18	ejournal.uksw.edu Internet Source	<1 %
19	etheses.uin-malang.ac.id Internet Source	<1 %
20	Togi Tampubolon, Rita Juliani, Juniar Hutahean, Jeddah Yanti. "Retrieving Surface Soil Moisture from Optical Satellite Imagery"	<1 %

over Medan and Surrounding", Journal of Physics: Conference Series, 2020

Publication

21

sinta.ristekbrin.go.id

Internet Source

<1 %

22

www.hindawi.com

Internet Source

<1 %

23

www.scribd.com

Internet Source

<1 %

24

digilib.uin-suka.ac.id

Internet Source

<1 %

25

ojs.uho.ac.id

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off