

Activity Test of CLA Synthesized from Castor Oil by in Vivo White Mice (*Rattus norvegicus*)

by Tita Juwitaningsih

THE
Character Building
UNIVERSITY

Submission date: 18-May-2023 09:00AM (UTC+0700)

Submission ID: 2095878004

File name: zed_from_Castor_Oil_by_in_Vivo_White_Mice_Rattus_norvegicus.pdf (377.62K)

Word count: 2184

Character count: 11492

PAPER • OPEN ACCESS

Activity Test of CLA Synthesized from Castor Oil by in Vivo White Mice (*Rattus norvegicus*)

To cite this article: Marham Sitorus *et al* 2020 *J. Phys.: Conf. Ser.* **1485** 012062

View the [article online](#) for updates and enhancements.

You may also like

- 2 [Evaluation of the Morpho-physiology characteristics of maize inbred lines introduced from CIMMYT to identify the best candidates for planting in acidic soil in Jasinga, Indonesia](#)
K Lubis, S H Sutjahjo, M Syukur et al.
- 4 [Identification of feeding pattern and their impact on milk fatty acid profiles from traditional dairy cows in Pangalengan Sub-district](#)
D Anzhany, Despal, T Tohamat et al.
- 5 [Preliminary observations on the effects of milk fortification with conjugated linoleic acid in yogurt preparation](#)
R V Salamon, I Albert, C D Andrés et al.

ECS The Electrochemical Society
Advancing the world state of electrochemical science & technology

242nd ECS Meeting
Oct 9 - 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD

Submit your abstract

Activity Test of CLA Synthesized from Castor Oil by in Vivo White Mice (*Rattus norvegicus*)

Marham Sitorus^{1,*}, Manihar Situmorang¹, Tita Juwitaningsih¹, Minat Karya Nia Hia¹ and Rini F Hutabarat¹

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Negeri Medan, Medan 20221, Indonesia

*Corresponding author: marham@unimed.ac.id

Abstract. Malondialdehyde (MDA) is a metabolite resulting from lipid peroxidation by free radicals that can be formed when hydroxyl free radicals such as Reactive Oxygen Species (ROS) react with fatty acid components of the cell membrane so that a chain reaction is known as fat peroxidation. The fat peroxidation will break the chain of fatty acids into various toxic compounds and cause damage to the cell membrane. Thus MDA is an parameter of the presence of free radicals in the body. In this research, the Conjugated Linoleic Acid (CLA) antioxidant test was synthesized from castor oil in vivo against white mice (*Rattus norvegicus*). White mice exposed to free radicals through cigarette smoke for 2 hours per day for 14 days. Mice were given an antioxidant intake of CLA with a concentration of 200, 400, 600, 800, 1000 μg / body weight every day with three repetitions. The same thing is done with intake of Ascorbic acid (Vitamin C) and α -tocopherol (Vitamin E) as a comparison. As a control also made negative control without treatment and positive control by exposure to cigarette smoke without intake of antioxidants. The CLA can inhibit free radicals by reduction MDA in blood an inhibitory equivalent to Ascorbic acid (Vitamin C) and α -tocopherol (Vitamin E).

1. Introduction

Antioxidants are compounds that can inhibit the rate of oxidation or prevent the reaction of free radical formation by neutralizing free radicals in cells. Chemically free radicals cause cell damage and will cause cancer and tumors, which today is one type of disease that causes high mortality [1]. Compounds that are widely used as antioxidants are α -tocopherol or vitamin E and ascorbic acid or vitamin C, which are abundant in nature, whereas when synthesized requires high costs.

The urgency of this research is the study of alternative antioxidants that are effective, inexpensive, abundant and renewable. Conjugated Linoleic Acid (CLA) which is found in large livestock (ruminant), milk and its by-products are also antioxidants, but the abundance is very limited. Synthetically, CLA can be obtained from ricinoleate, which is a major component of castor oil. Castor oil is obtained from castor bean seeds (*Ricinus communis* Linn) with oil content of 40-50% and ricinoleic content of 80-90%, castor oil is one of the oil that is not consumed (non edible oil) and this is an advantage because it does not compete with edible oil [2]. In vitro the antioxidant activity of Conjugated Linoleic Acid (CLA) synthesized from castor oil has been tested against $4 \mu\text{g}$ / mL α , α -diphenyl-2-picrylhydrazyl (DPPH) with ethanol as a spectrophotometer^[3]. The results show that at



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

concentrations starting at 4 $\mu\text{g} / \text{mL}$, it can inhibit more than 80% DPPH equivalent to the inhibitory power of α -tocopherol and ascorbic acid which is a commonly used natural antioxidant. The IC50 (inhibitory concentration of 50%) of CLA measured at 8 $\mu\text{g} / \text{mL}$ DPPH is 2,133.4 $\mu\text{g} / \text{mL}$ equivalent to α -tocopherol 2,839.4 $\mu\text{g} / \text{mL}$ and ascorbic acid 2,947.4 $\mu\text{g} / \text{mL}$. The very high antioxidant activity of CLA and the very strong IC50 category (<50 $\mu\text{g} / \text{mL}$) against DPPH indicate that CLA has the potential to be used as an antidote [3].

Furthermore, in vivo, the antioxidant activity of CLA has been tested on experimental animals which are conditioned to be exposed to free radicals through cigarette smoke by measuring blood Malondyaldehyde (MDA) levels, spectrometers appear as parameters that animals are exposed to free radicals. The presence of MDA in the blood is the result of free radical lipid peroxidation as an indicator of the presence of free radicals in the body of experimental animals [4, 5, 6, 7].

2. Materials and Methods

2.1. Chemicals and instruments

Chemicals materials used are qualification Brands namely: Tiobarbituric acid (TBA), Trichloroacetic (TBA), Acetic acid glacial, 11,33, Tetra methoxy propane (TMP), Aquadest and white mice (*Rattus norvegicus*). The research instrument are, EDTA tube (vortex), smoking chamber, centrifuge and UV – Vis SIMADZU Spectrophotometer.

2.2. Activity test of CLA synthesized from castor oil by in vivo white mice (*Rattus norvegicus*)

The experimental animals (*Rattus norvegicus*) used were 2.5 - 3.0 months old, body weight 100 - 200 g, active, not deformed and male. Animal experiments were divided into 7 treatment groups randomly. Each treatment group consisted of 3 replications for 3 types of antioxidants (CLA, Vitamin C and D), so that the total number of mice used was 65 animals including positive and negative controls. The cigarettes used are clove cigarettes. Free radical exposure was carried out using 3 cigarettes for each treatment group, except the negative control group. The exposure was carried out for 14 days in a smoking chamber. One hour after exposure to cigarette smoke continued with given of CLA, Vitamin C and Vitamin D according to the dose in each treatment group. Furthermore, blood sampling was performed on rats that were fasted for 24 hours. Mice were anesthetized using chloroform then dissected and their blood drawn through the heart organ using a 3 mL syringe. The blood obtained is collected in a blood tube and centrifuged for 10 minutes at 3000 rpm. Blood plasma located at the top is separated and taken to be analyzed for MDA concentration with Visible spectrophotometer at wavelength ($\lambda = 535 \text{ nm}$).

3. Results and Discussion

The initial stage carried out in this in vivo test was the conditioning of mice in the experimental cage before treatment. After conditioning the exposure of cigarette smoke with a duration of 2 hours / day for 14 days and continued with the treatment of intake of CLA, ascorbic acid (vitamin C) and α -tocopherol (vitamin E) as a comparison and ascorbic acid with a variation of concentration of 200, 400, 600, 800 and 10 mg / kg / day with three repetitions, respectively. Besides that, treatment was also carried out for positive control (fumigation without antioxidant intake) and negative control (without fumigation and without antioxidants). After completion of the treatment, blood from the heart of the experimental animal is then surgically collected in an EDTA tube to prevent clotting and then centrifuged for blood serum collection [8, 9].

Table 1. Absorbance data and antioxidant inhibition of CLA, ascorbic acid (vitamin C) and α -tocopherol (vitamin E).

Dosage (mg/kgWeight)	CLA		Asam Ascotbat (Vit C)		A-tokoferol (Vit D)	
	A_{av}	%inhibition ^{*)}	A_{av}	% inhibition ^{*)}	A_{av}	% inhibition ^{*)}
200	0,0493	3,52	0,0413	19,18	0,0477	6,65
400	0,0482	5,68	0,0468	8,41	0,0474	6,85
600	0,0474	7,63	0,0481	5,87	0,0477	6,65
800	0,0478	6,46	0,0483	5,48	0,0479	6,26
100	0,0447	12,52	0,0465	9,01	0,0446	12,72

$$A_{\text{kontrol(-)}} = 0,0397$$

$$A_{\text{kontrol(+)}} - A_{\text{kontrol(-)}} = 0,0114 \text{ (MDA in the body)}$$

$$*) \% \text{ inhibition} = [(A_{\text{kontrol(+)}}) - (A_{\text{antooksidan}})] / (A_{\text{kontrol (+)}})$$

Furthermore, measured levels of Malondyaldehyde (MDA) blood serum, by spectrometer appear. Malondialdehyde (MDA) is a metabolite resulting from lipid peroxidation by free radicals that can be formed when hydroxyl free radicals such as Reactive Oxygen Species (ROS) react with fatty acid components of the cell membrane so that a chain reaction is known as fat peroxidation. The fat peroxidation will break the chain of fatty acids into various free radicals that are toxic and cause damage to the cell membrane. Thus MDA is an indicator of the presence of free radicals in the body with the results in Table 1 and the graph in Figure 1.




Figure 1. Graph of the inhibitory power of antioxidant CLA, ascorbic acid (vitamin C) and α -tocopherol (vitamin E) at concentration (mg/ kg weight).

Based on table 1 and figure 1, the CLA can inhibit free radicals and the relative pattern is the same as the inhibitory power of ascorbic acid (Vitamin C) and α -tocopherol (Vitamin E). Thus the CLA has the potential to be used as an inexpensive and abundant natural antioxidant. In addition it is also a renewable antioxidant, through the cultivation of Ricinus tree as a source of castor oil. Many plants are easy to grow because they do not need strict soil requirements and can grow on marginal land. The contribution will raise the economic value of the castor plant in general and castor oil in particular.

4. Conclusions

As in vitro CLA can inhibit DPPH with IC_{50} (inhibitory concentration of 50%) 2,133 4 $\mu\text{g} / \text{mL}$ equivalent to α -tocopherol 2,839 4 $\mu\text{g} / \text{mL}$ and ascorbic acid 2,947 4 $\mu\text{g} / \text{mL}$. The very high antioxidant activity of CLA and the very strong IC_{50} category ($<50 \mu\text{g} / \text{mL}$) against DPPH indicate that CLA has the potential to be used as an antioxidant. Furthermore, the antioxidant activity in vivo to experimental animals conditioned by free radicals can inhibit free radicals measured by decreasing blood Malondyaldehyde (MDA) by visible spectrometers as parameters that animals are exposed to free radicals.

References

- [1] Yu L, Adams D and Gabel M. 2002 Conjugated linoleic acid isomers differ in their free radical scavenging properties *Journal of Agricultural and Food Chemistry* **50** 4135-4140.
- [2] Patrick L. 2006 Lead toxicity part II: the role of free radical damage and the use of antioxidants in the pathology and treatment of lead toxicity *Alternative Medicine Review* **11** 114-127.
- [3] Bhowimic D N and Sarma A N. 1987 Dehydration of castor oil. Ind. Eng. Chem. Prod; Departement of Oil and Paint Technology Kampur India.
- [4] Muniz P, Coma M J and Teran J. 2014 Oxidative stress and vascular damage in hypoxia processes. Malondialdehyde (MDA) as biomarker for oxidative damage *Electron J Biomed* **2** 50-53.
- [5] Sinaga F A. 2017 Stress oksidatif dan status antioksidan pada aktivitas fisik maksimal *Generasi Kampus* **9** 176-189
- [6] Sas K, Robotka H, Toldi J and Vécsei L. 2007 Mitochondria, metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders *Journal of the Neurological Sciences* **257** 221-239.
- [7] Raharja K T Wirjatmadi B and Adriani M. 2016 Pemberian buah kawista menghambat peningkatan kadar malondialdehid serum tikus wistar yang dipapar asap rokok *Jurnal Kedokteran Brawijaya*, **29** 196-201.
- [8] Pryor WA. 1997 Cigarette smoke radicals and the role of free radicals in chemical carcinogenicity *Environmental health perspectives* **105** 875-882.
- [9] Tyndale R F and Sellers E M. 2001 Variable CYP2A6-mediated nicotine metabolism alters smoking behavior and risk *Drug Metabolism and Disposition* **29** 548-552.
- [10] Simorangkir M, Hutabarat W, Nainggolan B and Silaban S. 2019 Antioxidant and antibacterial activities of nonpolar to polar solvent extracts of Sarang Banua (*Clerodenum fragrans* Vent Willd) leaves *Rasayan Journal of Chemistry* **12** 959-965.

Acknowledgement

Research grant from Direktorat General of the Ministry of Research, Technology and Higher Education who funded this research through the Grant of Centralization of Competency-Based Financial Year 2019 is gratefully acknowledged. Thanks also to Dr. Tutik Dwi Wahyuningsih, M.Sc Head of the Organic Chemistry Laboratory and Instrument Gajah Mada University who recorded, GC-MS and FTIR and Drs. Marudut Sinaga, M.Si, Head of the Unimed Chemistry Laboratory which records the antioxidant activity by UV Spectrophotometer. Thank you also disgiused to Minat Karya Nia Hia and Rini F Hutabarat students who were involved in this research in order to completed her Thesis.

Activity Test of CLA Synthesized from Castor Oil by in Vivo White Mice (*Rattus norvegicus*)

ORIGINALITY REPORT

12%

SIMILARITY INDEX

12%

INTERNET SOURCES

1%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Wolverhampton Student Paper	3%
2	toc.proceedings.com Internet Source	2%
3	www.bezpecnostpotravin.cz Internet Source	2%
4	animres.edpsciences.org Internet Source	1%
5	china.iopscience.iop.org Internet Source	1%
6	digilib.unimed.ac.id Internet Source	1%
7	vital.seals.ac.za:8080 Internet Source	1%
8	www.mdpi.com Internet Source	1%
9	adoc.tips Internet Source	<1%

10

worldwidescience.org

Internet Source

<1 %

11

eureka.patsnap.com

Internet Source

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On



THE
Character Building
 UNIVERSITY