

DAFTAR PUSTAKA

- Anggraini, T. (2017). *Proses dan manfaat teh*. Padang: CV. Rumahkayu Pustaka Utama.
- Anggraini, T., Neswati, Nanda, R. F., & Syukri, D. (2021). Effect of Processing on Green and Black Tea DPPH Radical Scavenging Activity, IC₅₀ Value, Total Polyphenols, Catechin and Epigallocatechin Gallate content. 709(1), 012017. <https://doi.org/10.1088/1755-1315/709/1/012017>
- Anis, U., Nurbawita, A., Marniza, M., Marsigit, W., & Gusriani, I. (2023). Pengaruh ketinggian tempat tumbuh terhadap kadar air, kadar abu, total fenol, dan aktivitas antioksidan daun teh. *Jurnal Industri Hasil Perkebunan*, 18(2), 24. <https://doi.org/10.33104/jihp.v18i2.8217>
- Anoto, D. A., Basuki, S. P. H., & Setiyabudi, R. (2024). Efektifitas Pemberian Konsumsi Teh Hijau terhadap Penurunan Gula Darah Diabetes Melitus Tipe 2 (DMT2)”. *Jurnal Ilmiah Permas: Jurnal Ilmiah STIKES Kendal*, 14(2), 479-488.
- Astill, C., Birch, M. R., Dacombe, C., Humphrey, P. G., & Martin, P. T. (2001). Factors affecting the caffeine and polyphenol contents of black and green tea infusions. *Journal of Agricultural and Food Chemistry*, 49(11), 5340-5347. <https://doi.org/10.1021/jf010759+>
- Astuti, R., Sulastri, D., & Setyaningsih, R. (2020). Standardization of Brewing Methods for Indonesian White Tea. *Jurnal Pangan dan Gizi*, 15(1), 30–36.
- Badan Standardisasi Nasional. (2011). SNI 3143:2011 Teh Hitam. Jakarta: Badan Standardisasi Nasional.
- Badan Standardisasi Nasional. (2013). SNI 3836:2013 Teh Kering dalam Kemasan. Jakarta: Badan Standardisasi Nasional.
- Blois, M. S. (1958). Antioxidant determinations by the use of a stable free radical. *Nature*, 181(4617), 1199-1200.
- BPS. (2023). Statistik Perkebunan Teh Indonesia Tahun 2022.
- Candioti, L. V., De Zan, M. M., Cámara, M. S., & Goicoechea, H. C. (2014). Experimental design and multiple response optimization using the

- desirability function in analytical methods development. *Talanta*, 124, 123–138.
<https://doi.org/10.1016/j.talanta.2014.01.034>
- Chadijah, S., Musdalifah, & Qaddafi, M. (2021). Optimalisasi suhu dan waktu penyeduhan daun teh hijau (*Camellia sinensis* L.) P+3 terhadap kandungan antioksidan kafein, katekin dan tanin. *Bencoolen Journal of Pharmacy*, 1(1), 59–65.
- Chen, Zhen-Yu; Zhu, Qin Yan; Tsang, David; Huang, Yu (2001). Degradation of Green Tea Catechins in Tea Drinks. *Journal of Agricultural and Food Chemistry*, 49(1), 477–482. doi:10.1021/jf000877h
- Chung S. Yang; Hong Wang; Guang Xun Li; Zhihong Yang; Fei Guan; Huanyu Jin (2011). Cancer prevention by tea: Evidence from laboratory studies. , 64(2), 113–122. doi:10.1016/j.phrs.2011.03.00
- Deng, S., Cao, Q.-Q., Gao, Y., Wu, W., Chen, J.-X., Wang, F., Zou, Q., Xu, F., Cao, X., Sun, W., Yin, J.-F., & Xu, Y.-Q. (2025). Elucidating the effect of brewing temperature on the sensory quality of Longjing tea based on multi-scale molecular sensory science. *Food Chemistry: X*, 28, 102635. <https://doi.org/10.1016/j.fochx.2025.102635>
- Derringer, G., & Suich, R. (1980). Simultaneous Optimization of Several Response Variables. *Journal of Quality Technology*, 12(4), 214–219. <https://doi.org/10.1080/00224065.1980.11980968>
- Dinas Kesehatan Provinsi Sumatera Utara. (2023). Profil Kesehatan Provinsi Sumatera Utara Tahun 2023. Medan: Dinas Kesehatan Provinsi Sumatera Utara.
- Direktorat Jenderal Perkebunan. (2022). Statistik Perkebunan Indonesia Komoditas Teh 2020–2022. Kementerian Pertanian RI.
- Effendi, D. S., Syakir, M., Yusron, M., & Hartati, R. S. (2017). *Budidaya dan pascapanen teh*. Pusat Penelitian dan Pengembangan Perkebunan, Kementerian Pertanian RI.
- FAO. (2022). FAOSTAT Statistical Database: Tea Production. Food and Agriculture Organization of the United Nations.

- Fajar, R. I., Wrasati, L. P., & Suhendra, L. (2018). Kandungan senyawa flavonoid dan aktivitas antioksidan ekstrak teh hijau pada perlakuan suhu awal dan lama penyeduhan. *Jurnal Rekayasa dan Manajemen Agroindustri*, 6(3), 196–204
- Fajrina, A., Jubahar, J., & Sabirin, S. (2016). Penetapan kadar tanin pada teh celup yang beredar di pasaran secara spektrofotometri UV-Vis. *Jurnal Farmasi Higea*, 8(2), 133–142.
- Fellows, P. J. (2009). *Food processing technology: Principles and practice* (3rd ed.). Woodhead Publishing.
- Hilal, Y. and Engelhardt, U. (2007) Characterisation of White Tea—Comparison to Green and Black Tea. *Journal für Verbraucherschutz und Lebensmittelsicherheit*, 2, 414-421.
- Ho, C.-T., Zheng, X., & Li, S. (2015). Tea aroma formation. *Food Science and Human Wellness*, 4(1), 9-27. <https://doi.org/10.1016/j.fshw.2015.04.001>
- International Organization for Standardization. (2019). ISO 3103:2019 Tea—Preparation of liquor for use in sensory tests. Geneva: International Organization for Standardization.
- Jakubczyk, Karolina; Justyna; Kochman, Joanna; Janda, Katarzyna (2020). Chemical Profile and Antioxidant Activity of the Kombucha Beverage Derived from White, Green, Black and Red Tea. *Antioxidants*, 9(5), 447–. doi:10.3390/antiox9050447
- Kodagoda, K. H. G. K., & Wickramasinghe, I. (2017). Health benefits of green and black tea: A review. *International Journal of Advanced Engineering Research and Science*, 4(7), 107–112. <https://doi.org/10.22161/ijaers.4.7.16>
- Kosińska, A., & Andlauer, W. (2014). Antioxidant capacity of tea: Effect of processing and storage. In V. Preedy (Ed.), *Tea in health and disease prevention* (pp. 109–120). Academic Press. doi:10.1016/B978-0-12-404738-9.00012-X
- Lawless, H. T., & Hemann, H. (2010). *Sensory Evaluation of Food : Principles and Practice* (2nd ed.). Springer.

- Lee, J., & Chambers, D. H. (2010). Descriptive Analysis and U.S. Consumer Acceptability of 6 Green Tea Samples from China, Japan, and Korea. *Journal of Food Science*, 75(2). <https://doi.org/10.1111/J.1750-3841.2009.01503.X>
- Leslie, P. J., & Gunawan, S. (2019). Uji fitokimia dan perbandingan efek antioksidan pada daun teh hijau, teh hitam, dan teh putih (*Camellia sinensis*) dengan metode DPPH. *Tarumanagara Medical Journal*, 1(2), 383–388.
- Leung, L. K., Su, Y.-L., Chen, R., Zhang, Z., Huang, Y., & Chen, Z.-Y. (2001). *Theaflavins* in Black Tea and Catechins in Green Tea Are Equally Effective Antioxidants. *Journal of Nutrition*, 131(9), 2248–2251. <https://doi.org/10.1093/JN/131.9.2248>
- Li, M.-Y., Liu, H.-Y., Wu, D.-T., Kenaan, A., Geng, F., Li, H.-B., Gunaratne, A., Li, H., & Gan, R.-Y. (2022). L-theanine: A unique functional amino acid in tea (*Camellia sinensis* L.) with multiple health benefits and food applications. *Frontiers in Nutrition*, 9, Article 853846
- Lipscomb, K. A. (2016). Effect of temperature on sensing intensity of basic tastes: sweet, salty and sour. *Journal of Field Robotics*, 5(4), 1. <https://doi.org/10.5539/JFR.V5N4P1>
- Liu, Y., Luo, L., Liao, C., Chen, L., Wang, J., & Zeng, L. (2018). Effects of brewing conditions on the phytochemical composition, sensory qualities and antioxidant activity of green tea infusion: A study using response surface methodology. *Food Chemistry*, 269, 24–34. <https://doi.org/10.1016/J.FOODCHEM.2018.06.130>
- Lorenz, M. (2013). The Role of Individual Tea Compounds in Cardiovascular Protective Effects of Green and Black Tea (pp. 829–840). Academic Press. <https://doi.org/10.1016/B978-0-12-384937-3.00070-7>
- Minitab Support. (2024). Response optimizer and desirability function. Minitab LLC. Retrieved from Minitab Support website.

- Molyneux, P. (2004) The Use of Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*, 26, 211-219.
- Montgomery, D. C. (2013). *Design and analysis of experiments* (8th ed.). John Wiley & Sons.
- Musial, C., Kuban-Jankowska, A., & Gorska-Ponikowska, M. (2020). Beneficial properties of green tea catechins. *International Journal of Molecular Sciences*, 21(5), 1744.
- Mutmainnah, N., Chadijah, S., & Qaddafi, M. (2018). Penentuan suhu dan waktu optimum penyeduhan batang teh hijau (*Camellia sinensis* L.) terhadap kandungan antioksidan kafein, tanin dan katekin. *Lantanida Journal*, 6(1), 1–10.
- Nugraheni, Z. V., Rachman, T. M., & Fadlan, A. (2022). *Ekstraksi Senyawa Fenolat dalam Daun Teh Hijau (Camellia Sinensis)*. *Akta Kimia Indonesia*, 7 (1), 69–76.
- Panagiotakos, D. B., Lionis, C., Zeimbekis, A., & Polychronopoulos, E. (2009). Long-term tea intake is associated with reduced prevalence of (type 2) diabetes mellitus among elderly people from Mediterranean islands: MEDIS epidemiological study. *Journal of Nutrition, Health & Aging*, 13(6), 376–381. <https://doi.org/10.1007/s12603-009-0087-0>
- Piyasena, K. G. N. P., Napagoda, M. T., Kalinga, J., & Jayasinghe, L. (2026). Health benefits of different types of tea: A review. *Food & Nutrition Health*, 3. <https://doi.org/10.1007/s44403-026-00056-1>
- Putra, I. W. E. P., Wrsiati, L. P., & Wartini, N. M. (2020). Pengaruh suhu awal dan lama penyeduhan terhadap karakteristik sensoris dan warna teh putih silver needle (*Camellia assamica*) produksi PT. Bali Cahaya Amerta. *Jurnal Rekayasa dan Manajemen Agroindustri*, 8(4), 492–501.
- Riandani, A. P. (2024). Atribut sensori dan persepsinya. Dalam M. I. F. Gunawan dkk., *Teknik evaluasi sensori produk pangan* (hlm. 13–22). CV HEI Publishing Indonesia.

- Sakanaka, S., & Ishihara, Y. (2008). Comparison of antioxidant properties of persimmon vinegar and some other commercial vinegars in radical-scavenging assays and on lipid oxidation in tuna homogenates. *Food Chemistry*, 107(2), 739–744.
- Saklar, Sena; Ertas, Erdal; Ozdemir, Ibrahim S.; Karadeniz, Bulent (2015). Effects of different brewing conditions on catechin content and sensory acceptance in Turkish green tea infusions. *Journal of Food Science and Technology*, 52(10), 6639–6646. doi:10.1007/s13197-015-1746-y
- Salamah, N., Astuti, S. F. W., Ar Rahma, N. A., & Guntarti, A. (2024). Analysis of total flavonoid content and antioxidant activity of green tea (*Camellia sinensis*) with variations in type and brewing. *Jurnal Farmasi Sains dan Praktis (JFSP)*, 10(3), 249–258.
- Sanlier, N., Atik, İ., & Atik, A. (2018). A minireview of effects of white tea consumption on diseases. *Trends in Food Science & Technology*, 82, 82–88. <https://doi.org/10.1016/j.tifs.2018.10.004>
- Sayuti, K., & Yenrina, R. (2015). *Antioksidan alami dan sintetik*. Padang: Andalas University Press.
- Sen, G., Sarkar, N., Nath, M., & Maity, S. (2020). Bioactive components of tea. 4(1), 001–009. <https://doi.org/10.29328/JOURNAL.AFNS.1001020>
- Senanayake, S. P. J. N. (2013). Green tea extract: Chemistry, antioxidant properties and food applications – A review. *Journal of Functional Foods*, 5(4), 1529–1541. <https://doi.org/10.1016/j.jff.2013.08.011>
- Setyaningsih D, *et al.* (2017). *Analisis Sensori untuk Industri Pangan dan Agro*. Bogor: IPB Press.
- Tang, Y., Yu, X., Zhang, Y., Lu, X., Liu, Q., Jia, Y., & He, C. (2020). Sensory descriptive analysis of green tea: correlation with chemical components. 461(1), 012013. <https://doi.org/10.1088/1755-1315/461/1/012013>
- Tiradritti, M., & Del Casino, C. (2022). Tea brewing parameters: Influence of temperature and time on composition and sensory properties. *Beverages*, 8(2), 18. <https://doi.org/10.3390/beverages8020018>

- Unachukwu, U.J., Ahmed, S., Kavalier, A., Lyles, J.T. and Kennelly, E.J. (2010), White and Green Teas (*Camellia sinensis* var. *sinensis*): Variation in Phenolic, Methylxanthine, and Antioxidant Profiles. *Journal of Food Science*, 75:C541-C548. <https://doi.org/10.1111/j.1750-3841.2010.01705.x>
- Winarsi, H. (2007). *Antioksidan alami dan radikal bebas*. Kanisius.
- Winiarska-Mieczan, A., & Baranowska-Wójcik, E. (2024). The Effect of Brewing Time on the Antioxidant Activity of Tea Infusions. *Applied Sciences*, 14(5), 2014. <https://doi.org/10.3390/app14052014>
- Zargar, B., Majeed, D., Ganai, S. A., Mir, S. A., Dar, B. N., & Dar, B. N. (2018). Effect of different processing parameters on antioxidant activity of tea. *Journal of Food Measurement and Characterization*, 12(1), 527–534. <https://doi.org/10.1007/S11694-017-9664-5>
- Zhang, L., Ho, C.-T., Zhou, J., Santos, J. S., Armstrong, L., & Granato, D. (2019). Chemistry and biological activities of processed *Camellia sinensis* teas: A comprehensive review. *Comprehensive Reviews in Food Science and Food Safety*, 18(5), 1474–1495. <https://doi.org/10.1111/1541-4337.12479>